




GOLETA SANITARY DISTRICT
NPDES MONITORING PROGRAM
2013 ANNUAL REPORT

Submitted: March 2014

GOVERNING BOARD

JOHN R. FOX, PRESIDENT
JOHN S. CARTER
GEORGE W. EMERSON
SHARON ROSE
JERRY D. SMITH



**GENERAL MANAGER/
DISTRICT ENGINEER**

KAMIL S. AZOURY, P.E.

A PUBLIC AGENCY
www.goletasanitary.org

March 31, 2014

California Regional Water Quality Control Board
Central Coast Region
Attn: Monitoring and Reporting Review Section
895 Aerovista Place, Suite 101
San Luis Obispo, CA 93401

Dear Mr. Harris:

Facility Name: Goleta Sanitary District

Address: One William Moffett Place
Goleta, CA 93117

Contact Person: Robert Hidalgo
Job Title: Operations Supervisor
Phone Number: (805) 967-4519

WDR/NPDES Order Number: R3-2010-0012
WDID Number: 3/420102001

Type of Report (circle one): Monthly Quarterly Semi-Annual Annual

Month(s) (circle applicable months*): JAN FEB MAR APR MAY JUN
JUL AUG SEP OCT NOV DEC

*Annual Reports (circle the first month of the reporting period)

Year: 2013

Violation(s) (Place an X by the appropriate choice): X **No** (there are no violations to report) ___ **Yes**

If Yes is marked (complete a-g):

a) Parameter(s) in Violation: _____

b) Section(s) of WDR/NPDES Violated: _____

c) Reported Value(s)

d) WDR/NPDES

Limit/Condition:

e) Dates of Violation(s)

(reference page of report/data sheet):

f) Explanation of Cause(s):

(attach additional information as needed)

g) Corrective Action(s):

(attach additional information as needed)


Comment

In accordance with the Standard Provisions and Reporting Requirements, I certify under penalty of law that this document and all attachments were prepared under my direction or supervision following a system designed to assure that qualified personnel properly gather and evaluate the information submitted. Based on my knowledge of the person(s) who manage the system, or those directly responsible for data gathering, the information submitted is, to the best of my knowledge and belief, true, accurate, and complete. I am aware that there are significant penalties for submitting false information, including the possibility of fine and imprisonment.

If you have any questions or require additional information, please contact me at the number provided above.

Sincerely,

GOLETA SANITARY DISTRICT



Name: Kamil S. Azoury, P.E.
Title: General Manager/ District Engineer

Prepared by: K. Womersley
Reviewed by: R. J. Hidalgo

GOLETA SANITARY DISTRICT


NPDES Monitoring and Reporting Program

2013 Annual Report

Quarterly and Annual Receiving Water Monitoring
Conducted by
Aquatic Bioassay and Consulting Laboratories, Inc.
29 North Olive Street
Ventura, California 93001
(805)643-5621

Submitted March 2014

I certify under penalty of law that this document and all attachments were prepared under my direction or supervision in accordance with a system designed to assure that qualified personnel properly gather and evaluate the information submitted. Based on my inquiry of the person or persons who manage the system, or those persons directly responsible for gathering the information, the information submitted is, to the best of my knowledge and belief, true, accurate, and complete. I am aware that there are significant penalties for submitting false information, including the possibility of fine and imprisonment for knowing violations.



Kamil S. Azoury, P.E.
General Manager/District Engineer
Goleta Sanitary District

Date: _____

3/31/14

AQUATIC BIOASSAY OCEANOGRAPHIC TEAM

MARINE BIOLOGISTS

T. Mikel, President

S. Johnson, Principal Scientist

D. Laur, Ph.D., Chief Biologist

Jim Mann, Biologist

Karin Wisenbaker, Biologist

MARINE CHEMISTS

R. Gossett (CRG Marine Laboratories)

TAXONOMISTS

D. Laur, Ph.D., Miscellaneous

Phillips, Crustacean Taxonomist

S. Johnson, Mollusk Taxonomist

BOAT CREW

J. Gelsinger, Captain

CHAPTER 1

INTRODUCTION

The Goleta Sanitary District (GSD) treatment plant operates under Clean Water Act Section 301(h) which waives secondary treatment requirements. On November 19, 2004 the California Regional Water Quality Control Board, Central Coast Region (RWQCB), adopted Waste Discharge Requirements (WDR) Order R3-2004-0129 and the United States Environmental Protection Agency (EPA), Region IX issued NPDES permit CA 0048150 to the Goleta Sanitary District (GSD). A settlement agreement was made a part of the NPDES 301(h) waiver permit issued in 2004. The settlement agreement required GSD to upgrade its wastewater treatment plant to full secondary treatment by November of 2014.

As required by waste discharge requirements GSD submitted an NPDES permit renewal application to the RWQCB and the EPA in May 2009. At the time of the application submittal, the District was five years into the ten year conversion schedule described in the settlement agreement of 2004. Both the RWQCB and the EPA agreed to renew the 301(h) waiver permit for another five years while GSD continued to make progress to upgrade its treatment facility. The treatment plant is operating under WDR Order No. R3-2010-0012 and NPDES Permit No. CA0048160 which became effective September 2010.

Although GSD continues to operate the wastewater treatment facility under the 301(h) waiver provision of the Clean Water Act the final full secondary tie-in of the newly built structures to the existing plant was completed on May 15 to 16, 2013. Final effluent parameters measured on a regular basis, such as suspended solids and BOD, show a dramatic decrease in concentrations from May 15th to May 16th, when the plant went from a blended effluent to a full secondary effluent.

This annual report will discuss both treatment processes, under the following section: **WASTEWATER TREATMENT PROCESS**. The first part of each section describes the process that generated a blended secondary effluent from January to May 2013 and the second part focuses on the full secondary process in operation from June to December 2013.

As a condition of the NPDES permit, GSD is required to conduct an extensive monitoring and reporting program to assess compliance with limitations established by the California Ocean Plan and the federal Clean Water Act. For GSD, these limitations have been met by blending primary and secondary treated effluent as allowed for ocean dischargers under Section 301(h) of the Clean Water Act. Under conditions set forth in the permit, GSD must monitor the influent, effluent, biosolids (sludge), the outfall and diffuser, receiving water, bottom sediment, and biology to demonstrate that the discharge of wastewater is not causing adverse impacts on the ocean environment.

The Goleta wastewater treatment plant (WWTP) is located in an unincorporated coastal area of Santa Barbara County, California. Treated wastewater is discharged to the Pacific

Ocean approximately one mile offshore of Goleta Beach County Park via a south-trending ocean outfall. The outfall lies within and extends outside of a small embayment formed by Goleta Point directly to the west.

The Goleta WWTP treats wastewater from the service areas of the Goleta Sanitary District (GSD), the Goleta West Sanitary District, the University of California at Santa Barbara, the Santa Barbara Municipal Airport, and certain Santa Barbara County facilities. Existing agreements among the agencies establish GSD as the owner of the joint wastewater treatment facilities and assign the responsibility of operation and maintenance of the facilities to GSD. However, each agency “owns” an “indeterminate, perpetual and exclusive capacity right” in the facilities and an “easement right of flow through” the facilities.

WASTEWATER TREATMENT PROCESS

The following discussion focuses on the principal features of GSD's bended secondary process of wastewater and sludge treatment. The performance capacities and characteristics of the treatment plant are detailed in Chapter 2.

Treatment Plant Facilities

The Goleta Sanitary District Wastewater Treatment Plant is located at One William Moffett Place, in an unincorporated area of Santa Barbara County, CA. The plant site is approximately 10 miles west of the City of Santa Barbara, near the Pacific Coast. A regional view of the study area is shown in Figure 1-1.

On average, over the past 10 years, 2004 to 2013, the plant has discharged about 3.9 million gallons per day (MGD) of treated effluent to the open coastal waters of the Santa Barbara Channel via an ocean outfall. The treatment plant is currently discharging municipal wastewater in accordance with NPDES permit CA 0048160. The treatment plant's discharge meets the state water quality standards as set forth in the Water Quality Control Plan for Ocean Waters of California (California Ocean plan) and the federal Clean Water Act.

Facilities Description

The Goleta wastewater treatment plant underwent its first substantial upgrade completed in June 1988. The upgraded plant was designed to assure compliance with monthly 30-day average discharge limitations of 63 mg/L for suspended solids and 98 mg/L for BOD under an average dry weather flow 9.0 MGD. The facilities operate utilizing a split-stream process of physical and biological treatment. Biological treatment is provided by the trickling filter and solids contact process. The following sections describe the treatment process.

The second substantial upgrade was completed in December 2013 almost a year before the regulatory deadline of November 2014. The process uses two trickling filters and an aeration basin to achieve full secondary treatment.

Collection System

Over 190 miles of pipelines collect wastewater that flows almost entirely by gravity to pump stations located in each agency's service area. These stations pump the flow to the treatment facility.

Pump Station and Headworks

Influent from the collection system of each agency is pumped to the treatment plant headworks where raw wastewater flows through a bar screen which removes large debris. Influent is then routed to aerated grit tanks where sand and grit are allowed to settle out. This debris and grit is then transported via truck to a local landfill. Air collected from the influent pump stations and headworks is scrubbed in odor reduction towers equipped with activated carbon.

The upgrade of 2013 also included upgrading structures that had reached the end of their useful life. The headworks upgrade included the installation of two new bar screens with a smaller screen spacing, $\frac{1}{4}$ inch in order to better remove more inorganic materials and the installation of two new screening washer/compactor units. The odor reduction tower was removed and replaced with a biological odor reduction tower.

Primary Sedimentation

Wastewater then flows into one of three circular primary sedimentation basins (primary clarifiers) where solids settling to the bottom and floatable materials rising to the surface are mechanically collected and pumped to digesters. The primary effluent flow is then split with one portion of the effluent stream receiving additional secondary treatment and the remaining primary treated flow is discharged directly after being disinfected. On average 69% of the solids were removed in the primary treatment process during 2013, with an average of 62 % from January to May and 74% from June to December.

No new structures were added to the primary treatment stage as part of the upgrade. However as part of the renovations performed under the treatment plant upgrade all three primary clarifiers were drained and inspected. Renovations included replacement of the boom sweeps, removal and replacement of the sweep motors, the catwalks were sand blasted and both the catwalks and troughs were painted. Additionally, the concrete around the effluent trough was deteriorating and this area was patched.

Secondary Treatment

Secondary treatment involves three treatment elements: the biofilter, a solids contact channel, and secondary sedimentation tanks. In the biofilter, primary effluent trickles over plastic media where bacteria feed on organic wastes, thus removing these wastes from the water. Effluent from the trickling filter flows to a solids contact channel where air is injected and the effluent is mixed with recirculated sludge from the secondary sedimentation basins. The resulting biological action coagulates these fine particles and the organic solids settle out as sludge in two secondary sedimentation tanks. Effluent from this secondary process is combined with primary effluent at the chlorine contact tank. A portion of the secondary process flow can be diverted to the reclamation facilities for tertiary treatment with gravity filters.

The upgrade included the construction of a new biofilter identical to the existing, demolition of the solids contact channel and construction of a three train aeration basin, with structures stubbed out for the construction of a fourth train in the future if needed. New construction also included a new blower building, two new secondary clarifiers and construction of various supporting structures, such as pumping stations, interstage pump, RAS station, etc.

Chlorine Contact Tank

The primary and secondary effluent flows are combined at the head of the chlorine contact chamber where sodium hypochlorite is injected to kill bacteria in the effluent. According to the District's permit, a minimum of 5 mg/L total chlorine residual (calculated as a 7-day average) must be maintained at the end of the chlorine contact channel to ensure a sufficient bacterial kill. Prior to discharge into the ocean, sodium bisulfite is added for dechlorination, thus completing the disinfection process.

No changes or upgrades were made to the chlorine contact tank under the 2013 upgrade.

Sludge Treatment and Biosolids Disposal

On average throughout 2013, settleable solids and floatable materials from the primary clarifiers were treated in three heated anaerobic sludge digesters for approximately 38 days, with the average from January to May of 41 days and then from June to December of 36 days. Anaerobic digestion decomposes organic material and produces digester gas composed primarily of methane. This digester gas fuels boilers used to heat sludge in the digesters. Sludge from the digesters then flows to one of three stabilization basins where it settles and bacteria can continue the organic decomposition. Stabilized sludge is dredged from the bottom of these basins and is dewatered by compression through a belt filter press.

The 2013 upgrade involved the demolition of the third stabilization basin which was converted to an equalization basin. The belt filter press was removed and replaced with a new solids handling building which included two mechanical thickeners and two screw presses.

Because of the extensive construction taking place to upgrade the secondary portion of the treatment facilities, the sludge that would be air dried in the sludge drying beds and converted into Class A biosolids, for use by the local community has been temporarily discontinued. The belt pressed and then the screw pressed biosolids, identified as Class B, were transported under a new three year contract signed in 2012 by Western Express, Inc. to the Holloway Solid Waste Facility in Lost Hills, CA. The biosolids are used to reclaim the depleted gypsum mine pits.


A complete biosolids report describing the treatment and disposal process is prepared each year and submitted to the EPA. The deadline for submittal of this report is February 19th of each year.

Figure 1-1. Regional View of the Goleta Valley.



Figure 2-1. Treatment Process Flow Diagram for Blended Secondary Process Prior to Upgrade

PRIMARY TREATMENT **Collection System:** Wastewater is collected from homes and businesses, and flows by gravity or is pumped to the treatment plant through a network of sewers.



SECONDARY TREATMENT






Figure 2-1. Treatment Process Flow Diagram Prior to Upgrade (continued)

RECLAMATION TREATMENT



SLUDGE TREATMENT



Reclamation Facilities

On September 13, 1991, the California Regional Water Quality Control Board, Central Coast Region approved Order No. 91-03 that permits the Goleta Sanitary District to produce up to 3.0 MGD of reclaimed water. The reclaimed water produced at the Goleta Sanitary District is distributed by the Goleta Water District for use within their service area. Reclaimed water is used for landscape irrigation and for incidental uses including construction dust control and compaction, and to flush toilets within several buildings within Goleta. The Goleta Water District is regulated by separate water reclamation requirements.

Secondary effluent enters the reclamation facilities where a flash mixer disperses aluminum sulfate (alum) and polymer into the water. The flocculated suspension is then filtered through a bed of anthracite coal where the floc is removed. The filtered water then flows to a chlorine contact tank where sodium hypochlorite is added for disinfection. The highly chlorinated treated water then flows to a 3 million-gallon underground storage tank where it is stored until needed. Reclaimed water is distributed throughout the Goleta Valley by a distribution system operated and maintained by the Goleta Water District.

An annual report describing the reclamation treatment process, operational parameters, water quality, and production rates is prepared and submitted to the RWQCB by January 31st.

No changes were made to the reclamation facilities as part of the 2013 upgrade.

Ocean Outfall

The treated blended effluent is discharged to the ocean through an outfall pipe that extends 5800 feet offshore and terminates at a depth of approximately 92 feet below Mean Lower Low Water (MLLW) level. At the pipe terminus, a multi-port diffuser with 36, four inch diameter ports mixes one part of effluent with approximately 122 parts of seawater (Tetra Tech, Inc. 1993) to achieve a high initial wastewater dilution.

No changes were made to the outfall as part of the 2013 upgrade.

Staff

Mr. Kamil Azoury, P.E., serves as GSD's General Manager and District Engineer. The General Manager is responsible for overall operation and performance of the treatment plant.

Eight state certified treatment plant operators operate the wastewater treatment plant under the direction of Mr. Jeffrey Salt, the District Operations Manager. Mr. Salt also supervises the treatment plant's industrial waste staff. Mr. Chuck Smolnikar, supervises the maintenance staff and the laboratory is under the direction of Ms. Kathleen Werner, the technical services and laboratory supervisor. The grade and certification number of operations, maintenance, and laboratory personnel employed during the 2012 operational year are shown in Table 1-1.

Table 1-1. Goleta Sanitary District Operation Staff, 2013

Staff	Grade	California Certification No.
Operators		
Paul Buckley	V	7728
Robert Hidalgo	IV	6905
Stephen Conklin	III	7065
Todd Frederick	IV	IV-27633
Ricardo Lopez	III	III-10756
John Crisman	III	28857
Jose Sanchez	II	6400
Francisco M. Lemus	III	10893
Jes Hulbert	I	I-28266
Morgan Lee	I	I-28400
River Ferrara	I	I-28488
Lab Technologist		
Kathleen Werner	IV	070134001
Ray Giordano	III	070733003
Robert Hidalgo	I	741
Paul Buckley	I	1181
Teresa Kistner	I	99076111
Todd Frederick	I	060731013
Maintenance Technologist		
Charles Smolnikar		
Carl Easter	II	110662004
Mark Baumgartner	II	080722022
Paul Buckley	I	301
John Corral	I	770
Robert Hidalgo	I	1087
Mike Sullivan		
Jose Hernandez		

Monitoring and Reporting Program

The Goleta Sanitary District monitoring and reporting program was conducted in accordance with the requirements of the NPDES permit CA0048160. The objectives of the monitoring program and this report are to:

- Document short- and long-term effects of discharge on receiving waters, sediment, biota, and beneficial uses of the receiving waters.
- Determine compliance with NPDES permit terms and conditions.
- Document training and certification of wastewater treatment facility operators.

- Assess treatment plant performance and the effectiveness of industrial pretreatment and toxics control programs.
- Evaluate the monitoring and reporting program and make recommendations for improving the program.

The receiving water monitoring program consists of assessing water quality and ocean sediment chemistry, evaluating community structures of benthic biota, bottom fish, and epibenthic macroinvertebrates, and determining the bioaccumulation of pollutants in various marine organisms. Table 1-2 summarizes the sampling schedule for various elements of the monitoring and reporting program conducted during 2013.

Table 1-2. Schedule for NPDES Monitoring, Goleta Sanitary District, 2013

Monitoring Program Component	Frequency	Schedule
Standard Wastewater Parameters	Daily - Weekly	As Specified
Influent and Effluent Metals	Monthly	Every Month
Acute Toxicity	Quarterly	Jan, April, July, and Oct
Chronic Toxicity	Quarterly	Jan, April, July, and Oct
Influent and Effluent Priority Pollutants	Annually	October
Surf-Zone Bacteria	Weekly	Every Month
Receiving Water Bacteria	Quarterly	Jan, April, July, and Oct
Ocean Water Quality	Quarterly	Jan, April, July, and Oct
Benthic Sediments	Annually	October
Benthic Biota	Annually	October
Fish Trawls	Annually	October
Outfall Inspection	Annually	October
Bioaccumulation	Annually	October

Influent, effluent, and receiving water monitoring is conducted in accordance with U.S. Environmental Protection Agency approved test procedures as stipulated under Title 40 of the Code of Federal Regulations, Section 136 (40 CFR 136): *Guidelines establishing test procedures for the analysis of pollutants*. Water quality analyses for compliance monitoring are performed by analytical laboratories certified by the California Department of Health Services. Bioassay testing is conducted in accordance with guidelines approved by the State Water Resources Control Board and the EPA.

In order to comply with a request from the Central Coast RWQCB in a letter dated June 27, 2008 the District is no longer submitting hard copies of NPDES reports to the RWQCB. All documents are converted into a searchable PDF format and are submitted electronically. In light of this change the District contacted all other interested parties to whom monitoring reports are sent and except for the Division of Water Quality, all agreed to accept their routine reports electronically. The first report submitted this way was the June 2008 monthly reclaimed water report.

WASTEWATER TREATMENT UPGRADING PROJECT BACKGROUND

A condition of the November 2004 301(h) permit renewal included a provision to upgrade the current blended effluent treatment process to full secondary. Under a negotiated settlement agreement between the RWQCB and GSD the District agreed to follow a detailed conversion schedule to ensure that the treatment plant was discharging full secondary treated effluent by November 2014. The conversion schedule is shown below. The District completed the project in December 2013, the ninth year of the 10 year conversion timeline, almost a full year before the November 2014 required date.

The District awarded the facility planning contract to HDR Engineering and the environmental review contract was awarded to Tetra Tech. A preliminary draft of the facilities plan was sent to the Regional Water Quality Control Board in December 2006 and was circulated for review to all treatment plant contract users and other interested parties. The final facilities planning document was completed in June 2008.

A separate contract was then awarded to HDR Engineering to initiate and complete the design of the new treatment plant. The treatment plant design process addressed concerns regarding cultural resources and construction costs and allowed GSD to retain the use of current structures while planning for future regulatory changes. The new secondary treatment structures include the construction of a second biofilter, identical to the existing, an aeration basin and two new secondary sedimentation tanks. Primary and secondary solids will be co-thickened in mechanical thickeners located in a newly constructed solids handling building. The solids treatment will continue with anaerobic digestion, lagoon stabilization and finally, conversion to biosolids with two new screw presses.

Other features of the proposed upgraded plant include:

- ❑ Conversion of stabilization basin #1 into a flow equalization basin
- ❑ Construction of a second biofilter with a total media depth of 6 feet to match the existing biofilter
- ❑ Construction of a three train aeration basin with stub outs to add a fourth train at a future date if needed
- ❑ Construction of two new secondary clarifiers, and
- ❑ Construction of a solids handling building that will house the mechanical thickeners, polymer tanks and screw presses.

The District met all timelines specified in the conversion schedule except for C. 2. Environmental Review & Permitting, Certification of Final CEQA Document. The January 31, 2009 deadline to complete this milestone proved to be unattainable due to a flaw in the original negotiated agreement. The District requested and received, from the RWQCB, an extension for this task. The extension request was based on force majeure reasons caused by unforeseen cultural resource issues at the treatment plant site. The District conducted an extended phase 1 archaeology study in the areas of the new biofilter, proposed DAFTs and new pipeline corridors. Two inch geoprobes were drilled

approximately every 10 meters and the contents of the geoprobe were examined by a geomorphologist, an archaeologist and a Native American representative. The results of this extensive archaeological investigation indicated that the location proposed for the DAFTs had the possibility of containing some Native American artifacts and the District was advised to relocate these structures. Eventually the design was modified and the DAFT structures were removed and replaced with mechanical thickeners that were located in the southern portion of the plant.

No indication of artifacts were found in the location proposed for the new biofilter and corresponding pipeline corridors. These structures did not need to be relocated, however archaeologists and native American monitors were on site during the excavation of these areas.

CONVERSION SCHEDULE

Tasks	Date of Completion*
A. <u>Preliminary Activities:</u>	
1. Submittal of Detailed Conversion Plan and Timeline to Owners of Capacity in District's Plant	01/01/05
2. Coordination of Conversion concepts w/Owners of capacity in District's Plant (Education regarding participation in conversion)	06/30/05
3. Send Requests for Environmental & Consulting Engineering Contracts	12/31/05
4. Award of Environmental & Consulting Engineering Contracts	06/30/06
B. <u>Facilities Planning:</u>	
1. Complete Draft Facilities Plan	12/31/06
2. Complete Final Facilities Plan	06/30/08
C. <u>Environmental Review & Permitting:</u>	
1. Complete & Circulate Draft CEQA Document	06/30/08
2. Certify Final CEQA Document	01/31/09 06/30/10
3. Submit Applications for all Necessary Permits	01/31/09
4. Obtain all Necessary Permits	01/31/11
D. <u>Financing:</u>	
1. Complete Draft Plan for Project Design & Construction Financing	01/30/07
2. Complete Final Plan for Project Design & Construction Financing	03/31/08
3. Submit Proof that all Necessary Construction Financing has been Secured, Including Compliance with Proposition 218	12/31/10
E. <u>Design & Construction:</u>	
1. Initiate Design	06/30/08
2. 30% Design	12/31/08
3. 60% Design	11/30/09
4. 90% Design	03/31/10

5. 100% Design	09/30/10
6. Issue Notice to Proceed to Contractor	04/30/11
7. Construction Progress Reports	Quarterly (w/self monitoring reports)
8. Complete Construction & Commence Debugging and Startup	04/30/14
9. Full Compliance w/Secondary Requirements	11/01/14

*Any completion date falling on a Saturday, Sunday or State Holiday shall be extended until the next business day. The district shall submit proof of completion of each task within 30 days after the due date for completion.

By the end of December 2010, the District was successful in meeting all regulatory conditions and received all permits necessary to complete the project. For reference purposes, the following permits have been approved:

Permitting Agency	Type of Permit	Permit Number
Santa Barbara County	Government Code Consistency	09GOV-00000-00001
Santa Barbara County	Revised Development Plan	09RVP-00000-00001
Santa Barbara County	Grading Permit	09GRD-00000-00073
Santa Barbara County	Coastal Development Permit	09CDP-00000-00099
California Coastal Commission	Coastal Development Permit	4-09-011
Santa Barbara County	Land Use Permit	10LUP-00000-00235
Santa Barbara County	Land Use Permit	10LUP-00000-00360
Santa Barbara County Air Pollution Control District	Authority to Construct	13378
Santa Barbara County	Grading Permit	10GRD-00000-00075
Santa Barbara County	Building Permit	10BDP-00000-00553
Goleta Water District	Can & Will Serve Letter	

PCL Construction company was the low bidder and was awarded the construction contract. Their bid submittal was for \$28.6 M. The final cost of construction is still to be determined as change orders and costs of those change orders are still under discussion. To date the cost of the project has reached \$31 M. Mobilization took place in April 2011 and construction started in May 2011. A total of ten quarterly construction progress reports were prepared and submitted to the state and regional water quality control boards and several other interested parties. The last quarterly construction report was submitted on January 27, 2014 and covered the last quarter of construction work from July 1, 2013 to September 30, 2013. By the end of December 2012 all new structures had been built. The new biofilter, the aeration basin and one of the new secondary clarifiers had been put on line and were operational. Throughout 2013 some of the existing structures were taken off line for extensive renovations.

The plant began producing full secondary treated wastewater on May 16, 2013 when the final tie in was completed. PCL construction demobilized September 2013 and the project was deemed complete by December 2013.

REPORT ORGANIZATION

This report summarizes data collected during the 2013 monitoring and reporting program, and analyzes this data to determine compliance with the discharge permit terms and conditions. Chapters in this report have been organized to parallel sections of the monitoring and reporting program. The chapter sequence also follows the flow of wastewater as it undergoes treatment in the plant, as it is discharged to the marine receiving waters, and as it encounters nearby sediments and resident biota. Chapter 9 presents a summary of the lift station and collection system overflows, the causes of the overflows, the corrective actions taken, and any corrective actions planned. Chapter presentation is as follows:

Chapter 1	Introduction
Chapter 2	Treatment Plant Performance
Chapter 3	Receiving Water Environment
Chapter 4	Physical Characteristics of Benthic Sediments
Chapter 5	Chemical Characteristics of Benthic Sediments
Chapter 6	Biological Characteristics of Benthic Sediments
Chapter 7	Fish Populations
Chapter 8	Chemical Characteristics of Fish and Mussel Tissue
Chapter 9	Collection System Summary
	Appendices including the outfall dive survey

CHAPTER 2

TREATMENT PLANT PERFORMANCE

The performance of a wastewater treatment plant is measured by its ability to reduce influent contaminants to levels acceptable for discharge to the environment. Federal and state authorities mandate these levels of treatment in order to protect the marine environment. Proper operation of the Goleta Sanitary District's wastewater treatment plant is assured through the monitoring of several effluent parameters such as flow, total suspended solids, biochemical oxygen demand, residual chlorine, hydrogen-ion concentration (pH), turbidity, ammonia, settleable solids, oil and grease, and toxicity concentration. Metals, pesticides, and other priority pollutants are also analyzed to aid in determining the impact the wastewater discharge has on receiving waters, evaluating compliance with discharge permit limitations, and monitoring the effectiveness of the industrial pretreatment and toxic control program.

WASTEWATER CHARACTERIZATION

Goleta Sanitary District's NPDES monitoring program requires measurement of many parameters at frequencies ranging from continuous to once per year. During 2013, influent, effluent, biosolids (sludge), and surf zone samples were collected by treatment plant personnel, and analyzed by the Goleta Sanitary District wastewater treatment plant laboratory and various contract laboratories such as; Aquatic Bioassay Laboratories for ocean monitoring, Aquatic Testing Laboratories (ATL) for acute and chronic toxicity, FGL Environmental Laboratories and Exova, Vista Analytical Laboratory, Weck Laboratories as subcontractors to FGL. Treatment plant personnel monitored and analyzed wastewater for performance-evaluating parameters including wastewater flow, suspended solids, biochemical oxygen demand (BOD), pH, turbidity, settleable solids, ammonia, oil and grease, temperature, residual chlorine, coliform and enterococcus bacteria. Monthly analyses for influent and effluent metals were performed by FGL Environmental Laboratories of Santa Paula, CA. FGL Environmental Laboratories, and their certified subcontract laboratories performed annual analysis of priority pollutants and other parameters in influent, effluent, and biosolids samples. Influent and effluent samples were also analyzed for radioactivity. Bioassay tests for acute and chronic toxicity concentration were performed quarterly by Aquatic Testing Laboratory.

Analytical methodologies used by Goleta Sanitary District Laboratory and other contract laboratories used by GSD are based on approved U.S. Environmental Protection Agency (EPA) methods (EPA 1983; Federal Register 1984) and other methods in *Standard Methods for the Examination of Water and Wastewater, 21st ed.* (Standard Methods 2005). All methodologies employed during 2013 were approved for NPDES monitoring programs. Quality assurance and quality control procedures followed those presented in *Standard Methods for the Examination of Water and Wastewater, 21st edition.*

Results of the wastewater chemical analyses used to monitor proper operation of the treatment plant during 2013, and the respective discharge permit limitations, are presented


in Tables 2-1 and Table 2-2. All monthly averaged data presented in these tables are calculated from daily values at the treatment plant, with the exception of removal efficiencies, which are calculated from the monthly averages of the respective influent and effluent parameters.

Influent Flow

The daily influent flow into the treatment plant was monitored continuously throughout 2013. Influent flow without the internal plant recirculated flow, averaged 5.174 million gallons per day (MGD) a 6% increase over the average of 4.862 MGD that was treated in 2012.

Overall, the average monthly influent flows for 2013 were stable throughout the year, fluctuating from a low of 4.56 MGD in December to a 5.40 MGD in April. This is similar to the range of flows seen in 2012 and contrasts with a range of 1.6 MGD for 2011. No sharp spikes during the rainy winter months were observed as was seen in March of 2011. See Figure 2-1.

Figure 2-1. Influent Flows Monthly Average Comparison for 2011, 2012 and 2013



The highest flows into the plant during 2013 occurred in April, and may be associated with heavy rains that occurred in March.

Since 2001 the Goleta West Sanitary District and Goleta Sanitary District have maintained an aggressive collection system rehabilitation program. Numerous sections of the collection system in both Districts have been relined or replaced to correct structural deficiencies while significantly reducing the inflow and infiltration (I&I) problems. However,

even with the reduction of I&I the amount of rainfall during the year can affect the total amount of influent flow measured. The District's storm water pollution prevention plan requires all storm water collected from process areas to be treated before disposal. After several dry years the low ground water table and dry creeks can reduce the potential for ground water intrusion into the collection systems.

Effluent Flow

The effluent flow from the treatment plant was monitored continuously during 2013 and averaged 4.2 MGD for the year. The difference between the influent and effluent flow is due to the production of reclaimed water, which is not discharged into the ocean but is distributed throughout the community for landscape irrigation and other uses.

Figure 2-2. Influent and Effluent Flows 2013 Monthly Averages




Figure 2-2 shows the monthly average influent and effluent flows for 2013. Higher wastewater effluent flow generally occurs during the winter months when influent flow is also the highest and recycling is minimal. The most important factor contributing to fluctuations in the effluent flow is the amount of wastewater that is processed into reclaimed water and used for irrigation. The lowest effluent flow occurred during July when the amount of flow discharged to the Pacific Ocean dropped to 3.57 MGD as depicted in Figure 2-2. The temporal variations in the monthly average effluent flow seen in 2013 fluctuated from a low of 3.57 MGD in July, when the daily production of reclaimed water was the highest of the year and averaged 1.52 MGD for the month to a high of 5.149 MGD during January when the reclaimed facility was on line for 2 days out of the month and a total of 1.94 million gallons were filtered. January was also the rainiest month of the year with approximately 3.7 inches of rain. Figure 2-2 is a time history of the influent and effluent flows and Table 2-1 shows the actual monthly flow average values.

Table 2-1. Monthly Averages Flow, Suspended Solids and BOD, Goleta Sanitary District, 2013.

Month	Flow		Total Suspended Solids				Biochemical Oxygen Demand			
	Influent MGD	Effluent MGD	Influent mg/L	Effluent mg/L	Removal (%)	Mass Emission (lbs/day)	Influent mg/L	Effluent mg/L	Removal (%)	Mass Emission (lbs/day)
Jan	5.200	5.14	304	43	86	1870	295	68	77	2938
Feb	5.307	4.97	281	53	81	2239	280	83	70	3471
Mar	5.265	4.63	295	48	84	1853	272	62	77	2396
Apr	5.403	4.54	292	54	81	2067	287	85	70	3203
May	5.344	3.95	302	30	90	993	292	51	82	1720
Jun	5.013	3.62	310	8	97	242	303	6	98	194
Jul	5.084	3.57	393	7	98	215	310	5	98	159
Aug	5.115	3.69	389	6	98	183	335	5	98	152
Sep	5.260	3.90	354	10	97	322	314	5	98	171
Oct	5.387	4.24	325	9	97	332	315	5	98	174
Nov	5.152	4.44	359	9	97	326	312	4	99	161
Dec	4.560	4.06	370	8	98	280	349	4	99	135
Average	5.174	4.23	331	24	92	910	305	32	89	1240
Limit	NL	7.64	NL	63		4010	NL	98		6240

**NL = No Limit

Suspended Solids


Influent and effluent suspended solids were measured five days per week on 24-hour composite samples. The effectiveness of the treatment plant in removing suspended solids is demonstrated by the variation of influent solids versus the low-level and consistent output of effluent solids (see Figure 2-3). Influent suspended solids concentrations averaged 331 mg/L for the year an increase of about 10% from the 2012 annual average of 300 mg/L which was a 5% increase from the 286 mg/L annual average of 2011. For the past three years the concentration of suspended solids entering the plant has been on a steady increase. Figure 2-3 below shows a marked increase in concentration of suspended solids in the influent TSS beginning in July. This may be the start of the community reducing their water usage because of drought conditions.

The treatment process reduced the concentration of total suspended solids in the effluent to an annual average of 24 mg/L a 40% annual decrease of the 43 mg/L average of 2012. The average effluent TSS for the first five months of the year when the plant was still operating as a blended secondary was 46 mg/L and the average from June to December under full secondary conditions was 8 mg/L.

All 30-day monthly averages were well below the 63-mg/L monthly average limitation. The maximum daily value for 2013 was 94.5 mg/L, below the 100 mg/L maximum at any time limitation and occurred on Sunday, March 31, 2013..


Overall removal efficiency for the year was an average of 92 percent. Again the efficiency was higher after the completion of construction to full secondary, see Table 2-1.

Figure 2-3. Influent and Effluent Total Suspended Solids 2013 Monthly Averages



Average monthly suspended solids mass loading rates for 2013 are represented graphically in Figure 2-4. Mass loading calculations factor in flow rates and as such they correspond very closely with total plant flows and rainfall. Loadings are the highest during the wet winter months and drop to the lowest values during the dry summer months. The mass emission limit is based on average dry weather flow (ADWF) and is a limit applied to dry weather flows (DWF). There is no limit for mass emissions on wet weather flows.

The maximum average monthly mass emission loading for 2013 occurred in February at a high of 2,239 lbs/day, which is approximately half of the permitted monthly 30-day average limit of 4,010 lbs/day. The highest one-day maximum load for the year occurred on March 31, 2013 when 3,486 lbs of suspended solids were discharged. This maximum day discharge of total suspended solids loadings occurred on the same day as the maximum concentration occurred. The results of the transition to full secondary treatment are shown clearly in Figure 2-4. The discharge loading rates drop in May when the plant was still operating as a blended secondary during the first half of the month and then as a full secondary process during the second half of the month. Loading rates drop even further throughout the rest of the year.

Figure 2-4. Effluent Discharge Total Suspended Solids Mass Loading, 2013 Monthly Averages, lbs/day

Biochemical Oxygen Demand

Biochemical oxygen demand (BOD) levels were measured on 24 hour composite samples of the influent and effluent, at least three and five days per week, respectively.

During 2013 influent BOD averaged 305 mg/L showing a small increase from the annual influent average of 282 for 2012, and 271 for 2011. The increase in the influent BOD concentration mirrored that of the influent total suspended solids concentrations and may be reflective of a decrease in water use by the community in response to the beginning of drought conditions. The influent BOD increase slightly throughout the year, ranging from a monthly average low of 272 mg/L in March to a high of 349 mg/L in December.

The monthly average final effluent BOD concentration dropped dramatically throughout the year with the annual average of 32 mg/l and the range extending from a low of 4 in November and December to a high of 85 in April, (Table 2-1). This drop in BOD concentration occurred in May when the newly construction secondary treatment structures were tied into the existing facility and all of the flow could receive full secondary treatment. The difference between influent and effluent BOD represents an overall removal rate of 89 percent.

The maximum effluent concentration for 2013 was measured on April 21, 2013 at a concentration of 146 mg/L. Except for a slightly increased amount of flow through the reclamation facility which would increase the primary effluent/ secondary effluent flow ratio, no apparent reason could be given for this high value. The NPDES effluent BOD monthly average limitation and the maximum at any time limitation are 98 mg/L and 150 mg/L, respectively. All BOD NPDES limitations were achieved throughout the year.

Figure 2-5. Influent and Effluent Biochemical Oxygen Demand 2013 Monthly Averages




Table 2-2. Monthly Averages of Influent and Effluent Parameters, Goleta Sanitary District, 2013

	pH		Turbidity Effluent (NTU)	Settleable Solids Effluent (mL/L/hr)	Ammonia Effluent (mg/L)	Oil and Grease			Toxicity	
	Influent	Effluent				Influent	Effluent	Mass Emission (lbs/day)	Acute Effluent (TUa)	Chronic Effluent (TUC)
Jan	7.6	7.1	41	0.33	20	29.7	7.6	325		
Feb	7.7	7.2	46	0.39	27	46.9	6.8	272	0.91	17.9
Mar	7.7	7.2	37	0.28	29	25.0	5.3	200		
Apr	7.8	7.3	46	0.39	16	34.1	6.4	249	1.39	17.9
May	7.7	7.2	24	0.23	32	24.7	6.4	196		
Jun	7.6	6.9	3	0.14	0.1	25.1	< 4	< 122		
Jul	7.6	6.8	4	0.13	5	28.9	3.0	87	1.14	17.9
Aug	7.5	6.8	3	0.14	8	27.9	2.2	67		
Sep	7.4	6.7	4	0.15	0.2	51.4	1.8	62		
Oct	7.7	6.8	4	0.22	0.4	48.0	3.0	96	0.00	5.6
Nov	7.6	6.6	3	0.19	0.1	26.9	2.0	77		
Dec	7.8	6.6	3	0.12	0.1	32.1	1.2	47		
Average	7.6	6.9	18	0.23	11	33.4	4	153	0.86	14.8
Limit	NL	6 to 9	75	1.0	74	NL	25	1590	4.0	123

**NL = No Limit

In 2013, all effluent BOD mass emission values were below all limitations. The maximum

monthly average mass emission was 3,203 lbs/day for April. The maximum at any time mass emission was 5,683 lbs/day and occurred on April 14, 2013. The mass emission limit is based on average dry weather flow (ADWF) and is a limit, which is only applied to dry weather flows (DWF). There is no limit for mass emissions on wet weather flows. The mass emissions monthly average limitation of 6,240 lbs/day and the maximum at any time limitation of 9,560 lbs/day were never exceeded during 2013.

Hydrogen-Ion Concentration (pH)


Influent and effluent pH levels were monitored five days per week to ensure that the effluent remained within an acceptable range when discharged into the ocean. Influent pH averaged 7.6 units for the year; effluent pH averaged 6.9 units. Monthly averages of effluent pH dropped to below 7.0 after the full secondary treatment facility was put on-line in May 2013 (Table 2-2). However, as the NPDES effluent pH limitations are established as a minimum of 6.0 and a maximum of 9.0 pH units, all pH values were well within these limitations for 2013.

Ammonia

The effluent was monitored monthly to determine the concentration of ammonia. The permit specifies six-month median, daily maximum, and instantaneous maximum limitations of 74 mg/L, 300 mg/L, and 740 mg/L, respectively. The monthly average ammonia concentration ranged from 0.1 mg/L in June, November and December up to 32 mg/L in May (Table 2-2). The unusually low ammonia concentrations measured in June, November and December are indicative of some ammonia stripping that may be taking place in the newly constructed aeration basin. The increased aeration basin volume seems to have a direct impact on decreasing the concentration of the ammonia in the secondary treated effluent. Operations staff adjusted operational parameters during the start up and troubleshooting period for the new facilities. The monthly average for the year was 11 mg/L. The values for ammonia were well below all their respective permit limitations.

Turbidity

Effluent turbidity was monitored five days per week. The permit limitations for effluent turbidity consists of a monthly average of 75 Nephelometric Turbidity Units (NTU), a weekly average of 100 NTU, and a maximum at any time limitation of 225 NTU. Effluent turbidity data are shown graphically in Figure 2-6. The maximum value at any time, 67 NTU, occurred on May 15 and could be associated with the plant shut down that was taking place that night to tie in the newly constructed secondary structures. Monthly averages ranged from a low of 3 NTU to a high of 46 NTU. The dramatic drop in monthly averaged turbidity due to the conversion to full secondary treatment. (Table 2-2). All values were significantly below their respective permit limitations.

Figure 2-6. Effluent Discharge Turbidity 2013 Monthly Averages, NTU

Acute Toxicity Concentration

All quarterly acute toxicity tests were performed on 24-hour composite effluent samples. The acute toxicity has a daily maximum limit of 4.0 acute toxicity units (TU_a). All four quarterly acute toxicity samples for 2013 were collected under the conditions of the new NPDES WDR Order No. R3-2010-0012 which requires the District to use Topsmelt as the acute toxicity test species, replacing fathead minnow larvae. The annual average acute toxicity value was 0.86 TU_a . (See Table 2-2). All values were below the permit limitation of 4 TU_a and saw and improvement with full secondary treatment.

Chronic Toxicity Concentration

The effluent was analyzed for chronic toxicity (TU_c) on a quarterly basis in February, April, July, and October. The special testing conducted during 2011 to identify the most sensitive chronic toxicity organism showed that the abalone development test was the most sensitive. All results were well below the daily maximum limitation of 123 TU_c .

Settleable Solids

The effluent was monitored for settleable solids concentrations 5 days per week. The permit specifies that the monthly average, weekly average, and maximum at any time may not exceed 1.0 milliliters/liter/hour (ml/L/hr), 1.5 ml/L/hr, and 3.0 ml/L/hr, respectively.

Monthly averages ranged from 0.12 ml/L/hr to 0.39 mL/L/hr. The maximum value at any time was 1.0 mL/L/hr which occurred on January 10th. All values were well below their respective permit limitations.


Oil and Grease

Influent and effluent oil and grease were monitored bi-weekly (once every two weeks) and weekly, respectively. Monthly average results are shown graphically in Figure 2-7. Prior to August 2007 Freon was the solvent used in the standard method to extract oil and greases from water samples. According to EPA regulations, in August 2007 the GSD laboratory ceased using Freon as the extraction solvent and began using hexane as the required solvent. The District continued to use the liquid-liquid extraction method, the only change at this time was the solvent. In December 2010 the GSD laboratory began analyzing for oil and grease using the approved standard solid phase extraction (SPE) method.


Influent grease and oil results were very inconsistent throughout the year. Average monthly concentrations spiked in February, September and October. February had one high sample on the 14th of 78 mg/L which caused the increase in the monthly average as did October with one high sample collected on the 7th that resulted in 86 mg/L of grease and oil.

Effluent grease and oil concentrations show what is now becoming a typical result of decreased concentration of measured parameters after the full secondary treatment plant became operational in May 2013.

The influent annual average value of 33 mg/L was reduced to an annual average of 4 mg/L in the final effluent resulting in an 88 percent annual average removal rate. All monthly, weekly, and maximum permit limits were met. Mass emissions values ranged from a monthly average low of 47 lbs/day in December to a high of 325 lbs/day in January. Both are well below the permit limitation of 1,590 lbs/day. Monthly average oil and grease concentrations in the effluent ranged from 1.2 mg/L in December to 7.6 mg/L in January. (Table 2-2). All permit limitations for effluent oil and grease were met during 2013.

Figure 2-7. Influent and Effluent Grease and Oil 2013 Monthly Averages**Temperature**

Effluent temperature was sampled five days per week throughout 2013. The data reflect a typical response to seasonal changes (Figure 2-8). The coolest temperatures occurred during January and December with an average monthly temperature of 20.0 °C. A warming trend continued throughout the spring and summer months to reach a monthly averaged high in July and August of 25.6 °C. As expected, the year ended with a cooling trend during the fall and winter months (October through December).

Figure 2-8. Effluent Discharge Temperature 2013 Monthly Averages

Wastewater Disinfection

Sodium hypochlorite is used to disinfect the treated wastewater at the Goleta Sanitary District. The sodium hypochlorite is flash mixed into the wastewater at the beginning of the chlorine contact channel. At an average effluent flow rate of 4 MGD, the chlorine is in contact with the wastewater for approximately 2½ hours (145 minutes). The NPDES permit specifies that the District must maintain a total chlorine residual of at least 5 mg/L at the end of the chlorine contact channel under total suspended solids peak loading conditions. The Goleta Sanitary District maintains its chlorine contact tank to provide maximum chlorination effectiveness at all times. The chlorine residual at the end of the chlorine contact channel averaged 8.0 mg/L during 2013. The average monthly values are reported in Table 2-3.

After the disinfection process is completed, the sodium hypochlorite is neutralized (dechlorinated) by adding sodium bisulfite to the wastewater stream. This process lowers residual chlorine to levels that are environmentally safe, before discharge to the ocean such that the chlorine poses no risk to the receiving water environment. Treatment plant personnel continuously monitor the residual chlorine levels as required by the NPDES permit.

The permit limitations for residual chlorine in the effluent immediately prior to discharge and after dechlorination are as follows: 6-month median of 0.25 mg/L, daily maximum of 0.98 mg/L, and instantaneous maximum of 7.4 mg/L. After dechlorination, the monthly average residual chlorine levels were very consistent throughout the year; at or below the detection limit of 0.1 mg/L for all months. The monthly average values are shown in Table 2-3. No chlorine residual exceedences occurred during 2013.

Effluent Coliform Bacteria

The effluent was analyzed five days a week for coliform bacteria. The monthly average values for total coliform, fecal coliform, and enterococcus bacteria detected in the effluent are presented in Table 2-3. Monthly average values ranged from 25 to 433 MPN/100 mL for total coliform and from 5 to 243 MPN/100 mL for fecal coliform. The permit prohibits more than 10 percent of the final effluent samples, in any thirty-day period, to exceed a total coliform density of 2,400 MPN/100mL with no sample exceeding a total coliform concentration of 16,000 MPN/100mL. A total of 302 final effluent total coliform samples were analyzed throughout the year with no samples exceeding either the 30-day limitation or the 16,000 MPN/100mL limit. The maximum total coliform concentration was measured on June 13, 2013 at 9,200 MPN/100mL.

Effluent Enterococcus Bacteria

The effluent was also analyzed five days a week for enterococcus bacteria. The monthly mean values are presented in Table 2-3 and the values were consistently low throughout the entire year, thereby demonstrating the effectiveness of the chlorination process.

Table 2-3. Chlorine and Bacteria Monthly Averages, 2013

Month	Chlorine at the end of the CCC	Chlorine after Dechlorination	Total Coliform	Fecal Coliform	Enterococcus
	mg/L	mg/L	MPN/100mL		
January	9.2	< 0.1	54	18	1.9
February	8.8	< 0.1	84	18	2
March	8.7	< 0.1	63	17	2
April	9.0	< 0.1	39	17	2
May	8.4	< 0.1	56	22	2
June	7.8	< 0.1	433	243	2
July	8.1	< 0.1	25	13	3
August	7.5	< 0.1	34	20	2
September	7.3	< 0.1	49	19	2
October	6.8	< 0.1	65	13	2
November	7.1	< 0.1	57	5	2
December	7.1	< 0.1	61	5	2

SURF ZONE BACTERIA

The Goleta Sanitary District has an extensive bacteria monitoring program that measures the concentrations of enterococcus, total coliform, and fecal coliform groups of bacteria at the end of the treatment process immediately before discharge to the ocean, at the end of the pipeline in the zone of initial dilution, at far shore and near shore ocean sampling locations and in the surf zone at stations extending west from Goleta Point to 1,000 meters east of the outfall line. Table 2-4 summarizes the locations and frequency of all bacteria monitoring conducted at the Goleta Sanitary District.

Table 2-4. Bacteria Monitoring Program

Location	Frequency of Total Coliform, Fecal Coliform and Enterococcus Bacteria Testing
Final Effluent prior to ocean discharge	5 days/week
Zone of Initial Dilution in the discharge plume at 25 m and 100 m from outfall pipe	Quarterly: 3 samples at each location; 1m below surface, mid-depth and 1 m above bottom
Far Shore (ocean) Stations; B1, B2, B3, B4, B5 and B6	Quarterly: 3 samples at each location; 1m below surface, mid-depth and 1 m above bottom
Near Shore (ocean) Stations; K1, K2, K3, K4 and K5	Quarterly: 3 samples at each location; 1m below surface, mid-depth and 1 m above bottom
Surf Zone Stations; A, A1, A2, B, C, D, E	Weekly

Final effluent samples and weekly receiving water surf zone samples are collected and analyzed in-house by GSD personnel the results of which are discussed in this chapter. Zone of initial dilution, far shore and near shore bacteria samples are collected and analyzed by ABC Laboratories of Ventura. Results of this testing is presented in chapter 3.

Approximately 336 samples are collected each year from the surf zone and each sample is analyzed for total coliform, fecal coliform and enterococcus for a total of approximately 1,008 bacteria tests conducted every year. These samples are collected and indicator organism concentrations are monitored in order to ensure that the beneficial uses of the Goleta Beach coastal area are protected. The following section discusses the 2013 bacterial trends found in the surf zone environment.

Surf-zone Stations.

Consistent with historical trends, bacteria monitoring at surf-zone stations usually yield more frequent and higher amounts of coliform bacteria than at the near shore and farshore (ocean) stations and even from the final effluent that is discharged to the ocean. The occurrence of bacteria in the shoreline monitoring area is often in response to the drainage, tidal flushing, and dredging of Goleta Slough. Over the years it has been determined that coastal bird populations, organic beach debris (including dog waste), and most importantly, the urban flushing effects of storm water runoff can be contributors to high surf zone bacteria concentrations. There has never been any indication that the treatment plant discharge has contributed to bacteria concentrations along the shoreline.

Goleta Slough, which is the confluence of the San Jose, Atascadero, and San Pedro creeks, is a slow-flowing, estuarine water body, which discharges directly into the Pacific Ocean between two of the Goleta Sanitary District's monitoring stations (stations D and E). Because the slough receives little flushing (except during storm runoff episodes) and is a rich waterfowl habitat, slough waters are relatively high in organics and coliform bacteria with respect to surf-zone waters.

Concentrations of bacteria at surf-zone stations in 2013 in general, were higher than that observed in the effluent, offshore and near shore ocean stations. This is consistent with the results of earlier years. Throughout the year, levels of bacteria at surf-zone monitoring stations ranged from < 1.8 to $\geq 1,600$ MPN/100mL for total coliform, <1.8 to 920 MPN/100mL for fecal coliform bacteria and <1.8 to 240 MPN/100mL for enterococcus bacteria. Several maximum one time exceedences occurred throughout the year and were reported in the corresponding monthly report. Table 2-5 is a summary of the 2013 surf zone exceedences.

Table 2-5. Surf Zone Exceedences 2013

Date	Station	Exceedence	Possible Cause	Final Effluent Result
1/7/2013	D	One time enterococcus \geq 104 MPN/100mL	Rain runoff contamination	< 1.8 MPN/100mL
	E	One time total coliform \geq 10,000 MPN/100mL	Rain runoff contamination	18 MPN/100mL
1/13/2013	B	One time enterococcus \geq 104 MPN/100mL	No clear reason	< 1.8 MPN/100mL
1/28/2013	D	One time fecal coliform \geq 400 MPN/100mL	Residual Goleta Slough contaminated runoff	< 18 MPN/100mL
4/18/2013	D	One time fecal coliform \geq 400 MPN/100mL	Residual Goleta Slough contaminated runoff	< 18 MPN/100mL
9/6/2013	A2	One time enterococcus \geq 104 MPN/100mL	Large amount of kelp and large number of sea birds.	< 1.8 MPN/100mL
9/18/2013	A2	One time enterococcus \geq 104 MPN/100mL	Large amount of kelp and large number of sea birds.	< 1.8 MPN/100mL
Sept 2013		30 day enterococcus gmean \geq 35 MPN/100mL	Large amount of kelp and large number of sea birds.	
10/6/2013	A2	One time enterococcus \geq 104 MPN/100mL	Large amount of kelp and large number of sea birds.	< 1.8 MPN/100mL
10/20/2013	A1	One time enterococcus \geq 104 MPN/100mL	Large amount of kelp and large number of sea birds.	< 1.8 MPN/100mL
12/6/2013	A2	One time fecal coliform \geq 400 MPN/100mL	Large amount of kelp and large number of sea birds.	5 MPN/100mL

In January the exceedences were most likely the result of contaminated rain water runoff that occurred as a result of the two rainstorms in January as the sampling occurred in days immediately following a rainstorm. Exceedences in surf zone bacteria that occurred later in the year were almost exclusively limited to Stations A1 and A2. Observations recorded on the sample days noted that there was a large amount of kelp on the shore between Stations A1 and A2 which typically attracts a large number of sea birds and which could be the source of the higher than normal bacteria.

Throughout the year the final effluent samples analyzed previous to and on the surf zone collection days indicated no or very low concentrations of coliform and/or enterococcus bacteria, see Table 2-5.

No samples were collected from Station E on January 13th because Station E was inaccessible due to high flows in the Goleta Slough and the Slough could not be crossed safely by District staff.

Although the range of bacteria concentrations was large throughout the year, the average values for the year were 67 for total coliform, 49 for fecal coliform and 8 for enterococcus. These values were somewhat lower than 2012 and may be due to the mouth of the Goleta Slough remaining closed for a good part of 2013. Santa Barbara County Flood Control

District had in the past a program where they regularly dredged the mouth of the slough open, but this program was suspended during 2013. As a result the mouth of the Slough was closed off by a build up of a natural sand berm between the ocean and the slough.

Figure 2-9. Surf Zone Annual Average Bacteria Concentrations 2013




Figure 2-9 shows the impact of the Goleta Slough discharge on the surf zone samples. Goleta Slough empties between station location D and E which show some of the highest overall annual average bacteria concentrations for all three indicator organisms measured weekly. Except for station A2 which had a large amount of kelp washed up on the beach for several months of the year and accompanying sea birds contributing to coliform bacteria concentrations, the further the station is from the slough mouth the lower the concentration of bacteria measured until Station A, located at Campus Point, the furthest point west with the “cleanest” samples.

Effluent bacteria samples collected at the end of the treatment and disinfection process, during these same time periods showed low or undetected concentrations of bacteria discharged from the treatment plant demonstrating that the effluent was not a source for the high surf zone bacteria concentrations.

The impact of Goleta Slough on bacteria water quality in the surf zone of the study area has been documented for the past 22 years. This historical data has shown, year after year that the highest concentration of indicator organisms are found in and adjacent to the Goleta Slough mouth and are associated with storm water run off.

Metals

Twenty four-hour composite samples of influent and effluent were collected monthly and analyzed for metals (Table 2-6). In all instances, the concentrations of metals in the effluent for 2013 (Table 2-6) were low or undetected and were well below all permit limitations. Although the wastewater treatment process is not particularly efficient at removing metals, hence the need for the pretreatment program, with the upgrade of the treatment plant to full secondary treatment in May 2013 the concentrations of some metals in the final effluent showed a marked decrease from concentrations detected with the blended secondary process. Particularly, chromium, copper, lead and zinc appear to have been removed from the wastewater with the full secondary treatment. Metal concentrations in the influent were consistent throughout the year.

Table 2-6. Influent and Effluent Metals (ug/L), Goleta Sanitary District, 2013.

	Arsenic	Cadmium	Chromium	Copper	Lead	Mercury	Nickel	Silver	Zinc
Influent (ug/L)									
January	< 2	0.2	3	84	1.5	0.06	5	< 1	130
February	< 2	< 0.2	4	77	2.9	0.09	6	< 1	120
March	1.45	0.176	3.36	87	2.28	0.0906	5.44	1.08	137
April	1.75	0.219	2.74	93.5	1.68	0.0679	5.93	0.911	120
May	1.45	0.285	3.68	109	5.23	0.0813	6.13	1.82	152
June	1.60	0.161	2.93	88.4	2.04	0.109	4.8	0.604	144
July	1.49	0.223	4.17	116	2.08	0.103	6.22	1.56	151
August	1.24	0.218	2.53	96.2	2.26	0.116	5.25	0.668	115
September	1.49	0.283	6.64	138	2.50	0.180	7.58	1.82	171
October	1.26	0.234	3.32	79.4	1.50	0.251	5.28	1.31	117
November	1.67	0.263	6.67	120	2.30	0.2260	8.01	0.912	168
December	1.12	0.291	4.52	37.2	2.06	0.0107	6.4	0.425	170
Effluent (ug/L)									
January	< 2	< 0.2	2	32	0.6	< 0.02	5	< 1	60
February	< 2	< 0.2	2	32	0.6	0.02	10	< 1	70
March	1.08	0.088	1.76	32.8	0.978	0.0156	4.52	0.33	74.2
April	1.31	0.077	1.25	23.4	0.548	0.0267	4.86	0.173	49
May	1.06	0.176	1.57	32.8	5.54	0.0224	5.24	0.361	50.1
June	1.07	< 0.2	0.594	7.07	0.851	0.0077	4.53	0.025	41.9
July	0.807	< 0.2	0.583	8.08	0.241	0.00727	4.36	0.031	26.4
August	0.813	< 0.2	0.299	5.67	0.168	0.00944	3.20	0.018	18.2
September	0.783	< 0.2	0.469	5.09	0.178	0.0466	3.32	0.018	29.6
October	0.812	0.0310	0.498	6.12	0.172	0.0147	4.24	< 1	39.0
November	0.968	0.0380	0.520	8.74	0.035	0.00964	4.14	< 1	38.5
December	0.957	0.0480	0.636	6.67	0.474	0.0122	4.26	0.016	53.8
Effluent Limits (ug/L)									
6-month median	620	120	250	120	250	4.9	620	67	1,500

Priority Pollutants

The NPDES permit requires priority pollutant analyses to be performed on influent and effluent composite samples annually. Compounds detected in the influent and/or effluent samples are presented in Table 2-7; complete copies of all the laboratory reports listing all the chemical compounds and analytical methods are available for review at the Goleta Sanitary District laboratory. Fourteen compounds were detected in the influent and eleven in the effluent. Diethylphthalate, phenol, carbon disulfide, methylene chloride and trichloroethylene were detected in the influent but not in the effluent. Whereas a small amount of chloromethane and bromoform were detected in the effluent but not in the influent. Acetone was detected in both the influent and effluent. It has been one of the most consistently detected chemicals in the wastewater stream. The most likely source of acetone entering the treatment plant is probably the University of California at Santa Barbara where acetone is used extensively in many of the research laboratories. Concentrations of detected chemicals are all reported as parts per billion.

Table 2-7. Detected Priority Pollutants, Goleta Sanitary District, 2013

Parameter, units	Influent, ug/L	Effluent, ug/L
Acetone	553	5.31
Antimony	1.76	1.06
Bis(2-Ethylhexyl)phthalate	1.45	15.3
Bromodichloromethane	1.26	32.7
Chloroform	7.73	40.3
Chloromethane	ND	0.252
Dibromochloromethane	0.690	15.4
Diethylphthalate	1.66	ND
TCDD, equivalents, pg/L	0.24567	0.00388
Phenol	10.5	ND
2-Butanone (MEK)	5.65	1.61
Carbon Disulfide	15.7	ND
Methylene Chloride	1.43	ND
Toluene	0.448	0.702
Trichloroethylene	0.150	ND
Bromoform	ND	0.788
Radioactivity, gross Alpha pCi/L	0.127 +/- 1.90	1.49 +/-1.62
Radioactivity, gross Beta pCi/L	3.76 +/-2.31	4.92 +/-2.01
ND = Not Detected		

Results of influent and effluent radioactivity determinations for 2013 are also presented in Table 2-7. Limits for radioactivity are defined in Title 17 of the California Code of Regulations section 30269, which state limitations of 3×10^{-8} $\mu\text{Ci/mL}$ (or 30 pCi/L) for alpha emission and 3×10^{-6} $\mu\text{Ci/mL}$ (or 3000 pCi/L) for beta emission. Samples collected during 2013 were below these limitations.

DISCHARGE COMPLIANCE

Throughout 2013 the wastewater discharge from Goleta Sanitary District complied with all applicable permit effluent limitations. All monitored parameters were below their respective limitations as required by the permit. All metals, priority pollutants, and pesticides were low or undetected throughout the year.

OCEAN OUTFALL CONDITIONS

The outfall pipeline, diffuser section, and armor rock protection were inspected by divers from Aquatic Bioassay and Consulting Laboratories, Inc. on October 30, 2013. A report was prepared and videotape was made of the diffuser section and along the outfall pipeline and armor rock.

During the diffuser dive survey, 36 diffuser ports were carefully inspected for flow and general efficiency. The remainder of the outfall pipe was inspected for damage, leaks or evidence of leaks and general stability of the pipe and armor rock. Inspection of the outfall yielded no evidence of damage, holes, cracks, or erosion. The pipe and associated armor rock appeared stable with little or no displacement.

The complete report of the outfall dive survey is included as an appendix to this report. Copies of the outfall dive on DVDs are available at the District for review.

CHAPTER 3

Receiving Water Environment

3.1. Scope and Period of Performance

This report covers the period of field and laboratory studies conducted from January 1, 2013 through December 31, 2013. The Aquatic Bioassay consulting team conducted water quality surveys in the vicinity of the of the Goleta Sanitary Districts outfall on January 24th, April 4th, July 16th, and November 13th, 2013. The team evaluated the local effect of the discharge within the immediate vicinity of the outfall terminus, and compared conditions there with those at control sites up-coast and down-coast of the outfall. During each field survey, the team recorded general observations of weather, etc., sampled for bacteria and water column variables (temperature, salinity, pH, transmittance and dissolved oxygen). On July 16th, the team deployed a series of caged mussel arrays for bioaccumulation analysis and on October 31st, the team retrieved the mussels. On October 15th, the team collected epibenthic fish and macroinvertebrates by otter trawl, and collected benthic sediments for physical, chemical, and infaunal analysis using a Van Veen Grab.

3.2. Station Locations and Descriptions

Water-column monitoring was conducted at ocean stations that are located at fixed distances from the midpoint of the diffuser (Figure 3-1). Stations B4 and B5 are located at the boundary of the zone of initial dilution (ZID), 25 meters (m) west and east of the diffuser, respectively. Station B2 and B3 are near-field stations located 500 and 250 m west of the diffuser, respectively. Station B1 is a far-field station located 1500 m west of the diffuser offshore Goleta Point. Station B6 is a reference station located 3000 m east of the diffuser. Plume stations WCZID and WC100 are respectively located 25 and 100 m away from the discharge in the direction of current flow. Nearshore Stations K1 through K5 are also at fixed distances west and east of the outfall in 20 m of water. Historically, the location of the 20 m depth contour represents the offshore limit of kelp beds in the study area.

Mussel arrays were deployed at Stations B3, B4, and B6. Trawl sampling was initiated at Stations B3 moving west for ten minutes and at Station B6 moving east for ten minutes (trawl stations TB3 and TB6, respectively).

3.3. Navigation and Positioning

The outfall diffuser and all sampling stations were located using a *Lowrance Global Map 2000* differential global positioning system (DGPS). DGPS positions were checked visually and by bottom-finder. Once the outfall terminus location was verified, a water quality analyzer cast was taken directly over the diffuser and water quality profiles were simultaneously downloaded to an onboard computer. Aquatic Bioassay biologists inspected the water quality traces for excursions from ambient such as higher temperature or lower salinity, dissolved oxygen, light transmittance, or pH. Any of these would reflect the presence of the wastewater plume. Once the plume was identified, a sail-drogue was deployed over the diffuser at the same depth as the discharge plume signature. The drogue was allowed to move with the current until an obvious direction and velocity could be determined. Stations WCZID (25 m from terminus) and WC100 (100 m from terminus) were then positioned along the drogue's line of travel.






Figure 3-1. Goleta Sanitary District receiving water monitoring stations. Trawl stations are represented by arrows (--->).

3.4. Statistical Analysis

For this report, two types of statistical tests were performed; trend analysis using correlation coefficient analysis, and comparative analysis using t-tests and analysis of variance (ANOVA). For this report, statistical significance is highlighted at two levels. For most ecologists, a pattern that is strong enough so that there is only a one chance or less in 20 that it is random is said to be statistically significant. In other words, the probability (p) is that there is only a 5% chance (0.05) or less that the pattern is random ($p < 0.05$). A pattern that has only one chance in ten or less (but more than one chance in 20) is said to be "marginally significant". That is, the probability is less than 10% but greater than 5% of being random ($0.05 < p < 0.10$).

3.5.1. Correlation Coefficients. Correlation analysis compares two variables to determine if they tend to increase or decrease in the same way. If two measurements tend to vary in opposite ways, their correlation coefficient (r -value) will tend to have a negative sign. If two measurements tend to vary in the same way, their r -value will tend to have a positive sign.

In addition to its sign, the size of an r -value is important. r -values range from -1.000 to $+1.000$. An r -value of -1.000 means that the two measurements being compared vary exactly opposite from each other, an r -value of $+1.000$ means that the two measurements vary



exactly in the same way, and an r-value of 0.000 means that the two measurements have no relationship to each other at all. Most r-values, however, fall somewhere among these three values. Depending upon the number of samples that are used to represent the true population, we have more confidence in our r-values when they are high. If an r-value is large enough so that the chance that the relationship could be random is only one in 20 or less ($p \leq 0.05$), we can have confidence that the relationship is probably real. We would have less confidence in a relationship between two variables if the probability was only one in ten ($0.05 < p \leq 0.10$) and no confidence if it was greater than ten ($0.10 < p$).

Based upon experience from past studies, we know that wastewater discharges can negatively impact the marine environment in very specific ways. If the outfall discharge is causing chemicals to accumulate in sediments and/or tissues, it follows that their concentrations would be higher nearer the diffuser than farther away. In this report, the distances of the stations from the diffuser were correlated against the concentration of the individual chemical components that were measured from these stations. Thus, the sign of the correlation coefficient between distance from outfall and chemical concentration would be *expected* if that chemical correlation was *negative*. That is, as the distance from the outfall becomes *larger*, the concentration of the compound becomes *smaller*. Another r-value that is expected to be negative is temperature. The effluent is always warmer than the ocean water, so temperatures, like chemicals, would be expected to become smaller with larger distances.

If the discharge were disrupting biological communities; abundance, diversity, etc., it would be expected to be lower near the outfall than farther away. Thus, population variables would be *expected* to correlate *positively* with distance from outfall, i.e. as distance becomes *larger* these variables would become *larger*. However, it is well documented that infauna populations can thrive near the nutrient enriching effects of ocean outfall where nutrients have enriched the area (Pearson and Rosenberg 1978). A positive and significant correlation between distance from the outfall abundance, numbers of species and diversity could signal that this is the case. Other r-values that are expected to be positive with distance are salinity, pH, dissolved oxygen, surface transparency, and light transmissance. This is because effluents are usually less saline, less clear, and lower in dissolved oxygen and pH than ocean water. If the discharge were affecting the receiving waters, an increasing pattern of these variables with distance from outfall would be expected.

In conclusion, variables that vary in patterns that are both expected and significant should be those which bear further scrutiny.

3.5.2. T-tests. This statistic is used to compare variables when there are only two. Unlike correlation coefficients, the trend with distance is not evaluated. For most variables, the mean of values near the outfall and the mean of values farther away will be different. The t-test determines whether or not that difference is statistically significant. Note that trend with distance or sign of the statistic is not of importance for this test. The question asked is only if they are different beyond what might be expected of random chance.

T-tests are used in this report for trawled fish and invertebrate population metrics and chemical compounds in fish tissue, since these variables were replicated and collected at two locations (i.e. TB3 and TB6). If the average difference in concentration of a chemical compound between these two stations is large enough that the probability is less than or equal to 5% ($p \leq 0.05$), the difference is said to be statistically significant. If the difference is large enough so that the probability is less than or equal to 10% but greater than 5% ($0.05 < p \leq 0.10$), the difference is said to be marginally significant. If the concentration of the compound



is larger at the near-outfall station, and the t-test is significant, the pattern should be further evaluated.

3.5.3. Analysis of Variance (ANOVA). ANOVA is similar to the t-test, except it can be used test for significant differences among more than two stations. ANOVAs were used for population variables and tissue analysis of bivalves. ANOVA analysis requires two steps. In the first step, differences in a variable among stations are evaluated to determine if they are sufficiently large to be statistically significant ($p \leq 0.05$). If they are, then a second test must be performed to determine which stations' variables are significantly larger than which other station or stations. In this report, this second step is called the comparison of means. For example, a comparison of means stating: $B1 > B2$, $B3 > B4$, indicates that, for that particular variable, Station B1 is significantly larger than Stations B2, B3, and B4, and Stations B2 and B3 are also significantly larger than Station B4. For chemical contaminants, if stations near the outfall are significantly higher than stations farther away, that compound should be evaluated further. For population variables, the opposite is true.

3.6. General Oceanographic Conditions

With the exception of somewhat sporadic freshwater runoff from non-point sources, the aquatic conditions in Goleta offshore area are controlled by the oceanographic conditions in the Southern California Bight. The mean circulation in the Southern California Bight is dominated by the northward-flowing Southern California Countercurrent, which may be considered as an eddy of the offshore, southward-flowing California Current (Daily, et. al. 1993). Nutrient rich, upwelled waters from the California Current can enter the western end of the Santa Barbara Channel promoting primary productivity (Dugdale and Wilkerson, 1989). The California Countercurrent transports nutrient poor, warmer water northward into the eastern Santa Barbara Channel (Hickey 1998). The California Countercurrent is seasonal in nature and is usually well developed in the summer and fall and weak (or absent) in winter and spring (SCCWRP 1973). This causes relatively nutrient-poor waters to predominate in the warmer water months and nutrient rich waters to predominate in the colder water months (Soule, et. al. 1997).

Superimposed upon annual trends are the sporadic occurrences of the El Nino Southern Oscillation (ENSO) that can be described as an oceanographic anomaly whereby particularly warm, nutrient-poor water moves northward from the tropics and overwhelms the typical upwelling of colder nutrient-rich water. The El Nino Watch (<http://coastwatch.pfel.noaa.gov/erddap/index.html>) program continuously monitors global sea surface temperatures. These temperature data are compared to the long-term sea surface temperatures generated from data collected from 1950 to 2013. Comparison of the monthly sea temperature with this long term average creates a temperature anomaly so that the average monthly temperature falls either above or below the average. This anomaly allows us to determine how a given month or time period deviates from the long term ocean temperature trend. The water temperatures offshore Goleta was at or up to one degree below the long term trend for January, March, April and December. Beginning in May and lasting through October, temperatures were 0.5 °C above the average, except in September when temperatures were 2.5 °C above the average (Figure 3-2).






Figure 3-2. Sea surface anomaly temperatures for 2013 compared with long term trends.

3.7. Anthropogenic Inputs

In addition to the Goleta discharge, several other natural and anthropogenic sources could potentially impact the coastal area. Three marshes (Devereux Lagoon, Campus Lagoon, and Goleta Slough) and several creeks discharge into the local area. All are a potential source of contaminated water and sediments, coliform and enterococcus bacteria, and nutrients; particularly during the rainy season. Several sources of crude oil are also present. Natural seeps occur west of the diffuser in the vicinity of Coal Oil Point and Goleta Point, and offshore production activity occurs throughout the Santa Barbara Channel.

3.8. Rainfall

Total rainfall is not as important in terms of impacting an area as the timing of the rainfall, the amount in a given storm, and the duration of a storm (or consecutive storms). Relative to timing, the first major storm of the season will wash off the majority of the pollutants and nutrients accumulated on the land over the preceding dry period. An early, large, long duration storm would have the greatest impact on the waters. In addition, determining the impact of the rainfall and runoff is also a function of the timing of the sampling surveys. With a greater lag between runoff and survey sampling, mixing with oceanic waters would reduce observable impacts (Soule, et. al. 1996).

The rainfall reported in this document is for Santa Barbara Airport obtained from the Western Regional Climate Center in Reno, Nevada. Data is summarized in Table 3-2 and Figure 3-3, where periods of precipitation and water column survey days are highlighted. The rainfall for this period (4.71 inches) was 13.57 inches below the average yearly rainfall since 1981 (18.96 inches). The wettest month was January (2.04 in), followed by March (1.06 in), and November (0.74). No rain fell in August and September. Rain in all other months ranged from 0.01 to 0.18 inches. Each of the water quality surveys occurred following periods of no rain, except in May with a (0.01 in) rainfall and January when rain fell before, during and after the sampling event.



Receiving Water Environment

Table 3-2. Daily 2013 Santa Barbara Airport rainfall (inches) with dates of water column surveys bordered and rain days in gray.

Day/Month	January	February	March	April	May	June	July	August	September	October	November	December
1	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
2	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
3	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.00	0.00	0.00	0.00	0.00
4	0.00	0.00	0.00	0.01	T	0.00	0.00	0.00	0.00	0.00	0.00	0.00
5	0.10	0.00	0.00	0.00	T	0.00	0.00	0.00	0.00	0.00	0.00	0.00
6	0.23	0.00	0.01	0.00	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00
7	0.00	0.00	0.62	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.16
8	0.00	0.05	0.06	0.00	0.00	T	0.00	0.00	0.00	0.00	0.00	0.00
9	0.00	0.00	0.00	0.00	0.00	0.02	0.00	0.00	0.00	0.03	0.00	0.00
10	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
11	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
12	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
13	0.00	0.00	0.00	0.00	0.00	T	0.00	0.00	0.00	0.00	0.00	0.00
14	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
15	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
16	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
17	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
18	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
19	0.00	0.13	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
20	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.31	0.00
21	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
22	0.00	0.00	0.00	0.00	0.00	0.00	0.07	0.00	0.00	0.00	0.02	0.00
23	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
24	1.48	0.00	0.00	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
25	0.18	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
26	0.03	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
27	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
28	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	T	0.00	0.00
29	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.36	0.41	0.00
30	0.00	0.00	0.00	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
31	0.00	0.00	0.37	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Monthly Total	2.04	0.18	1.06	0.03	0.01	0.03	0.07	0.00	0.00	0.39	0.74	0.16
Annual Total	4.71											

T =Trace, some precipitation fell but not enough to measure.






Figure 3-3. Santa Barbara rainfall for 2013.



3.9. Water Quality Materials and Methods

Sampling and data collection for water quality assessment was conducted quarterly at the 13 stations described above. Temperature, conductivity (later converted to salinity), dissolved oxygen, pH, and light transmittance were measured continuously through the water column using a SeaBird 25plus CTD Water Quality Analyzer with associated WetLabs 25-cm Transmissometer. All probes were calibrated immediately prior to each field excursion and, if any data were questionable, they were calibrated again immediately after the instruments were returned to the laboratory. Measurements of light penetration were measured using a Secchi disk. At all stations, water samples were collected at the surface, at mid-depth, and above the bottom with a Nauman sampler.

Water was distributed into sterile 125 mL polypropylene bottles for bacterial analysis. At all stations, temperature and pH were measured directly at the surface using an NBS traceable standard mercury thermometer and hand-held, buffer-calibrated pH meter (respectively). Extra water samples were also collected and set for dissolved oxygen and chloride titration in the field. These extra samples and measurements were used as a check and back up to the water quality analyzer.

All samples from all stations were placed in coolers containing wet ice and were returned to the Ventura laboratory the same day. Immediately upon return, the bacterial samples were set for total and fecal coliform and enterococcus bacteria via multiple-tube fermentation methods. Check samples were titrated for dissolved oxygen by Winkler titration and chloride (converted to salinity) by the argentometric titration. All water analyses were performed in accordance with *Standard Methods for the Examination of Water and Wastewater* (American Public Health Association, 22nd Edition).

After all analyses were completed, the five water quality analyzer variables were correlated against the check samples measured or collected in the field: thermistor probe versus mercury thermometer, conductivity probe versus chloride titration, dissolved oxygen probe versus Winkler titration, field pH probe versus hand-held pH meter, and transmissometer versus Secchi disk (see Appendix Figure 10-1 for calibration curves). The Seabird Water Quality Analyzer was downloaded and water column graphs were generated. Two tables were also prepared containing the results of the physical, chemical, bacterial, and observational water measurements. Check sample correlations, water column graphs, and data tables were joined with a narrative report and were presented to the Water Quality Control Board quarterly. The results and conclusions of all water column measurements and analyses are presented and summarized in Section 3.10 below.



3.10. Results

3.10.1. Physical and Chemical Water Quality

3.10.1.1. Temperature

Coastal water temperatures vary considerably more than those of the open ocean. This is due to the relative shallowness of the water, inflow of freshwaters from the land, and upwelling. Seawater density is important in that it is a major factor in the stratification of waters. The transition between two layers of varying density is often distinct; the upper layer, in which most wind-induced mixing takes place, extends to a depth of 10 to 50 m in southern California waters.

During the winter months, there is little difference in temperature between surface and deeper waters, while in the summer a relatively strong stratification (i.e. thermocline) is evident because the upper layers become more heated than those near the bottom do. Thus, despite little difference in salinity between surface and bottom, changes in temperature during the summer result in a significant reduction of density at the surface. Stratified water allows for less vertical mixing. This is important because bottom waters may become lower in oxygen without significant replenishment from the surface (Soule et. al. 1997).

Spatial temperature patterns. Examination of 3D contours for each quarterly survey showed that the water column was isothermal during January and November. In July temperatures warmed and a moderately strong thermal gradient was established. In November the water column temperatures were warmest of the four surveys and was isothermal (Figure 3-6 and Table 3-3). In January, water temperatures essentially the same through the water column (12.5 °C). The April survey had water temperatures that declined with depth, ranging from 14.3 °C near the surface to 12.1 °C at the bottom. Thermal stratification was strongest in July when water temperatures were ranged from 11.6 to 18.1 °C, representing a 6.5 °C decrease from surface to bottom. In November the water column had the highest average temperatures of the year, but the thermal stratification was weak with temperatures ranging from 16.7 °C at the surface to 15.1 °C near the bottom.

Influences of the outfall were not evident in the temperature profiles during any survey. Temperatures did not correlate with distance from the outfall in any survey. There were no significant temperature differences by t-test between near outfall and far field station groups during the four quarterly surveys.



Receiving Water Environment




Figure 3-6. Temperature contours for the K Station (depth = 18 m) and B Station (depth = 28 m) water quality transects. The Goleta Sanitary District outfall is depicted as a red line. The color legend is presented to the right.



Receiving Water Environment

Table 3-3. Water quality parameter averages and ranges for all stations and depths combined for each quarterly survey. The statistical significance of quarterly measurements with distance from the outfall was tested by correlation analysis and by t-test.

Parameter	Month	Average	Range	Expected & Significant Correlation w/ Outfall?	Significant t-test w/ Outfall?
Temperature	January	12.5	12.4 - 12.7	No	No
	April	13.4	12.1 - 14.3	No	No
	July	15.4	11.6 - 18.1	No	No
	November	16.0	15.1 - 16.7	No	No
Salinity	January	33.5	33.4 - 33.5	No	No
	April	33.6	33.6 - 33.7	No	No
	July	33.6	33.5 - 33.8	No	No
	November	33.6	33.5 - 33.6	No	No
pH	January	8.0	7.7 - 8.3	No	No
	April	8.2	8.0 - 8.3	No	No
	July	8.2	7.9 - 8.3	No	No
	November	8.2	8.1 - 8.3	No	No
DO	January	8.4	7.9 - 8.8	No	No
	April	7.7	5.3 - 8.7	No	No
	July	7.6	4.7 - 8.7	No	No
	November	7.6	6.9 - 8.1	No	No
Transmissance	January	77.2	57.0 - 82.6	No	No
	April	84.6	77.9 - 87.3	No	No
	July	73.3	61.2 - 80.2	No	No
	November	82.5	72.5 - 85.2	No	No
Transparency	January	5.3	4.8 - 6.3	Yes	Yes
	April	11.8	11.5 - 12.5	No	No
	July	7.2	6.0 - 9.0	No	No
	November	10.6	7.5 - 12.0	No	Yes



3.10.1.2. Salinity

Salinity (a measure of the concentration of dissolved salts in seawater) is relatively constant throughout the open ocean; however, it can vary in coastal waters primarily because of the inputs of freshwater from the land or because of upwelling. In a five-year study conducted by the U.S. Navy Research and Development Center, more than 1000 samples were analyzed for salinity. The mean salinity was 33.75 parts per thousand (ppt), and the range of 90% of the samples in southern California fell between 33.57 and 33.92 ppt (SCCWRP 1973).

Despite the general lack of variability, salinity concentrations can be affected by a number of oceanographic factors. During spring and early summer months, northwest winds are strongest and drive surface waters offshore. Deeper waters, which are colder, more nutrient-rich, and more saline, are brought to the surface to replace water driven offshore (Emery 1960). El Nino (ENSO) events can also affect coastal salinities. During these events northern flowing waters move into the Bight with waters that are also more saline, but are warmer and lower in nutrients than ambient water. Major seasonal currents (i.e. California current, countercurrent, or undercurrent) can also affect ambient salinity to some degree (Soule et. al. 1997).

Spatial salinity patterns. Average salinity in the survey area was nearly identical across the four surveys ranging from 33.5 ppt in January to 33.6 ppt in each of the other surveys. However, salinity provided the best opportunity to detect the effluent plume which is evident in the April and November surveys. In January, the water column was isohaline (same surface to bottom). In April, lower salinity water is seen as a surface and subsurface lens of slightly fresher water both to the west of the outfall. In July when the water column was most strongly thermally stratified, salinity was layered through the water column with no clear indication of the plume. Finally, in November a lens of fresher water was detected at the end of the outfall and stretched past station B3 to the west.

Salinity ranges and outfall effects. Table 3-3 shows the range of salinities for the 11 water column stations over the four quarterly sampling surveys. Salinities did not correlate with distance from the outfall and there were no significant salinity differences by t-test between near outfall and far field station groups.



Receiving Water Environment




Figure 3-7. Salinity (ppt) contours for the K Station (depth = 18 m) and B Station (depth = 28 m) transects. The Goleta Sanitary District outfall is depicted as a red line.



3.10.1.3. Hydrogen Ion Concentration (pH)

pH is defined as the negative logarithm of the hydrogen ion concentration. A pH of 7.0 is neutral, values below 7.0 are acidic, and those above 7.0 are basic (Horne 1969). Seawater in southern California is slightly basic, ranging between 7.5 and 8.6, although values in shallow open-ocean water are usually between 8.0 and 8.2 (SWQCB 1965). These narrow ranges are due to the strong buffering capacity of seawater, which rarely allows for extremes in pH.

Factors that can influence pH in the ocean are freshwater inputs, upwelling, and biological activity. Since freshwater pH values tend to be about 0.5 pH units less than seawater, any inflow from a freshwater source will tend to lower the pH slightly. When photosynthesis is greater than respiration, more carbon dioxide is taken up than generated, and pH may increase to higher values in the euphotic (i.e. light penetrating) zone. When respiration is greater than photosynthesis, more carbon dioxide is released than used and pH may decrease, especially when mixing is minimal such as in the oxygen minimum zone and towards the bottom (Soule et. al. 1997).

Spatial pH patterns. Average pH across the four quarterly surveys ranged from 7.7 to 8.3 (Figure 3-8 and Table 3-3). In January the pH sensor failed intermittently making evaluation of data difficult.

In April and July pH was stratified through the water column and was least near the bottom (8.0 and 7.9, respectively) and greatest near the surface (8.3, each). In November, pH was similar through the water column. There was no clear evidence of the effluent plume from the contours during any of the four surveys.

pH ranges and outfall effects. Table 3-3 shows the range of pH values for 11 water column stations for each of the four quarterly sampling surveys. There were no expected and significant correlations with distance to the outfall for any survey. Also, there were no significant differences in pH among station groups located near and far from the outfall by t-test for any survey. Analysis of each quarterly data set showed that all pH differences between stations near and away from the outfall were very low and well within the 0.2 pH unit limit specified in the California Ocean Plan (2009).



Receiving Water Environment




Figure 3-8. pH contours for the K Station (depth = 18 m) and B Station (depth = 28 m) transects. The Goleta Sanitation District outfall is depicted as a red line.



3.10.1.4. Dissolved Oxygen

The most abundant gases in the ocean are oxygen, nitrogen, and carbon dioxide. These gases are dissolved in seawater and are not in chemical combination with any of the materials composing seawater. Gases are dissolved from the atmosphere by exchange across the sea surface. The gases dissolved at the sea surface are distributed by mixing, advection (i.e. from currents), and diffusion. Concentrations are modified further by biological activity, particularly by plants and certain bacteria. In nature, gases dissolve in water until saturation is reached given sufficient time and mixing. The volume of gas that saturates a given volume of seawater is different for each gas and depends upon temperature, pressure, and salinity. An increase in pressure, or a decrease in salinity or temperature, causes an increase in gas solubility.

The amount of oxygen dissolved in the sea varies from zero to about 11 milligrams per liter. At the surface of the sea, the water is more or less saturated with oxygen because of the exchange across the surface and plant activity. In fact, when photosynthesis is at a maximum during a phytoplankton bloom, such as during a red tide event, it can become supersaturated (Anikouchine and Sternberg 1973). When these blooms die off, bacterial aerobic respiration during decomposition of these phytoplankton cells can rapidly reduce dissolved oxygen in the water. Dissolved oxygen typically decreases with depth due to respiration associated with the bacterial breakdown of organic material. However, if the water column is well mixed, oxygen will be fairly constant with depth. Temperature and/or salinity can affect the density structure of the water column and create barriers to vertical mixing.

Spatial oxygen patterns. During the January survey, dissolved oxygen concentrations ranged from 4.7 to 8.8 mg/L, were similar from surface to bottom in January and November, and less near the bottom in April and July (Figure 3-9 and Table 3-3). In April and July the water column was stratified for oxygen and ranged from 5.3 and 4.7 mg/L, near the bottom to 8.8 mg/L near the surface. This was clearly the result of upwelled, oxygen depleted deep water coming onshore and supports the decreases in temperature, pH and salinity discussed in previous sections.

Oxygen ranges and outfall effects. Table 3-3 shows the range of oxygen concentrations for the 11 water column stations over the four quarterly sampling surveys. Dissolved oxygen did not correlate significantly with distance to the outfall for any of the four surveys and there were no significant differences by t-test among sites located near the outfall and those further away. This indicates that dissolved oxygen was not influenced by the outfall diffuser. Dissolved oxygen concentrations between stations located near and away from the outfall remained within the Ocean Plan standards (2009) throughout the year, except in July when dissolved oxygen was depleted between the plume stations (8.29 mg/L) and sites B2, B3, B4 and B5 (range = 6.82 to 7.30 mg/L). These differences represented a 13% to 23% reduction in dissolved oxygen. It is most likely that the depressed oxygen offshore was due to upwelling since the plume sites (WCZID and WC100) had greater dissolved oxygen.



Receiving Water Environment




Figure 3-9. Dissolved oxygen contours for the K Station (depth = 18 m) and B Station (depth = 28 m) transects. The Goleta Sanitary District outfall is depicted as a red line.



3.10.1.5. Light Transmissance

Water clarity in the ocean is important both for aesthetic and ecological reasons. Phytoplankton, as well as multicellular marine algae and flowering plants are dependent upon light for photosynthesis and therefore growth. Since nearly all higher-level organisms are dependent upon plants for survival (except those animals living in deep-ocean volcanic vents and similar environments), the ability of light to penetrate into the ocean depths is of great importance. Seasonally, water is usually least clear during spring upwelling and winter rain. In early summer, increased day length can promote plankton growth and reduce water clarity, as well. In late summer and fall, days are shorter and the rains that bring sediments into the marine environment have yet to begin. Therefore, late summer and early fall are typically the periods of greatest water clarity. Anthropogenic influences such as wastewater effluents, storm drainage discharges, and non-point runoff can also influence water quality on a local basis.

Water clarity is determined using two completely different measuring techniques. Surface transparency is measured using a weighted, white plastic, 30 cm diameter disk (called a Secchi Disk) attached to a marked line. The disk is simply lowered through the water column until it disappears, and the depth of its disappearance is recorded. Surface transparency is a good estimate of the amount of ambient light that is available to plankton since the depth to which light is available for photosynthesis is generally considered to be about 2.5 times the Secchi disk depth.

Light transmissance is measured using a transmissometer, which is a 0.25 m open tube with an electrical light source at one end and a sensor at the other. The amount of light that the sensor receives is directly dependent upon clarity of the water between them. Results are recorded as percent light transmissance. Since transmissance is independent of ambient sunlight, it can be used at any depth and under any weather conditions. Surface light transmissance is usually positively correlated with surface transparency.

Spatial transmissance patterns. Water clarity was good throughout the water column during each of the four quarterly surveys (Figure 3-10). Average transmissance across the four surveys ranged from 73.3% in January to 84.6% in April (Table 3-3). In addition, clarity was similar with depth in each survey (range = 57% to 78%). In July the 3D contours show what may be the effluent plume as a slightly clearer patch of water spreading west and east of the outfall diffuser at 40 meters. In November there was a slightly less clear patch of water near the terminus of the outfall and further to the east near reference station B6.

Transmissance ranges and outfall effects. Table 3-3 shows the range of transmissance for the 11 water column stations over the four sampling surveys. Comparisons among stations showed there was no significant correlations with distance to the outfall or a significant difference among near and far field stations by t-test during any of the four surveys. In all cases, there was never a reduction in transmissance between near and far field stations that exceeded the Ocean Plan (2009) standard of 10%.



Receiving Water Environment




Figure 3-10. Transmissivity (%) contours for the K Station (depth = 18 m) and B Station (depth = 28 m) transects. The Goleta Sanitary District outfall is depicted as a red line.



3.10.1.6. Surface Transparency

As discussed in more detail in Section 3.10.1.5 above, surface transparency is recorded as the depth (m) at which a weighted, 30 cm, white plastic disk (Secchi Disk) disappears from view. Since only a single quarterly measurement is taken at each station, these data are presented as a line plot of transparency vs. quarter.

Transparency patterns and outfall effects. Figure 3-11 shows the range of transparency measurements for the 11 water column stations over the four sampling surveys. Average surface transparency ranged from 5.3 m in January to 11.8 m in April. The lowest transparency of the year in January coincided with the largest rainfall event of the year (1.5 inches) that was ongoing during sampling. Runoff from this event probably decreased surface transparency. Transparency correlated significantly with distance from the outfall in January and significantly by t-test among stations located near to the outfall compared to stations further away in January and November.




Figure 3-11. Average transparency vs. season for each of the 11 water quality stations.



3.10.2. Bacterial Water Quality

The three bacterial measurements of total coliforms, fecal coliforms and enterococcus, are used by health authorities to assess the potential risk of human exposure to pathogens in the aquatic environment (Soule 1997). The principle problem with these indicators is that analysis takes 72 hours, slowing the response of health officials to potentially hazardous conditions. Research has been underway to develop more rapid tests that are both sensitive and cost effective. Rainfall episodes have been closely associated with violations of all three bacterial standards, especially near areas where creeks or stormwater channels discharge into the ocean. At present, it is more prudent to post areas of potential or known contamination immediately following rain storm events than to wait for confirmation. Bacterial results are summarized in Tables 3-4 and 3-5.

3.10.2.1. Total Coliforms

Coliform bacteria (those inhabiting the colon) have been used for many years as indicators of fecal contamination; they were initially thought to be harmless indicators of pathogens at a time when waterborne diseases such as typhoid fever, dysentery and cholera were severe problems. Recently it was recognized that coliforms themselves might cause infections and diarrhea. However, the total coliform test is not effective in identifying human contamination because these bacteria may also occur as free living in soils, and are present in most vertebrate fecal material. The California Ocean Plan (SWRCB 2009) states that within 1,000 feet of shore, the single sample total coliform concentration cannot exceed 10,000 MPN/100 mL of water. Additionally, during a 30-day period the average concentrations cannot exceed 1,000 MPN/100 mL. Although no offshore stations are within 1000 feet of shore, this value was used as a criterion of concern.

Total coliform patterns over the year. Total coliform counts were very low during the year, ranging from <2 to 50 MPN/100 mL for all surveys (Table 3-4). In general values were very low throughout the year at all stations and depths with most samples below detection (<2 MPN/100 mL). These total coliform concentrations were far below either the single sample Ocean Plan standard (2009) of 10,000 MPN/100 mL or the monthly average total coliform standard of 1,000 MPN/100mL (Table 3-5).



Receiving Water Environment

Table 3-4. Annual summary of total and fecal coliforms and enterococcus bacteria (MPN/100 mL).

Sampling Station	Season	Offshore						Plume		Nearshore				
		B1	B2	B3	B4	B5	B6	WCZID	WC100	K1	K2	K3	K4	K5
SURFACE														
Total Coliform	Winter	5	2	5	<2	<2	<2	<2	23	<2	<2	<2	<2	<2
	Spring	<2	<2	2	<2	<2	<2	<2	<2	<2	<2	2	<2	<2
	Summer	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2
	Fall	<2	<2	<2	2	<2	<2	<2	<2	<2	<2	<2	<2	<2
Fecal Coliform	Winter	5	2	5	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2
	Spring	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2
	Summer	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2
	Fall	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2
Enterococcus	Winter	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2
	Spring	<2	<2	<2	<2	<2	2	<2	<2	<2	<2	<2	<2	2
	Summer	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2
	Fall	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2
MIDDLE														
Total Coliform	Winter	<2	2	2	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2
	Spring	<2	<2	<2	<2	20	2	20	<2	<2	<2	<2	<2	<2
	Summer	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2
	Fall	<2	<2	<2	50	<2	<2	<2	<2	<2	<2	<2	<2	<2
Fecal Coliform	Winter	<2	<2	2	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2
	Spring	<2	<2	<2	<2	20	2	20	<2	<2	<2	<2	<2	<2
	Summer	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2
	Fall	<2	<2	<2	20	<2	<2	<2	<2	<2	<2	<2	<2	<2
Enterococcus	Winter	<2	<2	<2	<2	2	<2	<2	<2	<2	<2	<2	<2	<2
	Spring	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2
	Summer	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2
	Fall	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2
BOTTOM														
Total Coliform	Winter	2	20	<2	8	20	50	<2	2	<2	20	<2	<2	<2
	Spring	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2
	Summer	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2
	Fall	<2	<2	<2	<2	<2	2	<2	<2	<2	20	<2	<2	<2
Fecal Coliform	Winter	2	20	<2	8	20	2	<2	2	<2	20	<2	<2	<2
	Spring	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2
	Summer	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2
	Fall	<2	<2	<2	<2	<2	2	<2	<2	<2	<2	<2	<2	<2
Enterococcus	Winter	<2	2	<2	<2	<2	2	<2	<2	<2	<2	<2	<2	<2
	Spring	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2
	Summer	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2
	Fall	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2	<2



Table 3-5. Indicator bacteria geometric averages and ranges for all stations and depths combined for each quarterly survey. Measurements for the year were compared individually against single sample event, REC-1 bathing water standards.

Parameter	Month	Average	Range	Water Quality Standard	Standard Exceedances
Total Coliform	January	5	<2 - 50	10,000	0
	April	3	<2 - 20	10,000	0
	July	2	<2 - 2	10,000	0
	October	4	<2 - 50	10,000	0
Fecal Coliform	January	4	<2 - 20	400	0
	April	3	<2 - 20	400	0
	July	2	<2 - 2	400	0
	October	3	<2 - 20	400	0
Enterococcus	January	2	<2 - 2	104	0
	April	2	<2 - 2	104	0
	July	2	<2 - 2	104	0
	October	2	<2 - 2	104	0

3.10.2.2. Fecal Coliforms

The fecal coliform test discriminates primarily between soil bacteria and those in warm blooded animals such as dogs, cats, birds, horses, barnyard animals, and humans. The California Ocean Plan (SWRCB 2009) states that within 1000 feet of shore, samples from each station shall have a density of fecal coliform organisms less than 400 MPN/100 mL of water for any single sample or average less than 200 for any 30 day period. Although no offshore stations are within 1000 feet of shore, this value was used as a criterion of concern.

Fecal coliform patterns over the year. Fecal coliform counts were very low during the year, ranging from <2 to 20 MPN/100 mL for all surveys (Table 3-4). In general values were very low throughout the year at all stations and depths with most samples below detection (<2 MPN/100 mL). These fecal coliform concentrations were far below either the single sample Ocean Plan standard (2009) of 400 MPN/100 mL or the monthly average fecal coliform standard of 200 MPN/100mL (Table 3-5).

3.10.2.3. Enterococcus

Enterococcus bacteria include species that are found in human wastes and are related to the Streptococcus bacteria. At one time they were believed to be exclusive to humans, but other Streptococcus species occur in feces of cows, horses, chickens, and other birds. Enterococci die off rapidly in the environment, making



them indicators of fresh contamination, but not exclusively from humans. The California Ocean Plan (SWRCB 2009) limitations within 1000 feet of shore are a 30 day average of 34 MPN/100 mL and a single sample limit of 104 MPN/100 mL.

Enterococcus bacteria patterns over the year. Enterococcus bacteria counts ranged from the method detection limit (<2 MPN/100 mL) to just above it (2 MPN/100 mL) during each survey (Table 3-4). Enterococcus concentrations at all stations and depths in the survey area were below the single sample Ocean Plan standard (2009) of 104 MPN/100 mL (Table 3-5).

3.11. Discussion

Quarterly water quality surveys were conducted offshore Goleta in January, April, July and November 2013. Measurements for temperature, salinity, pH, dissolved oxygen and water clarity showed that oceanographic conditions during the year were typical of nearshore areas in southern California. Rainfall for this period (4.71 inches) was 13.57 inches less than the average yearly rainfall since 1981 (18.96 inches). This lack of rainfall meant less nearshore surface runoff and may have led to the good water clarity and low bacteria counts throughout the year.

Salinity provided the best opportunity to detect the effluent plume which was evident in the April and November monthly contours. In January, lower salinity water was seen as a surface and subsurface lens of slightly fresher water both to the north and south of the outfall. In April, lower salinity water is seen as a surface and subsurface lens of slightly fresher water both to the north of the outfall. In November a lens of fresher water was detected at the end of the outfall and stretched past station B3 to the north. The depth of the plume in April and November are presumably due to the presence of a weak thermal gradient which held the buoyant freshwater plume beneath it. None of the other parameters showed evidence of the effluent plume.

Physical and chemical characteristic restrictions, which apply to waters outside of the zone of initial dilution, are addressed in the California Ocean Plan (2009):

- *The pH shall not be changed at any time more than 0.2 units from that which occurs naturally.*
- *The dissolved oxygen concentration shall not at any time be depressed more than 10 percent from that which occurs naturally, as the result of the discharge of oxygen demanding waste materials.*
- *Natural light shall not be significantly reduced at any point outside of the zone of initial dilution.*
- *Floating particulates and grease and oil shall not be visible.*
- *The discharge of waste shall not cause aesthetically undesirable discoloration of the ocean surface.*
- *Waste discharged to the ocean must be essentially free of: 1) Material that is floatable or will become floatable upon discharge.*



- *The waste discharged to the ocean must be essentially free of: 4) Substances that significantly decrease the natural light to benthic communities and other marine life.*

- *Waste discharged to the ocean must be essentially free of: 5) Materials that result in aesthetically undesirable discoloration of the ocean.*

The water quality parameters measured during the four quarterly surveys indicated that the outfall plume was not altering the condition of the water mass in the vicinity of the Goleta outfall. None of the above restrictions were exceeded outside the zone of initial dilution. Water color throughout the area was green, discharge related oil or floating particulates were never observed in the survey area. Water quality measurements taken near to and far from the outfall terminus did not correlate expectedly and significantly with distance from the outfall, except for transparency in January. In addition, in January and in November transparency was significantly different among sites close to and far from the outfall. While statistically significant, these differences were small and not ecologically significant:

1. In January transparency differences from near field to far field stations were exceedingly small, with the average difference among the plume stations (4.9 m) and the station with the greatest transparency (far field station B6, 6.3 m) a 1.4 m difference. In addition, January sampling occurred during the largest rainfall event of the year (1.5 inches). Surface runoff from this event was probably the cause of the small decrease in surface transparency.
2. In November transparency differences from near field to far field stations were also exceedingly small, with the average difference among the plume stations (10.7 m) and the station with the greatest transparency (outfall station B4, 12.0 m) a 1.3 m difference.

Dissolved oxygen, pH and transmissance were within Ocean Plan (2009) standards during each of the four quarterly surveys. The only exception to this was for oxygen in July when dissolved oxygen was depleted between the plume stations (8.29 mg/L) and sites B2, B3, B4 and B5 (range = 6.82 to 7.30 mg/L). These differences represented a 13% to 23% reduction in dissolved oxygen. The Ocean Plan limits reductions in dissolved oxygen between sites near the ZID and those further away to 10%. It is most likely that the depressed oxygen offshore was due to upwelling since the plume sites (WCZID and WC100) had greater dissolved oxygen.

Bacteriological standards are addressed in the Ocean Plan and NPDES discharge permit, however these standards relate primarily to shoreline waters used for recreation or shellfish harvesting (REC-1 bathing water standards). Total coliforms, fecal coliforms and enterococcus indicator bacteria concentrations were very low throughout the year in the Goleta survey area. A total of 156 samples were collected and analyzed for each indicator. None of these exceeded the single sample Ocean Plan standard (2009) during the year and over 95% of the measurements were below detection limits.

In conclusion, evidence from the four quarterly water column monitoring surveys conducted in 2013 indicate that the Goleta Sanitary District Wastewater Treatment Plant was in compliance with all water quality standards, and that the treatment plant was operating effectively.



CHAPTER 4

Physical Characteristics of the Benthic Sediments

4.1. Background

Marine sediments provide clues to the nature of the environment from which their constituent materials were derived, the transportation processes by which they arrived at the final site of deposition, and the physico-chemical and biological characteristics of the depositional environment. The Southern California Bight coastal shelf is characterized by sediments composed of varying combinations of sand, silt and clay. This is quite different in character from more northerly coastal reaches that are composed of rocky substrates. The distribution of benthic sediments can have a profound effect upon the diversity, abundance, and community structure of infaunal organisms and the accumulation of organic material and anthropogenic contaminants (Gray 1981). In general, finer sediments provide a more stable environment for benthic organisms, especially those that build tubes, burrow and feed there. Finer sediments, however, also tend to adsorb more organic and elemental contaminants than do coarser, sandier sediments. As a result, organisms that live closely associated with fine sediments can be exposed to higher concentrations of contaminants.

4.2. Materials and Methods

Benthic grab sampling was conducted in accordance with *Techniques for Sampling and Analyzing the Marine Macrobenthos* March 1978, EPA 600/3-78-030; *Quality Assurance and Quality Control (QA/QC) for 301 (h) Monitoring Programs: Guidance on Field and Laboratory Methods* May 1986, Tetra Tech; *The Southern California Bight Pilot Project Field Operations Manual* (SCCWRP 2008).

Samples were collected with a chain-rigged, tenth square-meter Van Veen Grab. At each station, the grab was lowered rapidly through the water column until near bottom, and then slowly lowered until contact was made. The grab was then slowly raised until clear of the bottom. Once on board, the grab was drained and initial qualitative observations of color, odor, consistency, etc. were recorded.

Sediments to be analyzed for physical properties were removed from the top 2 cm of the surface and placed in clean plastic Whirl-Pacs. These were analyzed for particle size distribution using a Horiba LA920 Particle Size Analyzer and in accordance with Standard Methods 2560 D (APHA, 2012). Sub-samples from each sediment sample were re-suspended in de-ionized water, and then injected into the analyzer. The analyzer is capable of measuring particle sizes ranging from silt and clay (<2 μm) up to coarse sand (2,000 μm). Results were recorded as the percentage each size distribution represented of the whole. When the LA920 detected particles in a sample that neared its upper detection limit (2,000 μm), a portion of the sample was dried at 105 °C, weighed, then sieved through a 2,000 μm mesh screen. Particles not passing through the screen were weighed and expressed as the percentage of particles in the sample >2,000 μm (gravel).

Data for each station were reduced to the median particle size (μm), percent fines and, the sorting index. The sorting index values range between sediments that have a very narrow distribution (very well sorted) to those which have a very wide distribution (extremely poorly sorted). This index is simply calculated as the 84th percentile minus the 16th percentile divided by two (Gray 1981). Well sorted sediments are homogeneous and are



typical of high wave and current activity (high energy areas), whereas poorly sorted sediments are heterogeneous and are typical of low wave and current activity (low energy areas).

4.3. Results

4.3.1. Station Event and Sea State Conditions

Sediment sampling, trawling and mussel retrieval was conducted on October 15th, 2013 under clear skies, and calm to moderate conditions (Table 4-1). Wave height was two feet from the southwest and winds were three knots from the northeast.

4.3.2. Particle Size Distribution

Tables 4-2 and 4-3, and Figure 4-1 illustrate the overall particle size distributions from the six sediment-sampling stations. Detailed raw and summary data for particle size are presented in Appendix 10.3. Results are presented for each size range as the percent of the whole. Two sediment characteristics can be inferred from the graphs. Position of the midpoint of the curve will tend to be associated with the median particle size (Figure 4-1). If the midpoint tends to be toward the larger micron sizes, then it can be assumed that the sediments will tend to be coarser overall. If the midpoint is near the smaller micron sizes, then it can be assumed that the sediments are mostly finer. Sediment sizes that range from 2000 to 63 μm are defined as sand, sediments ranging from 63 to 4 μm are defined as silt, and sediments that are 4 μm or less are defined as very fine silt and clay (Wentworth Sediment Scale, see Gray 1981). There are also subdivisions within the categories (e.g. very fine sand, etc., see Table 4-3). A second pattern discernible from the graph is how homogeneous the distributions of sediments are. Sediments that tend to have a narrow range of sizes are considered homogeneous or well sorted. Others, which have a wide range of sizes, are considered to be heterogeneous or poorly sorted.

4.3.2.1. General Description

A total of 36 replicate samples were successfully collected at the six sampling sites for all biological and chemical analyses (Table 4-2). The penetration depth of each grab exceeded the 5 cm minimum depth required by the Southern California Bight protocol. Surface sediments had the same descriptions at all stations. Surface sediments were composed of fine sand, the color was olive green and there was no odor.

4.3.2.2. Median Particle Size

Median particle sizes are depicted in Table 4-3. Similar to past years, median particle sizes were categorized as very fine sand, except at B1 which was fine sand. Median particle sizes ranged from 91 to 131 μm . Stations B1 and B4 had the greatest median particle sizes of all sites (122 and 131, respectively).



4.3.2.3. Sorting Index & Percent Fines

Particles at all stations were poorly sorted and sorting indexes ranged from 1.08 at station B4 to 1.73 at B5 (Table 4-3). The percent fine sediments ranged from 17% at station B4 to 26% at station B5 near the Goleta Outfall.

4.4. Discussion

Observational and analytical evaluations of the benthos in the vicinity of the Goleta outfall show that the sediments are heterogeneous and composed of very fine and fine sand. The percentage of fine sediments (silt and clay) ranged from 17% to 26% at each of the stations, which was in keeping with results from previous years. Hydrogen sulfide gas is a byproduct of bacterial decomposition of organic material under anoxic conditions. In 2012 the smell of hydrogen sulfide was present in two replicates at station B1, however, this year the smell was not present in any of the sediments collected.

There were no apparent differences in particle size between the outfall stations and those further away. Evidence from this analysis suggests that the discharge is not contributing finer particles to the benthos near the outfall terminus.



Table 4-1. Goleta Sanitary District locations, survey information and weather conditions during the sediment and trawling survey.

Stations	B1	B2	B3	B4	B5	B6	TB3	TB6
Date	15-Oct-13							
Time	10:04	9:39	9:18	8:56	8:25	7:52	11:22	2:05
Research Vessel	<i>Hey Jude</i>							
Survey Program	Benthic Sediment	Trawl, Bioaccum.	Trawl, Bioaccum.					
Dist. From Outfall (m)	1500	500	250	25	25	3000	250	3000
Dir. From Outfall (°M)	270	270	270	270	90	90	270	90
Depth (m)	27.0	27.0	27.0	27.0	27.0	29	22.8	25.9
Latitude (N)	34.58261	34.40192	34.40192	34.40192	34.40197	34.40283	34.40247	34.40063
Longitude (W)	119.84103	119.83069	119.82792	119.82547	119.82492	119.79269	119.83282	119.78075
Weather	Clear							
Tide	Outgoing	Incoming						
Swl. Ht. (ft)	2	2	2	2	2	2	2	2
Swl. Dir.	SW							
Wind Sp. (Kn)	3	3	3	3	3	3	3	3
Wind Dir.	NE	NE	NE	NE	NE	NE	NW	SW



Table 4-2. Sediment grab descriptions.

Station	Rep	Penetration (cm)	Surface Description	Surface Color	Odor	Analysis
B1	1	9.0	Fine Sand	Olive Green	None	Biology
B1	2	10.0	Fine Sand	Olive Green	None	Biology
B1	3	9.0	Fine Sand	Olive Green	None	Biology
B1	4	8.0	Fine Sand	Olive Green	None	Biology
B1	5	9.0	Fine Sand	Olive Green	None	Biology
B1	6	7.0	Fine Sand	Olive Green	None	Chemistry
B2	1	8.0	Fine Sand	Olive Green	None	Biology
B2	2	10.0	Fine Sand	Olive Green	None	Biology
B2	3	5.0	Fine Sand	Olive Green	None	Chemistry
B2	4	9.0	Fine Sand	Olive Green	None	Biology
B2	5	8.0	Fine Sand	Olive Green	None	Biology
B2	6	8.0	Fine Sand	Olive Green	None	Biology
B3	1	9.0	Fine Sand	Olive Green	None	Biology
B3	2	10.0	Fine Sand	Olive Green	None	Biology
B3	3	8.0	Fine Sand	Olive Green	None	Biology
B3	4	8.0	Fine Sand	Olive Green	None	Biology
B3	5	8.0	Fine Sand	Olive Green	None	Biology
B3	6	8.0	Fine Sand	Olive Green	None	Chemistry
B4	1	10.0	Fine Sand	Olive Green	None	Biology
B4	2	10.0	Fine Sand	Olive Green	None	Biology
B4	3	8.0	Fine Sand	Olive Green	None	Biology
B4	4	8.0	Fine Sand	Olive Green	None	Biology
B4	5	8.0	Fine Sand	Olive Green	None	Biology
B4	6	7.0	Fine Sand	Olive Green	None	Chemistry
B5	1	8.0	Fine Sand	Olive Green	None	Biology
B5	2	8.0	Fine Sand	Olive Green	None	Biology
B5	3	8.0	Fine Sand	Olive Green	None	Biology
B5	4	9.0	Fine Sand	Olive Green	None	Biology
B5	5	12.0	Fine Sand	Olive Green	None	Biology
B5	6	9.0	Fine Sand	Olive Green	None	Chemistry
B6	1	9.0	Fine Sand	Olive Green	None	Biology
B6	2	10.0	Fine Sand	Olive Green	None	Biology
B6	3	9.0	Fine Sand	Olive Green	None	Biology
B6	4	9.0	Fine Sand	Olive Green	None	Biology
B6	5	8.0	Fine Sand	Olive Green	None	Biology
B6	6	8.0	Fine Sand	Olive Green	None	Chemistry

Table 4-3. Grain size characteristics of each Goleta station.

Station	Median (microns) ^{1.}	Category	Sorting Index ^{2.}	Sorting	% Fines
B1	131	fine sand	1.58	poorly sorted	20
B2	91	very fine sand	1.43	poorly sorted	23
B3	105	very fine sand	1.11	poorly sorted	18
B4	122	very fine sand	1.08	poorly sorted	17
B5	94	very fine sand	1.73	poorly sorted	26
B6	109	very fine sand	1.21	poorly sorted	18

1. 0-4 = clay, 4-8 = very fine silt, 8-16 = fine silt, 16-31 = medium silt, 31-63 = coarse silt, 63-125 = very fine sand, 125-250 = fine sand, 250-500 = medium sand, 500-1000 = coarse sand.

2. <0.35 = very well sorted, 0.35-0.50 = well sorted, 0.50-0.71 = moderately well sorted, 0.71-1.00 = moderately sorted, 1.0-2.0 = poorly sorted, 2.0-4.0 = very poorly sorted, >4.0 = extremely poorly sorted.






Figure 4-1. Particle size frequency (%) at each station in the Goleta survey area.



CHAPTER 5

Chemical Characteristics of Sediments

5.1. Background

Sources of potential contaminants discharged into the Southern California Bight include treated municipal and industrial wastewater, storm water runoff from urbanized areas, disposal of dredged materials, aerial fallout, oil and hazardous material spills, boating and other sources. Bottom sediments are often the fate of these contaminants, where they can reside for long periods of time, exerting effects at various levels of biological organization (SCCWRP 1998). Organic and metal contaminants tend to adsorb more readily on finer particles and can thus accumulate in areas of deposition. This accumulation of contaminants can impact resident organisms living both within the sediments and on the surface.

5.2. Materials and Methods

Field sampling for all benthic sediment components is described in Chapter 4, Section 4.2, Materials and Methods. Single sediment grabs were collected at stations B1 through B6 (Figure 5-1). Sediment portions to be chemically analyzed were removed from the top two centimeters of the grab sample with a stainless steel spatula and placed in pre-cleaned glass bottles with Teflon-lined caps. During all collections, the sides of the grab were avoided. Samples were immediately placed on ice and returned to the laboratory. PHYSIS Environmental Laboratories, located in Anaheim, California, performed all chemical analyses. Results were standardized to $\mu\text{g/g}$ dry weight for undifferentiated organics and metals and $\mu\text{g/Kg}$ dry weight for complex organics.

Since replicate field samples are not required, results were correlated against distance from the outfall diffuser. When appropriate, correlations were designated as significant ($p \leq 0.05$) or marginally significant ($0.05 < p \leq 0.10$, see Section 3.5.) and expected (negative) or unexpected (positive) (see Section 3.5.1). Since grain size can have an important effect on the ability of contaminants to adhere to particles, results were also correlated against percent fine particle size. The expected sign for particle size would be negative (increasing concentrations with smaller size).

As described in (Section 4.4.), areas west of the diffuser are known sources of natural oil seepages; therefore, results were also correlated against distance from Goleta Point. Like distance from outfall, the expected sign would be negative. Spearman's correlation was used to assess spatial trends (see Sokal and Rohlf 1981).

In order to determine long-term trends, 2013 data were compared to results from monitoring surveys that began in 1991 (Brown and Caldwell 1992, 1993, 1994, 1995, 1996, 1997, 1998; Aquatic Bioassay 1999 to 2012). Data were also compared to results of "reference" sediments from uncontaminated areas collected and analyzed by the Southern California Bight Regional Monitoring Program (SCBRMP) in 1998, 2003 and 2008. Finally, results were compared to the limits presented in two NOAA studies (NOAA 1990 and Long, et. al. 1995). In these studies, researchers compiled published information regarding the toxicity of chemicals to benthic organisms. The data for each compound were sorted, and the lower 10th percentile and median (50th) percentile were identified. The lower 10th percentile in the data was identified as an Effects Range-Low (ER-L) and the median was identified as an Effects Range-Median (ER-M).



Per the NPDES permit, all contaminants were “normalized” to percent fine sediments and percent total organic carbon (TOC) at each station. NOAA scientists have determined that normalizing data from sediments that contain less than 20% silt and clay can cause erroneously high results; therefore, results from samples containing less than 20% fine components should be viewed with caution (NOAA 1990).

5.3. Results

Table 5-1 lists all of the chemical constituents measured from samples collected at each of the six benthic sediment stations. These compounds have been separated here into three main groups: undifferentiated organic compounds, heavy metals, and complex organic compounds. Complex organic compounds are further divided into chlorinated pesticides, polychlorinated biphenyls (PCB's), and polynuclear aromatic hydrocarbons (PAH's). Appendix tables 10-4 and 10-5 present data normalized to percent fine sediments (silt and clay fractions) and percent TOC. Appendix table 10-6 lists the constituents minimum detection limits (MDL), reporting limits (RL) and methods. Figure 5-2 shows the average (\pm standard deviation) concentration for all Goleta stations combined, for each constituent measured from 1991 to present. Tables 5-4 and 5-5 compare the Goleta sediment chemistry results with the 1998, 2003 and 2008 SCBRMP surveys and the NOAA ER-L and ER-M values.

5.3.1 Undifferentiated Organics

The undifferentiated organics discussed in this report includes groups of compounds whose concentrations can help to determine the extent of anthropogenic contaminant loading in an area. These groups are discussed below:

- Total organic carbon (TOC) is a measure of the amount of carbon derived from plant and animal sources. It is a better measure of the portion of a sample derived from these sources than is percent volatile solids (Soule et al. 1996).
- Sources of oil and grease can be attributed to storm water runoff and ocean going vessels. The extent that people dump used motor oil into storm drains is unknown. Also, the Goleta outfall is located in an area of natural oil seeps, which may be a natural source.
- Total Kjeldahl Nitrogen (TKN) is the method used for the measure of organic nitrogen in water and sediments. Organic nitrogen is present due to the breakdown of animal products and includes such natural materials as proteins and peptides, nucleic acids, urea, and numerous synthetic organic materials (APHA 1995).
- Acid volatile sulfide (H_2S) is an indicator of organic decomposition occurring particularly in anoxic sediments and characterized by a rotten egg smell. No sediment reference values are available for sulfides.

5.3.1.1 Undifferentiated Organics Spatial Patterns

The concentrations for each of the undifferentiated organics measured for this survey are listed in Table 5-1. Similar to 2012, the concentrations of oil and grease were greatest at Station B1 offshore Goleta Point (814 mg/L) and decreased at stations nearest to the outfall until the lowest concentration was measured at station B6 (184 mg/Kg). Total Kjeldahl nitrogen (TKN) concentrations were greatest near the outfall (B5) and least at station B6, 3,000 meters east of the outfall. TOC concentrations were least at station B3 (200 mg/L) and greatest at station B5 (5,100 mg/L) nearest the outfall (B5). Acid volatile sulfide (AVS)



was greatest at station B1 near Goleta Point (172.23 mg/L) and decreased to lowest concentrations at station B4 (4.11 mg/L) near the outfall.

Each undifferentiated organic correlated expectedly (decreased) with distance from the outfall, except oil and grease. None of the correlations with distance to the outfall were significant. All of the undifferentiated organics, except TOC, correlated expectedly with distance to Goleta Point; oil and grease significantly so. Each undifferentiated organic constituent correlated unexpectedly (increased with increasing particle size) and non-significantly with sediment particle size.

5.3.1.2 Undifferentiated Organic Ranges Compared with Past Years

Each of the undifferentiated organics measured during this survey were within their reported range since 1991 (Figure 5-2). Acid volatile sulfides which were historically high in 2011, dropped to background levels in 2012 and remained low in 2013. Concentrations of oil and grease, TKN, TOC and acid volatile sulfides in 2013 were variable but within range of the past 20 years with no sustained increasing or decreasing trends evident.

5.3.1.3 Undifferentiated Organics Compared with Reference Surveys

The average concentrations of undifferentiated organics reported in this survey were compared to concentrations found during three southern California regional surveys conducted in 1998, 2003 and 2008 (Table 5-4 and 5-5). O&G, TKN and AVS were not measured during these surveys. Average TOC concentrations in the Goleta survey area were lower or similar to concentrations measured by the other surveys. ER-L and ER-M threshold limits are not available for these constituents.

5.3.2 Heavy Metals

Heavy metals in the marine environment are relatively ubiquitous and, with the exception of mercury, can normally be detected in sediments in low amounts. When anthropogenic sources increase sediment concentrations above levels that can be assimilated by benthic organisms, their assemblages can be impaired. For example:

- Aluminum is generally considered to be nontoxic to organisms in its elemental state and is one of the most common elements on earth.
- Antimony is used for alloys and other metallurgical purposes. The salts, primarily sulfides and oxides are employed in the rubber, textile, fireworks, paint, ceramic, and glass industries (SWRCB 1973). Acute and chronic toxicity of antimony to freshwater aquatic life occur at water concentrations as low as 9000 to 1600 ppm, and toxicity to algal species occurs at about 610 ppm. There is no saltwater criterion available for antimony (Long and Morgan 1990).
- Arsenic is carcinogenic and teratogenic (causing abnormal development) in mammals and is mainly used as a pesticide and wood preservative. Inorganic arsenic can affect marine plants at concentrations as low as 13 to 56 ppm and marine animals at about 2000 ppm (Long and Morgan 1990). The USEPA (1983) gives a terrestrial range of 1-50 ppm, with an average of 5 ppm.
- Cadmium is widely used in manufacturing for electroplating, paint pigment, batteries and plastics. Toxicity in water to freshwater animals ranges from 10 ppb to 1 ppm, as low as 2 ppm for freshwater plants, and 320 ppb to 15.5 ppm for marine animals (Long and Morgan 1990). The USEPA (1983) places the terrestrial range for cadmium at 0.01 to 0.7 ppm, with an average of 0.06 ppm.
- Chromium is widely used in electroplating, metal pickling, and many other industrial processes. Chromium typically occurs as either chromium (III) or chromium (VI), the



latter being considerably more toxic. Acute effects to marine organisms range from 2,000 to 105,000 ppm for chromium (VI) and 10,300 to 35,500 ppm for chromium (III). Chronic effects range from 445 to 2,000 ppb for chromium (VI) and 2,000 to 3,200 ppb for chromium (III) (Long and Morgan 1990). The terrestrial range is 1 to 1,000 ppm with an average of 100 ppm (USEPA, 1983).

- Copper is widely used in anti-fouling paints. Saltwater animals are acutely sensitive to copper in water at concentrations ranging from 5.8 to 600 ppm. Mysid shrimp indicate chronic sensitivity at 77 ppm (Long and Morgan 1990).
- Iron is generally not considered toxic to marine organisms. Iron, in some organic forms, is a stimulator for phytoplankton blooms. Recent experiments in deep-sea productivity have shown a considerable increase in phytoplankton in normally depauperate mid-ocean waters when iron is added (Soule et al. 1996).
- Older paints and leaded gasoline are a major source of lead. Lead may be washed into the Harbor or become waterborne from aerial particulates. Adverse effects to freshwater organisms range from 1.3 to 7.7 ppm, although marine animals may be more tolerant (Long and Morgan 1990).
- Mercury is a common trace metal once used in industry and as a biocide. Acute toxicity to marine organisms in water ranges from 3.5 to 1678 ppm. Organic mercury may be toxic in the range of 0.1 to 2.0 ppm (Long and Morgan 1990).
- Nickel is used extensively in steel alloys and plating. Nickel is chronically toxic to marine organisms in seawater at 141 ppm (Long and Morgan 1990).
- Selenium is used as a component of electrical apparatuses and metal alloys and as an insecticide. Although there is no data available for selenium toxicity to marine organisms, the present protection criteria range is from 54 to 410 ppb (USEPA 1986). The normal terrestrial range is from 0.1 to 2.0 ppm with a mean of 0.3 ppm. Selenium and lead levels found and reported in Least Tern eggs from Venice Beach and North Island Naval Station in San Diego County were considered to be harmful to development (Soule et al. 1996).
- Silver has many uses in commerce and industry including photographic film, electronics, jewelry, coins, and flatware and in medical applications. Silver is toxic to mollusks and is sequestered by them and other organisms. Silver increases in the Southern California Bight with increased depth; high organic content and percent silt (Mearns et. al., 1991). The range in the rural coastal shelf is from 0.10 to 18 ppm, in bays and harbors from 0.27 to 4.0 ppm, and near outfalls 0.08 to 18 ppm (Soule et al. 1996). The normal terrestrial level ranges from 0.01 to 5.0 ppm, with a mean of 0.05 ppm.
- Soule and Oguri (1987, 1988) found the effects of tributyl tin can be toxic in concentrations as low as 50 parts per trillion in water. The terrestrial range for tin is 2 to 200 ppm, with a mean of 10 ppm. The California Department of Fish and Game considers tributyl tin to be the most toxic substance ever released in the marine environment. Tributyl tin may not be as bio-available in sediments as it is in seawater, and therefore may not affect the benthic biota in the same fashion.
- Zinc is widespread in the environment and is also an essential trace element in human nutrition. It is widely used for marine corrosion protection, enters the waters as airborne particulates, and occurs in runoff and sewage effluent. Acute toxicity of zinc in water to marine fish begins at 192 ppm, and chronic toxicity to marine mysid shrimp can occur as low as 120 ppm (Long and Morgan 1990). The normal terrestrial range is from 10 to 300 ppm, with a mean of 50 ppm (Soule et al. 1996).



5.3.2.1 Heavy Metal Spatial Patterns

The concentrations for each of the heavy metals measured for this survey are listed in Table 5-1. Of the fourteen metals measured, all were above detection at each of the sites. Differences in the concentrations of each metal among sites were small. Each of the fourteen metals correlated expectedly (decreased) with distance from the outfall. Four metals (aluminum, antimony, arsenic, and silver) correlated with marginal significance ($0.05 < p < 0.10$). Each of the fourteen metals correlated expectedly (decreased) with distance from Goleta Point and none correlated significantly. Three of the fourteen metals (antimony, chromium and selenium) correlated significantly ($p < 0.05$) with sediment particle size, but these relationships were not expected (metal concentration increased as particle size increased).

5.3.2.2 Heavy Metal Ranges Compared with Past Years

Each of the heavy metals measured during this survey were within their reported range since 1991 and there were no clear increasing or decreasing concentration trends, especially in recent years (Figure 5-2).

5.3.2.3 Heavy Metals Compared with Reference Surveys

The average concentrations of 14 of the heavy metals measured in this survey were compared to concentrations found during three SCBRMP surveys in 1998, 2003 and 2008 (Tables 5-4). Of the metals where comparisons could be made, several slightly exceeded concentrations measure in other surveys (aluminum, chromium, copper, mercury, nickel) (Table 5-5).

5.3.2.4 Heavy Metals Compared with NOAA Effects Range Thresholds

Metals concentrations measured at each station in the Goleta survey area during 2013 were compared to the ER-L and ER-M threshold values (Table 5-4). All metal concentrations were below both the ER-L and ER-M threshold limits.

5.3.3 Complex Organics

5.3.3.1 Pesticides, PCB's and PAH's

Pesticides, PCBs and PAHs are contaminants that are widespread in the environment, are toxic to marine organisms when concentrations are increased and can cause reproductive failure in organisms at higher levels in the food chain. The sources and relative toxicity of each of these organic chemical groups are discussed below.

- DDT is a pesticide that has been banned since the early 1970's, but the presence of non-degraded DDT suggests that either subsurface DDT is being released during erosion and runoff in storms, or that fresh DDT is still in use and finding its way into coastal waters (Soule et al. 1996). DDT has been found to be chronically toxic to bivalves as low as 0.6 ppb in sediment. Toxicity of two of DDT's breakdown products, DDE and DDD, were both chronically toxic to bivalve larvae as low as about 1 ppb (Long and Morgan 1990).
- Of the non-DDT pesticides, concentrations of chlordane between 2.4 and 260 ppm in water are acutely toxic to marine organisms. Heptachlor is acutely toxic in water from 0.03 to 3.8 ppm. Heptachlor epoxide, a degradation product of heptachlor, is acutely toxic to marine shrimp at 0.04 ppm in water. Dieldrin is acutely toxic to estuarine organisms from 0.7 to 10 ppb. Endrin shows acute toxicity within a range of 0.037 to 1.2 ppb. Aldrin is acutely toxic to marine crustaceans and fish between 0.32 and 23 ppb. The EPA freshwater and saltwater criteria for aldrin are 3.0 and 1.3 ppb,



respectively (Long and Morgan 1990). No toxicity data were found for any of the other chlorinated compounds measured during this survey.

- Although PCBs are not pesticides, their similarity to other chlorinated hydrocarbons makes their inclusion in this section appropriate. Before being banned in 1970, the principal uses of PCBs were for dielectric fluids in capacitors, as plasticizers in waxes, in transformer fluids, and hydraulic fluids, in lubricants, and in heat transfer fluids (Laws 1981). Arochlor 1242, a PCB congener, was acutely toxic in water to marine shrimp in ranges of 15 to 57 ppm (Long and Morgan 1990).
- The major sources of polynuclear aromatic hydrocarbons (PAH's) are believed to be the combustion of fossil fuels and petroleum or oil shales. PAH impact is characterized by altered community structure, abundance, and diversity near the pollutant source (Daily, et.al. 1993).

5.3.3.2 Pesticide, PCB, and PAH Spatial Patterns

Pesticides, PCB and PAH concentrations at the six sampling stations are listed in Table 5-1 and complex organic derivatives are listed in appendix table 10-7. Similar to some previous surveys there were no chlorinated hydrocarbons (DDTs, BHCs, total chlordane, or PCBs) detected in Goleta sediments.

Similar to past years, total PAHs were above detection at each site in the survey area, with concentrations ranging from 24.9 at station B6 to 148.1 at station B4 near the outfall. The dominant PAH congener was perylene which made up the largest percentages of the total PAHs at each site. Total PAHs correlated unexpectedly and non-significantly with the distance to the outfall, and expectedly and non-significantly with distance from Goleta Point.

5.3.3.3 Pesticide, PCB and PAH Ranges Compared with Past Years

Total DDT pesticides, chlorinated hydrocarbons and PAH concentrations were within the range of previous years (Figure 5-2). Total PCBs were below detection in the Goleta survey area for the tenth year in a row since 2004.

5.3.3.4 Pesticides, PCB's and PAH's Compared with Reference Surveys

The average concentrations of chlorinated pesticides (DDTs), PCBs and PAHs measured during the 2013 survey were compared to concentrations found during three southern California reference site surveys conducted in 1998, 2003 and 2008 (Table 5-4). Concentrations of each group of organics were similar to or less than those measured on the inner shelf and near SPOTWs in during each of the SCBRMP reference surveys.

5.3.3.5 DDT Pesticides & PCB's Compared with NOAA Effects Range Thresholds

Pesticide, PCB and PAH concentrations measured in the Goleta survey area were compared to the NOAA ER-L and ER-M threshold values (Table 5-4). Each group of constituents was well below these thresholds, except DDT which slightly exceeded the ER-L.



5.4 Discussion

Results from this survey support past studies in that the Goleta outfall discharge has little or no impact upon the chemical composition of local sediments. In order to confirm this, results from the chemical analysis of the benthos were compared among stations, compared to past surveys in the area, compared to other studies performed in southern California, and compared to levels known to have caused toxicity or other environmental impacts to resident marine infauna.

To determine if contaminant trends were significant across stations, results for each variable were correlated against three independent variables: distance from outfall diffuser, distance from Goleta Point, and median particle size. Goleta Point is a documented area of particularly heavy crude oil seepage. Since the diffuser is located relatively close to the Point (approximately 1,500 meters east) it is prudent to attempt to partition out the potential influences of seepages from the impact of the discharge. Correlation against particle size is important because it is well known that metals and other contaminants often adhere more readily to finer particles, and differences among stations may be due to differences in amount of fine material (Gray 1981).

Metal concentrations in the Goleta survey area were not as heavily influenced by distance from Goleta Point and particle size during 2013 as in past years (Aquatic Bioassay 1997 to 2009). In fact, the concentrations of each of the metals were similar across sites. Of the fourteen metals measured, all correlated expectedly but non-significantly (decreased) with distance from the outfall. In addition, all of the metals correlated expectedly and non-significantly with distance to Goleta Point. Antimony, chromium and nickel correlated unexpectedly and significantly with sediment particle size.

In 2013, chlorinated pesticides (DDTs, chlordane, dieldrin, etc.) and PCBs were below detection. In past surveys, total PAHs were nearly always measured in greatest concentrations near Goleta Point and declined on a gradient toward the outfall. However, in 2013 (as in 2011 and 2012) this was not the case with PAH concentrations being similar across sites. The reason for the reduction in sediment PAH concentrations are unclear, but indicate that oil seepage from Goleta Point is highly variable.

This year's results were compared to past measures made in the Goleta survey area since 1991. Concentrations of sediment contaminants have remained relatively stable over time and in 2013 were within the ranges of past years. Acid volatile sulfides (AVS) which were greater on average in 2011 compared to any survey in the past 20 years, returned to normal background concentrations in 2012 and remained low in 2013. Organic contaminants remained either low or below detection in 2013. Total DDTs were again elevated in 2013 after being below detection since 2010. In addition, total PCBs have not been detected in the survey area since 2004.

This year's results were compared to sediment contaminant concentrations measured during the 1998, 2003 and 2008 SCBRMP surveys on the inner shelf (depth < 30m) and near SPOTWs (SCBRMP 1998, 2003 and 2008). Of the metals where comparisons could be made, several slightly exceeded concentrations measure in other surveys (aluminum, chromium, copper, mercury, nickel). Of the organics measured, none were greater than the SCBRMP surveys.

The Goleta data were also compared to NOAA's Effects Range Low (ER-L) and Effects Range Median (ER-M) criteria. Based upon historical research, sediments with levels of chemical contaminants exceeding ER-L values have a "potential" of affecting sensitive benthic infauna or the sensitive live stages of the more tolerant organisms. Sediments containing



contaminants that exceed ER-M values will “probably” have a negative impact upon several groups of infauna organisms. In 2013 each constituent was well below the ER-L thresholds and far below the ER-M thresholds. The only exception to this was total DDT which slightly exceeded the ER-L. This indicates that Goleta sediments were not likely to have had an adverse effect on the benthic infauna community.

In summary, of the 22 constituents measured in Goleta sediments during the 2013 survey, none correlated expectedly and significantly with distance from the outfall. Since the concentration of the pollutants emanating from the plant are very low or below detection, the detection of contaminants in the vicinity of the outfall is likely due to other anthropogenic inputs such as runoff from Goleta Slough, areal deposition or naturally occurring processes such as the release of oil from the seeps located offshore of Goleta Point. Comparison of Goleta sediments with historical reference data from the southern California Bight showed that most constituents were similar to or below baseline concentrations. Additionally, all sediment chemical concentrations were below those levels thought to cause toxicity to sensitive infauna organisms.

Figure 5-1. Benthic sediment sampling locations (Stations B1 – B6) in the Goleta survey area.




Table 5-1. Sediment contaminant concentrations (dry weight) in the Goleta survey area.

Constituent ¹	Sediment Stations						Mean	S.D.	Correlations		
	B1	B2	B3	B4	B5	B6			Outfall	Point	Prt.Sz.
Undifferentiated Organics											
Oil and Grease (detection = 100 µg/g) ³	814	704	436	461	230	184	472	250	0.03	-0.94	0.03
TKN (detection = 0.6 µg/g) ³	308	433	377	365	719	296	416	156	-0.70	-0.09	0.54
TOC (detection = 100 µg/g) ³	2500	3400	200	4000	5100	1300	2750	1799	-0.64	0.14	0.43
AVS (detection = 0.05 µg/g) ³	172.23	11.26	4.40	4.11	112.17	3.63	51.30	72.98	-0.12	-0.66	0.66
Heavy Metals											
Aluminum (detection = 1.0 µg/g)	8072	8773	7832	8505	10540	6755	8413	1254	-0.73	-0.43	0.80
Antimony (detection = 0.025 µg/g)	0.15	0.14	0.11	0.13	0.17	0.11	0.135	0.024	-0.34	-0.49	0.86
Arsenic (detection = 0.025 µg/g)	5.10	6.03	5.42	5.47	5.54	4.82	5.40	0.41	-0.77	-0.41	0.45
Cadmium (detection = 0.0025 µg/g)	0.41	0.56	0.47	0.35	0.39	0.36	0.42	0.08	-0.30	-0.31	0.31
Chromium (detection = 0.0025 µg/g)	28.88	29.14	25.87	27.33	33.55	23.19	27.99	3.49	-0.63	-0.58	0.86
Copper (detection = 0.0025 µg/g)	4.11	5.95	4.29	5.08	6.75	3.53	4.95	1.22	-0.72	-0.29	0.79
Iron (detection = 1.0 µg/g)	8408	10916	9142	9650	11880	8253	9708	1436	-0.69	-0.19	0.79
Lead (detection = 0.0025 µg/g)	3.21	4.01	3.51	3.82	4.52	3.20	3.71	0.51	-0.72	-0.15	0.72
Mercury (detection = 0.00001 µg/g) ³	0.023	0.030	0.094	0.026	0.027	0.021	0.037	0.028	-0.52	-0.31	0.09
Nickel (detection = 0.01 µg/g)	12.65	17.17	13.45	13.98	17.93	11.95	14.52	2.46	-0.65	-0.24	0.83
Selenium (detection = 0.025 µg/g)	0.23	0.37	0.20	0.25	0.32	0.26	0.27	0.06	-0.20	0.08	0.70
Silver (detection = 0.01 µg/g) ³	0.04	0.06	0.04	0.06	0.07	0.04	0.05	0.01	-0.74	0.22	0.46
Tin (detection = 0.025 µg/g) ³	1.33	0.70	0.57	0.70	0.71	0.47	0.75	0.30	-0.20	-0.60	0.60
Zinc (detection = 0.025 µg/g)	20.81	28.14	22.45	23.81	30.38	20.11	24.28	4.13	-0.67	-0.20	0.81
Complex Organics (ng/g dry weight)²											
Chlorinated Pesticides											
DDTs ³	5.1	2.2	1.2	1.7	3.2	1.1	2.42	1.52	-0.20	-0.60	0.60
HCHs	0.0	0.0	0.0	0.0	0.0	0.0	0.00	0.00	0.00	0.00	0.00
Chlordane	0.0	0.0	0.0	0.0	0.0	0.0	0.00	0.00	0.00	0.00	0.00
Aldrin (detection = 1.0 µg/Kg)	1.0	1.0	1.0	1.0	1.0	1.0	1.00	0.00	0.00	0.00	0.00
Dieldrin (detection = 1.0 µg/Kg)	1.0	1.0	1.0	1.0	1.0	1.0	1.00	0.00	0.00	0.00	0.00
Heptachlor (detection = 1.0 µg/Kg)	1.0	1.0	1.0	1.0	1.0	1.0	1.00	0.00	0.00	0.00	0.00
Heptachlor epoxide (detection = 1.0 µg/Kg)	1.0	1.0	1.0	1.0	1.0	1.0	1.00	0.00	0.00	0.00	0.00
Mirex (detection = 1.0 µg/Kg)	1.0	1.0	1.0	1.0	1.0	1.0	1.00	0.00	0.00	0.00	0.00
Hexachlorobenzene (detection = 1.0 µg/Kg)	1.0	1.0	1.0	1.0	1.0	1.0	1.00	0.00	0.00	0.00	0.00



Table 5-1.continued

Constituent ¹	Sediment Stations						Mean	S.D.	Correlations		
	B1	B2	B3	B4	B5	B6			Outfall	Point	Prt.Sz.
Polychlorinated Biphenyls											
PCBs	0.0	0.0	0.0	0.0	0.0	0.0	0.00	0.00	0.00	0.00	0.00
Aroclors	0.0	0.0	0.0	0.0	0.0	0.0	0.00	0.00	0.00	0.00	0.00
Polycyclic Aromatic Hydrocarbons											
PAHs ³	74.3	36.5	29.0	148.1	49.0	24.9	60.30	46.54	0.55	-0.37	-0.09
1-Methylnaphthalene (detection = 1.0 µg/Kg)	1.3	1.0	1.0	1.0	1.0	1.0	1.05	0.12	0.26	-0.62	0.02
1-Methylphenanthrene (detection = 1.0 µg/Kg) ³	1.0	1.2	1.3	1.8	1.9	1.0	1.37	0.39	-0.97	0.29	0.06
2-Methylnaphthalene (detection = 1.0 µg/Kg)	1.5	1.1	1.1	1.0	1.1	1.2	1.17	0.18	0.53	-0.36	0.06
2,3,5-Trimethylnaphthalene (detection = 1.0 µg/Kg)	1.0	1.0	1.0	1.0	1.0	1.0	1.00	0.00	0.00	0.00	0.00
2,6-Dimethylnaphthalene (detection = 1.0 µg/Kg)	1.2	1.3	1.1	1.0	1.2	1.0	1.13	0.12	-0.32	0.58	0.77
Acenaphthene (detection = 1.0 µg/Kg)	1.0	1.0	1.0	1.1	1.0	1.0	1.02	0.04	-0.36	-0.05	-0.50
Biphenyl (detection = 1.0 µg/Kg)	1.0	1.0	1.0	1.0	1.0	1.0	1.00	0.00	0.00	0.00	0.00
Benz[a]anthracene (detection = 1.0 µg/Kg) ³	5.3	2.8	2.7	11.7	5.3	1.6	4.90	3.65	-0.66	-0.26	-0.03
Benzo[b]fluoranthene (detection = 1.0 µg/Kg) ³	6.1	3.6	2.8	11.6	3.9	2.0	5.00	3.52	-0.55	-0.74	-0.09
Benzo[e]pyrene (detection = 1.0 µg/Kg) ³	6.1	3.2	2.7	8.6	4.9	1.9	4.57	2.50	-0.55	-0.37	-0.09
Benzo[g,h,i]perylene (detection = 1.0 µg/Kg) ³	1.0	1.0	1.0	16.7	1.0	1.0	3.62	6.41	-0.53	0.13	-0.65
Fluoranthene (detection = 1.0 µg/Kg) ³	15.9	7.1	5.4	22.3	8.1	4.9	10.62	6.97	-0.55	-0.37	-0.09
Naphthalene (detection = 1.0 µg/Kg)	2.4	2.1	1.9	2.3	2.5	2.5	2.28	0.24	0.46	0.28	0.29
Perylene (detection = 1.0 µg/Kg) ³	72.1	30.3	17.4	19.3	33.4	13.6	31.02	21.54	-0.20	-0.60	0.60

Bold = Marginally significant (0.05 < p < 0.10)

Bold = Significant (p < 0.05)

1. Minimum detection limits, reporting limits and methods are listed in Appendix 10.4
2. Complex organic derivatives are listed in Appendix 10.4.
3. Non-normal data. Correlations by nonparametric Spearman's rho.






Figure 5-2. Average concentrations (\pm SD) of sediment contaminants measured between 1991 and 2013 in the Goleta survey area. TOC, acid volatile sulfide, aluminum, iron, selenium and tin were not measured from 1991 to 1995.






Figure 5-2. (continued)



Table 5-4. Comparison of sediment contaminants found in the Goleta survey area to the Southern California Bight Regional Monitoring Program (SCBRMP) data from 1998, 2003 and 2008; and, the NOAA status and trends ERL and ERM threshold values. The SCBRMP survey includes comparisons against stations located near SPOTWs and shallow water reference sites. Bolded reference surveys or thresholds were exceeded by 2013 Goleta contaminant concentration ranges.

Constituent	GOLETA S.D.		SCBRMP (2008) ¹				SCBRMP (2003) ²				SCBRMP (1998) ³		NOAA (1990) ⁴ , Long, et.al. (1995) ⁵	
	Mean	Range	Inner Shelf Mean	95% CI	So Cal Bight Mean	95% CI	Inner Shelf Mean	95% CI	Small POTW Mean	95% CI	SPOTW Mean	Shallow Mean	ER-L	ER-M
Undifferentiated Organics														
Oil and Grease	472	184 - 814	---	---	---	---	---	---	---	---	---	---	---	---
TKN	416	296 - 719	---	---	---	---	---	---	---	---	---	---	---	---
TOC	2750	200 - 5100	6600	4100	30000	100	2700	800.00	5400	1600	5500	4200	---	---
AVS	51.3	3.6 - 172.2	---	---	---	---	---	---	---	---	---	---	---	---
Heavy Metals														
Aluminum	8413	6755 - 10540	5256	726	15372	1594	9212	2233	13244	3585	---	---	---	---
Antimony	0.14	0.11 - 0.17	0.12	0.02	0.28	0.04	0.14	0.04	0.15	0.02	1.09	1.59	2	25
Arsenic	5.40	4.82 - 6.03	4.3	1.2	6.70	1.20	4.2	1.4	4.6	0.67	7.67	4.39	8.2	70
Cadmium	0.42	0.35 - 0.56	0.23	0.03	0.88	0.12	0.20	0.06	0.22	0.05	0.28	0.36	1.2	9.6
Chromium	27.99	23.19 - 33.55	16	3.8	56.0	9.9	27	6.8	27	5.6	24.72	19.02	81	370
Copper	4.95	3.53 - 6.75	4.4	0.8	23.00	5.80	6.6	1.8	9.0	2.5	17.41	6.82	34	270
Iron	9708	8253 - 11880	10239	2233	26218	3125	12952	2784	16255	3655	---	---	---	---
Lead	3.71	3.20 - 4.52	5.0	1.3	12.00	1.40	4.7	1.1	4.90	0.81	15.92	10.14	46.7	218
Mercury	0.037	0.021 - 0.094	0.02	0.01	1.600	2.800	0.03	0.01	0.05	0.03	0.050	0.036	0.15	0.71
Nickel	14.52	11.95 - 17.93	9	1.7	27.00	2.80	13	3.8	11	2.0	13.85	15.50	20.9	51.6
Selenium	0.27	0.20 - 0.37	0.44	0.11	3.50	2.60	0.69	0.22	0.55	0.12	0.97	0.47	---	---
Silver	0.05	0.04 - 0.07	0.12	0.06	0.91	0.40	0.13	0.06	0.14	0.06	0.12	0.19	1.0	3.7
Tin	0.75	0.47 - 1.33	---	---	---	---	---	---	---	---	---	---	---	---
Zinc	24.28	20.11 - 30.38	25	6.8	71.00	5.90	34	7.8	40	8.0	52.14	33.59	150	410
Complex Organics														
DDTs	0.0024	0.0011 - 0.0051	0.0023	0.0004	0.1260	0.0970	0.0023	0.0004	0.0012	0.0002	0.020	0.036	0.00158	0.0461
HCHs	0.0000	0.0000 - 0.0000	---	---	---	---	---	---	---	---	---	---	---	---
Chlordane	0.0000	0.0000 - 0.0000	0.0005	0.0001	0.0016	0.0008	0.00001	0.00001	0.00000	0.00000	---	---	---	---
PCBs	0.0000	0.0000 - 0.0000	0.0002	0.0000	0.1700	0.0067	0.0024	0.00001	0.0001	0.00001	0.004	0.005	0.0227	0.18
PAHs	0.0603	0.0249 - 0.1481	0.0512	0.0449	0.2860	0.0390	0.0512	0.0449	0.0249	0.0087	0.118	0.073	4.022	44.792

1. SCCWRP, 2012; 2. SCCWRP, 2006; 3. SCCWRP 2003; 4. Long and Morgan, 1990; 5. Long et al., 1995.



Table 5-5. Summary of sediment contaminant spatial trends and concentrations found in the Goleta survey area to the Southern California Bight Regional Monitoring Program (SCBRMP) data from 1998, 2003 and 2008; and, the NOAA status and trends ERL and ERM threshold values.

Constituent	Expected Correlation w/ Dist from Outfall	Expected & Significant Correlation	Exceeds Reference Surveys?						Exceeds	
			2008 Inner Shelf	2008 So Cal Bight	2003 Inner Shelf	2003 SPOTW	1998 SPOTW	1998 Shallow	ER-L?	ER-M?
Oil and Grease	No	No	---	---	---	---	---	---	---	---
TKN	Yes	No	---	---	---	---	---	---	---	---
TOC	Yes	No	No	No	Yes	No	No	No	---	---
AVS	Yes	No	---	---	---	---	---	---	---	---
Aluminum	Yes	No	Yes	No	No	No	---	---	---	---
Antimony	Yes	No	Yes	No	No	No	No	No	No	No
Arsenic	Yes	No	Yes	No	Yes	Yes	No	Yes	No	No
Cadmium	Yes	No	Yes	No	Yes	Yes	Yes	Yes	No	No
Chromium	Yes	No	Yes	No	Yes	Yes	Yes	Yes	No	No
Copper	Yes	No	Yes	No	No	No	No	No	No	No
Iron	Yes	No	No	No	No	No	---	---	---	---
Lead	Yes	No	No	No	No	No	No	No	No	No
Mercury	Yes	No	Yes	No	Yes	No	No	Yes	No	No
Nickel	Yes	No	Yes	No	Yes	Yes	Yes	No	No	No
Selenium	Yes	No	No	No	No	No	No	No	No	No
Silver	Yes	No	No	No	No	No	No	No	No	No
Tin	Yes	No	---	---	---	---	---	---	---	---
Zinc	Yes	No	No	No	No	No	No	No	No	No
DDTs	Yes	No	Yes	No	Yes	Yes	Yes	No	Yes	No
HCHs	No	No	---	---	---	---	---	---	---	---
Chlordane	No	No	No	No	No	No	---	---	No	No
PCB'S	No	No	No	No	No	No	No	No	No	No
PAH'S	No	No	Yes	No	Yes	Yes	No	No	No	No



CHAPTER 6

Benthic Infauna

6.1. Background

The benthic infauna community is composed of those species living in or on the bottom (benthos). This community is very important to the quality of the habitat because it provides food for the entire food web including juvenile and adult fishes who are bottom feeders. Usually polychaete annelid worms, molluscs, and crustaceans dominate the benthic fauna in shallow, silty, sometimes unconsolidated, habitats. In areas where sediments are contaminated or frequently disturbed by natural events such as storms or by manmade events, nematode round worms, oligochaete worms, or tolerant polychaetes or mollusks may dominate the fauna temporarily. Storms can cause organisms to be washed away or buried under transported sediment, or can cause changes in the preferred grain size for particular species. Excessive runoff may lower normal salinities, and thermal regime changes offshore may disturb the composition of the community. Some species of benthic organisms with rapid reproductive cycles or great fecundity can out-compete other organisms in recolonization, at least temporarily after disturbances, but competitive succession may eventually result in replacement of the original colonizers with more dominant species.

6.2. Materials and Methods

Field sampling for all benthic sediment components is described in Chapter 4. Sediments to be analyzed for infauna content were sieved through 1.0 millimeter screens. The retained organisms and larger sediment fragments were then washed into four-liter plastic bottles, relaxed with a magnesium sulfate solution, and preserved with 10% buffered formalin. Five replicates were collected from six benthic infauna stations (B1, B2, B3, B4, B5, and B6; see Figure 3-1). Screened and preserved sediments collected in the field were delivered to the Ventura laboratory for counting, sorting, and identification. Infauna were sorted out by Aquatic Bioassay staff biologists and separated into five groups: echinoderms, mollusks, polychaetes, crustaceans, and miscellaneous. For each station, organisms were counted per group in accordance with *Techniques for Sampling and Analyzing the Marine Macrobenthos* EPA 600/3-78-300, March 1978; *Quality Assurance and Quality Control (QA/QC) for 301(h) Monitoring Programs: Guidance on Field and Laboratory Methods*, Tetra Tech 1986; and *Southern California Bight Pilot Project Field Operations Manual*, 2008. Each sorted sample was re-checked by a second biologist for representatives not found during the first inspection. Infauna was identified by SCAMIT taxonomists Tony Phillips for crustaceans and polychaetes and other phyla, Megan Lily of the City of San Diego for echinoderms and, and Kelvin Barwick with the Orange County Sanitation District for mollusks. A complete list of infauna is included in Appendix 10.6. Aquatic Bioassay maintains and updates standardized type collections and voucher specimens for most southern California infauna.

Following enumeration of infauna organisms by species, the total and phyla group numbers of individuals, and numbers of separate species were compiled for each station replicate. In addition, several required biological indices were calculated: Shannon Weiner species diversity (H'), Margelef's richness index (d), Simpson's species diversity (SI), Schwartz's dominance (D), the infauna trophic index (ITI) and Benthic Response Index (BRI). Analysis of Variance (ANOVA) was used to compare average metrics values among stations. Species compositions were compared using numerical classification and ordination. Brief descriptions of the indices are presented below.



Shannon Diversity. The Shannon Diversity Index (H') (Shannon and Weaver 1963) is defined as:

$$H' = - \sum_{j=1}^s \{(n_j/N) \ln (n_j/N)\},$$

where: n_j = number of individuals of the j th species,
 s = number of species in the sample,
 N = number of individuals in the sample.

Margalef's Richness. Margalef's Species Richness Index (d) (Margalef 1958) is:

$$d = s-1 / \ln N,$$

where: s = number of species in the sample,
 N = number of individuals in the sample.

Simpson's Diversity. The Simpson's Diversity Index (SI) (Simpson 1949) is:

$$SI = 1 - \sum_{i=1}^s (p_i)^2,$$

where: p_i = proportion of individuals of the i th species in the community.

Schwartz' Dominance. Schwartz's Dominance Index (D) is defined as the minimum number of species required accounting for 75% of the individuals in a sample (Schwartz 1978).

Infauna Trophic Index. This index measures the prevailing feeding modes of benthic infauna. Higher values denote southern California species assemblages dominated by suspension feeders, which are more characteristic of unpolluted environments. Lower index values denote assemblages dominated by deposit feeders more characteristic of areas near major outfalls (Word 1980):

$$ITI = -33.33 \{n_2 + (2)(n_3) + (3)(n_4) / n_1 + n_2 + n_3 + n_4\},$$

where: n_1, \dots, n_4 = numbers of individuals in species trophic groups 1, ..., 4, respectively.

Benthic Response Index. The BRI is the abundance-weighted average pollution tolerance of species occurring in a sample (Smith *et al.* 2001). The general index formula is:

$$BRI_s = \frac{\sum_{i=1}^n a_{si}^f p_i}{\sum_{i=1}^n a_{si}^f} \quad (1)$$

where BRI_s is the BRI value for sampling unit s , n is the number of species in s , p_i is the pollution tolerance of species i , a_{si} is the abundance of species i in s , and f is an exponent used to transform the abundance values. The primary objective of BRI development is to assign pollution tolerance scores p_i to species based on their position on a pollution gradient. Once



assigned, the scores can be used to assess the condition of the benthic community by calculating the BRI. A reference threshold, below which natural benthic assemblages normally occur, was identified at an index value of 31, the point on the pollution vector where pollution effects first resulted in a net loss of species. Three additional thresholds of response to disturbance were defined at index values of 42, 53 and 73, representing points at which 25%, 50%, and 80% of the species present at the reference threshold were lost.

Analysis of Variance (ANOVA). ANOVA's were used to compare population variables and sediment chemistry concentrations among stations. ANOVA analysis requires two steps. In the first step, differences in a variable among stations are evaluated to determine if they are sufficiently large to be statistically significant ($p \leq 0.05$). If they are, then a second test must be performed to determine which stations are significantly different from another station or stations. In this report, this second step is called the comparison of means. For example, a comparison of means stating: OS1 > OS2, OS3 > OS4, indicates that, for that particular variable, Station OS1 is significantly larger than Stations OS2, OS3, and OS4, and Stations OS2 and OS3 are also significantly larger than Station OS4. For chemical contaminants, if stations near the outfall are significantly higher than stations farther away, that compound should be evaluated further. For population variables, the opposite is true.

Cluster Analysis. Cluster analysis was used to define groups of samples, based on species presence and abundance, which belong to the same community without imposing an *a priori* community assignment. Identified clusters were then evaluated to define the habitat to which they belong. In cluster analysis, samples with the greatest similarity are grouped first. Additional samples with decreasing similarity are then progressively added to the groups. The percentage dissimilarity (Bray-Curtis) metric (Gauch, 1982; Jongman et al., 1995) was used to calculate the distances between all pairs of samples. The cluster dendrogram was formed using the unweighted pair-groups method using arithmetic averages (UPGMA) clustering algorithm (Sneath and Sokal, 1973). All steps were completed using the computer program MVSP (Multivariate Statistical Package, v3.12, 2000). Only the most commonly occurring species were used in the analysis, in this case only those that occurred at more than one station and season.

For normal (station by station) classifications, the Bray-Curtis Index is:

$$B.C. = \sum_{i=1}^s \min(P_{ij}, P_{ik}),$$

where: P_{ij} = proportion of species i collected at station j ,
 P_{ik} = proportion of species i collected at station k ,
 s = number of species.

For inverse (species group by species group) classifications:

$$B.C. = \sum_{i=1}^N \min(P_{ij}, P_{ik}),$$

where now: P_{ij} = proportion collected at station i of species j ,
 P_{ik} = proportion collected at station i of species k ,
 N = number of stations.



Ordination analysis. Ordination analysis displays the sampling stations as points in a multidimensional space. The distances between the stations (points) in the space are proportional to the dissimilarity of the communities found at the respective stations. The different dimensions of the ordination space, called axes, define independent gradients of biological change in the community data. The projections of the station points onto the various axes are called scores. The axes are ordered so that the first axis displays a maximal amount of community change; the second axis defines a maximal amount of the remaining community change, and so on for subsequent axes. Often most of the relevant community changes are displayed in a few ordination axes.

6.3. Results

6.3.1. Benthic Infauna

6.3.1.1. Infauna Abundance

The simplest measure of resident animal health is the abundance of infauna collected per sampling effort. Measures of abundance include biomass and numbers of individuals, which is partially dependent upon the volume of sediment collected in the grab. For this survey, abundance was determined to be all of the non-colonial animals collected from one replicate Van Veen Grab (0.1 square meter surface area) and retained on a 1.0 mm screen (note that abundance per square meter can be easily calculated by multiplying individuals per grab by ten). Five replicates were collected from six sediment stations.

Spatial infauna abundance patterns. Infauna abundances at the six sediment sampling stations are listed in Table 6-1. Numbers of individuals were significantly greatest at station B5 (average = 1,796) near the Goleta outfall compared to each of the other sites. Numbers of individuals correlated unexpectedly and significantly with distance from the outfall, unexpectedly and non-significantly with distance from Goleta Point, and unexpectedly with particle size.

Infauna abundance patterns compared with past years. Figure 6-1 illustrates biological metric trends over time in the Goleta survey area during the past twenty years. The average numbers of individuals increased between 1990 and 1994 and then steadily declined through 1999. Low values during 1998 and 1999 may reflect the El Nino conditions present then. In 2000, values began to increase through 2002 (average = 700), dipped in 2003, and then nearly doubled to historic highs during the period between 2004 and 2006 (average = 1566). Infauna abundances declined in 2007 and 2008 to levels similar to the years previous to 2004. From 2009 thru 2013, abundances have remained relatively stable (average ~ 1,000).

Infauna abundance values compared with other surveys. Table 6-2 compares abundance and other variables with reference control stations from the Southern California Bight Regional Monitoring Program (SCBRMP) surveys conducted in 1998, 2003 and 2008. Average numbers of individuals collected in the Goleta survey area were far greater than the averages measured at reference site locations in each of the SCBRMP surveys.

6.3.1.2. Infauna Species

Another simple measure of population health is the number of separate infauna species collected per sampling effort (i.e. one Van Veen Grab). Because of its simplicity, numbers of species is often underrated as an index. If the sampling effort and area sampled are the same



for each station, however, this index can be one of the most informative. In general, stations with higher numbers of species per grab tend to be in areas of healthier communities.

Spatial infauna species patterns. Infauna species at the six sediment sampling stations are listed in Table 6-1. Numbers of species were significantly greater at station B5 (average = 164) compared to all other stations (average range = 101 to 132) by ANOVA ($p < 0.05$). Lowest numbers of species were collected at reference station B6. Numbers of species correlated unexpectedly and significantly with distance from the outfall, unexpectedly and significantly with Goleta Point, and unexpectedly and non-significantly with particle size.

Infauna species patterns compared with past years. Figure 6-1 illustrates biological metric trends over time in the Goleta survey area during the past twenty years. Similar to numbers of individuals, numbers of species increased between 1991 and 1994 and then steadily declined through 1999 possibly owing to an El Nino effect. Since 2000 the average number of species has steadily increased through 2006 when it reached a historic high (average = 181). Since 2006 the average number of species declined slightly thru 2013 (average = 124).

Infauna species values compared with other surveys. Table 6-2 compares numbers of species and other variables with reference control stations from SCBRMP surveys conducted in 1998, 2003 and 2008. Ranges for Goleta species counts were greater than ranges measured in each of the SCCWRP reference site surveys.

6.3.1.3. Infauna Diversity

Species diversity indices are similar to numbers of species; however they often contain an evenness component, as well. For example, two samples may have the same numbers of species and the same numbers of individuals. However, one station may have most of its numbers concentrated into only a few species while a second station may have its numbers evenly distributed among its species. The diversity index would be higher for the latter station. The diversity indices required in the Goleta permit are the Shannon Diversity Index, Margalef Richness Index, and Simpson Diversity Index. Since all of these indices are calculated from the same measures (numbers of individuals and numbers of species), they often show the same patterns, and are, thus, probably somewhat redundant (Table 6-1). Infauna population metrics are presented by station. Comparisons are made using correlation analysis and ANOVA.

Spatial infauna diversity patterns. Infauna diversities at the six sediment-sampling stations are listed in Table 6-1. Diversity, as measured by Shannon's, Margalef's, and Simpson's indices were similar across sites and uniformly elevated in the survey area. Shannon and Simpson Diversity were not significantly different among stations by ANOVA ($p < 0.05$). Margalef's Richness was significantly greatest at station B5 (average = 21.8) compared to the reference site B6 (15.7).

None of the correlations with distance to the outfall, Goleta Point or particle size were significant for Shannon or Simpson's Diversity. Margalef's Richness correlated unexpectedly and significantly with distance to the outfall and Goleta Point.

Infauna diversity patterns compared with past years. Figure 6-1 illustrates biological metric trends over time in the Goleta survey area during the past twenty years. Shannon Diversity has been high in the Goleta survey area during the entire time period, with averages ranging between 3.5 to over 4.0. Diversity was just below 4.0 through the 1990's and then began a slight decrease to a low in 2005. In 2006 diversity began to increase thru 2007 and 2008, and



reached a historic high in 2009 and 2010, before decreasing in 2011 and 2012. In 2013 diversity increased slightly.

Infauna diversity values compared with other surveys. Table 6-2 compares the Shannon Diversity Index reference stations from the SCBRMP surveys conducted in 1998, 2003 and 2008. Shannon Diversity measured in the Goleta survey area was similar to or greater when compared to each of the SCBRMP reference site surveys. Neither Margalef's nor Simpson's indices were calculated during the two SCCWRP programs.

6.3.1.4. Infauna Dominance

The Schwartz Dominance Index is defined as the minimum number of species required to account for 75% of the individuals in a sample. The infauna environment tends to be healthier when the dominance index is high, and it tends to correlate with species diversity.

Spatial infauna dominance patterns. Dominance at the six sediment-sampling stations is listed in Table 6-1. Dominance was not significantly different among sites by ANOVA. Dominance correlated expectedly and non-significantly with distance from the outfall, unexpectedly and non-significantly with distance from Goleta Point, and unexpectedly and non-significantly with sediment particle size.

Infauna dominance patterns compared with past years. Figure 6-1 illustrates biological metric trends over time in the Goleta survey area during the past twenty years. Dominance has been high in the Goleta survey area during the entire time period, ranging between 23 and 40. Dominance ranged between 35 and 40 through the 1990's and then began a slight decrease to a low in 2005. In 2006 dominance began to increase to an historic high (average = 36) in 2010, before decreasing back to 2005 levels in 2013.

Infauna dominance values compared with other surveys. Table 6-2 compares the dominance at reference sites from the SCBRMP surveys conducted in 1998, 2003 and 2008. Dominance in the Goleta survey area in 2013 was similar to or slightly less compared to the SCBRMP reference site surveys.

6.3.1.5. Infauna Trophic Index

The Infauna Trophic Index (SCCWRP 1978, 1980) was developed to measure the feeding modes of benthic infauna. Higher values denote California species assemblages dominated by suspension feeders, which are more characteristic of unpolluted environments. Lower index values denote assemblages dominated by deposit feeders more characteristic of sediments high in organic pollutants (e.g. near major ocean outfalls). SCCWRP has also provided definitions for ranges of infauna index values. Values that are 60 or above indicate "normal" bottom conditions. Values between 30 and 60 indicate "change", and values below 30 indicate "degradation". The infauna trophic index is based on a 60-meter depth profile of open ocean coastline in southern California. Therefore, its results should be interpreted with some caution when applied to Goleta's shallower stations (24 m).

Spatial Infauna Trophic Index patterns. Infauna Trophic Index (ITI) scores at the six sediment-sampling stations is listed in Table 6-1. ITI scores were significantly greatest at outfall stations B4, B5 and reference station B6 (average = 74 each). ITI values correlated expectedly and non-significantly with distance from the outfall, expectedly and significantly with distance from Goleta Point, and unexpectedly and non-significantly with particle size. ITI



scores at all stations were well above levels defining benthic communities that are changed (60) and far above levels defining benthic communities that are degraded (30).

Infauna Trophic Index patterns compared with past years. Figure 6-1 illustrates biological metric trends over time in the Goleta survey area during the past twenty years. Average ITI values have remained stable across years and were similar in 2013 to past surveys.

Infauna Trophic Index values compared with other surveys. The ITI was not calculated for the SCBRMP (1998, 2003 and 2008). This index has been replaced as a measure of biological condition by the Benthic Response Index (BRI).

6.3.1.6 Benthic Response Index

The Benthic Response Index (BRI) measures the condition of a benthic assemblage, with defined thresholds for levels of environmental disturbance (Smith et al. 2001). The pollution tolerance of each species is assigned based upon its distribution of abundance along a pre-established environmental gradient. To give index values an ecological context and facilitate their interpretation, four thresholds of biological response to pollution were identified. The thresholds are based on changes in biodiversity along a pollution gradient. A reference threshold, below which natural benthic assemblages normally occur, was identified at an index value of 31, the point on the pollution vector where pollution effects first resulted in a net loss of species. Three additional thresholds of response to disturbance were defined at index values of 42, 53 and 73, representing points at which 25%, 50%, and 80%, respectively, of the species present at the reference threshold were lost.

Spatial BRI patterns. Average BRI scores were significantly greatest by ANOVA at station B5 (average = 32) compared to all other sites with the lowest BRI score at reference station B6 (average = 26) (Table 6-1). The BRI scores correlated expectedly (increased) and significantly with distance to the outfall, expectedly and significantly with distance to Goleta Point, and non-significantly with particle size. Scores were below 31 for each station, except station B5 which was only slightly above (32), indicating there was no net loss of reference species in the survey area. This indicates that the sites in the Goleta survey area are similar to other shallow reference site locations in the Southern California Bight.

This was the fifth year the BRI was calculated for Goleta and therefore was not compared against past survey years. The BRI was calculated using reference site data collected throughout southern California, therefore the BRI results for the 2013 survey are comparable to reference site conditions.

6.3.1.6. Cluster & Ordination Analysis

Patterns of species composition in the receiving environment's infauna community were evaluated by comparing normal (station x station) and inverse (species group x species group) classifications using the Bray-Curtis pair-wise similarity index. As Bray-Curtis Index values between station groups approach zero, the population of animals that make up the community at those sites becomes more the same. A station dendrogram was constructed from the resulting pattern matrix (Figure 6-2). Rare species were excluded from the analysis so that 229 species that occurred at > three sites over the eight year period were retained for analysis (95% of the total number of individuals collected).

Stations clustered into four groups that were very similar to one another; with the Bray-Curtis Index values for all station nodes being less than 50% (Figure 6-2). The greatest Bray-Curtis



distance between any two station nodes was approximately 40%, which indicates very small differences in species abundances and composition between sites. Station group 1 included the outfall stations B4 and B5, group 2 included Goleta Point station B1, group 3 had stations B2 and B3, and reference station B6 was in group 4.

Of the twenty most relatively abundant species collected in each cluster group, six were shared across cluster groups, underscoring the community similarities among stations (Table 6-3). The most common species in the survey area were those typically found in coastal nearshore waters. In 2013 the polychaete *Mediomastus sp.* was the most relatively abundant species and was represented in each station group. Other abundant species included a polychaete worm (*Levinseniagracilis*) and crustaceans (*Idarcturusallelomorphus*, *Caprellacalifornica* and *Euphilomedescarcharodonta*).

When the biological metrics for each station cluster group were averaged together they showed that the infauna population in outfall and Goleta Point station groups 1 and 2 had somewhat greater abundances, numbers of species and diversity. However, average BRI scores for outfall station group 1 (32) were very slightly above the threshold (31) where reference species are being lost. However, station groups 2, 3 and 4 were just below the threshold (range = 26 to 28).

6.4. Discussion

Results from this infauna survey support past studies that indicated that the ocean outfall discharge does not appear to be strongly impacting the resident benthic infauna community. This was confirmed by statistically comparing results among stations both near and far from the diffuser, comparing results with historical surveys, comparing results with other studies performed in Southern California, and comparing stations by cluster analyses.

Evaluation of the biological metrics for the 2013 survey showed that there were significant differences among sites for abundance, numbers of taxa, ITI and BRI. Each of the standard metrics (excluding ITI and BRI) were greatest at outfall station B5, were least at the reference station (B6) and increased again near Goleta Point. This is in contrast to the 2012 survey when the infauna populations were slightly depressed near the outfall. It appears that Goleta Point plays a role in the distribution of infauna in the Goleta survey area. This pattern of increased infauna abundances and taxa near Goleta Point may be due to the increased availability of organic material emanating from the oil seeps that are present there (Pearson and Rosenberg 1978). These results indicate the difficulty with interpreting the results of hypothesis testing on infauna abundance data. To try to elucidate these patterns and assess what, if any, impacts might be occurring to the infauna community, two indices were calculated and cluster analysis was employed.

The Infaunal Trophic Index (ITI) assesses the health of the benthic community using trophic level feeding strategies. In 2013 ITI scores at all stations were well above levels defining benthic communities that are changed (60) and far above levels defining benthic communities that are degraded (30). ITI scores in the survey area ranged from least (70) at station B2 to greatest at the outfall stations B4, B5 and reference station B6 (all 74). The ITI has been employed to assess the health of benthic communities since the early 1980's. However, its use to assess communities residing at depths less than 60 m has been criticized.

The Benthic Response Index (BRI) scores (Smith et al. 2001) across all stations, except outfall station B5, were below 31 indicating that there was no net loss of reference species in the survey area. The BRI score at station B5 was only one point above the threshold. There was an expected and significant correlation with distance to the outfall and among stations by



ANOVA, with outfall station B5 having significantly greater (poorer) BRI scores compared to all other sites. The BRI approach differs from other multimetric techniques in using multivariate ordination as the basis for assigning pollution tolerance scores. The primary objective of BRI development is to assign pollution tolerance scores to species based on their position on a pollution gradient. Once assigned, the scores can be used to assess the condition of the benthic community. The BRI was developed using hundreds of infauna samples collected from throughout the southern California bight, at sites that were both degraded and in reference condition.

Biological metrics calculated for the 2013 survey were compared to results of past surveys at the same sampling locations since 1990. Each of the metrics measured in 2013 were within the ranges of past surveys.

Cluster analysis showed that the dissimilarity among both station and species groups were very low across the survey area. The three station clusters identified were at most 40% different from one another based on infauna abundances and taxa composition. Of the top twenty most abundant species in the survey area, six were shared by the three cluster groups, underscoring the community similarities among stations.

To further investigate the potential influence of the Goleta outfall on the infauna community, cluster analysis and ordination were conducted on infauna data sets collected from 2004 to 2013 (Figure 6-3). Ordination analysis showed that the largest portion of the variation in the infauna community during the time period could be described by ordination axis 1 (26%) which was closely associated with survey year. Stations clustered together on axis 1 by year with 2004 and 2005 infauna communities (cluster group 1) furthest from stations collected during 2011, 2012 and 2013 (cluster groups 5 and 6). This indicates that larger oceanographic conditions are defining the abundances and composition of species in the survey area. There was no clear outfall related gradient on either axis 1 or axis 2 which described 11% of the variation in the community.

The biological metrics for each site and survey were averaged by historic cluster group and showed there was very little difference across cluster groups indicating a relatively stable infauna population through time (Table 6-5). Of note was a reduction in average BRI scores from station group 1 (average = 31) in 2004 through 2006 to group 6 (average = 28) from 2011 to 2013. This indicates a gradual improvement in the biological condition of the survey area during the time period.

Finally, Goleta results were compared to measurements made of the inner continental shelf throughout southern California. All infauna population variables were comparable to or greater than those measured in regional surveys conducted by the SCBRMP in 1998, 2003 and 2008.

Although there are no specific numerical limitations regarding infauna animals, the California Ocean Plan (SWRCB 2007) states that:

The rate of deposition of inert solids and the characteristics of inert solids in the ocean shall not be changed such that benthic communities are degraded.

The dissolved sulfide concentration of waters in and near sediments shall not be significantly increased above that present under natural conditions.

The concentration of substances set forth in Chapter IV, Table B, in marine sediments shall not be increased to levels which would degrade indigenous biota.



The concentration of organic materials in marine sediments shall not be increased to levels which would degrade marine life.

Nutrient materials shall not cause objectionable aquatic growths or degrade indigenous biota.

Marine communities, including vertebrate, invertebrate, and plant species, shall not be degraded.

Waste management systems that discharge to the ocean must be designed and operated in a manner that will maintain the indigenous marine life and a healthy and diverse marine community.

Waste discharged to the ocean must be essentially free of: "2) Settleable material or substances that may form sediments which will degrade benthic communities or other aquatic life."

Waste discharged to the ocean must be essentially free of: "3) Substances which will accumulate to toxic levels in marine waters, sediments or biota."

Based upon spatial and temporal comparisons and analogies with other studies, the results of the infauna survey indicate that the discharge is in compliance with the general limitations and that it causes no adverse impact.



Table 6-1. Infauna population indices by replicate for each of the six Goleta survey area stations. Comparisons are made using correlation analysis and ANOVA ($p < 0.05$).

Constituent	Offshore Stations					
	B1	B2	B3	B4	B5	B6
INDIVIDUALS¹						
Repl. 1	855	971	839	658	2340	798
Repl. 2	1002	742	1096	1729	1647	621
Repl. 3	784	826	562	1057	2480	594
Repl. 4	677	1351	782	887	1202	405
Repl. 5	1464	790	940	906	1313	442
Mean =	956	936	844	1047	1796	572
Std. Dev. =	307	247	198	407	586	157
Lower Conf. Int. =	687	719	671	691	1283	434
Upper Conf. Int. =	1226	1153	1017	1404	2310	710
Overall Mean = 1025.3	r (outfall) = -0.49		r (point) = -0.28		r (prt.sz.) = -0.25	
Overall S.D. = 496.3	F = 6.98		Comp. of means = B5 > B1, B2, B3, B4, B5, B6			
SPECIES¹						
Repl. 1	122	120	136	103	163	128
Repl. 2	147	113	135	148	153	99
Repl. 3	141	120	97	145	191	108
Repl. 4	103	158	119	133	156	82
Repl. 5	147	120	143	132	155	86
Mean =	132	126	126	132	164	101
Std. Dev. =	19	18	18	18	16	18
Lower Conf. Int. =	115	110	110	117	150	84
Upper Conf. Int. =	149	142	142	148	177	117
Overall Mean = 130.1	r (outfall) = -0.56		r (point) = -0.42		r (prt.sz.) = -0.17	
Overall S.D. = 24.9	F = 6.28		Comp. of means = B5 > B4, B1, B2, B3, B6			
SHANNON DIVERSITY						
Repl. 1	3.75	3.67	3.9	3.39	3.42	3.89
Repl. 2	4.01	3.47	3.81	3.8	3.57	3.66
Repl. 3	4.03	3.61	3.63	3.64	3.58	3.69
Repl. 4	3.75	3.72	3.77	3.82	3.77	3.53
Repl. 5	3.47	3.79	3.77	3.63	3.53	3.47
Mean =	3.80	3.65	3.78	3.66	3.57	3.65
Std. Dev. =	0.23	0.12	0.10	0.17	0.13	0.16
Lower Conf. Int. =	3.60	3.55	3.69	3.50	3.46	3.51
Upper Conf. Int. =	4.00	3.76	3.86	3.81	3.68	3.79
Overall Mean = 3.68	r (outfall) = 0.07		r (point) = -0.24		r (prt.sz.) = 0.30	
Overall S.D. = 0.16	F = 1.51		Comp. of means = NA			
MARGALEF RICHNESS						
Repl. 1	17.92	17.30	20.05	15.72	20.88	19.01
Repl. 2	21.13	16.95	19.14	19.72	20.52	15.24
Repl. 3	21.01	17.72	15.16	20.68	24.31	16.75
Repl. 4	15.65	21.78	17.71	19.45	21.86	13.49
Repl. 5	20.03	17.84	20.74	19.24	21.45	13.95
Mean =	19.15	18.32	18.56	18.96	21.80	15.69
Std. Dev. =	2.34	1.97	2.21	1.89	1.49	2.25
Lower Conf. Int. =	17.10	16.59	16.62	17.30	20.50	13.72
Upper Conf. Int. =	21.20	20.04	20.50	20.62	23.11	17.66
Overall Mean = 18.75	r (outfall) = -0.53		r (point) = -0.44		r (prt.sz.) = -0.12	
Overall S.D. = 2.60	F = 4.57		Comp. of means = B5 > B6			

Bold = Marginally Significant ($0.05 < p < 0.10$)

Bold & Gray = Significant ($p < 0.05$)

1. The van Veen Grab collects samples one tenth of one square meter in area. To determine individuals per meter, multiply by ten.
2. Non-normal data: correlation coefficients and ANOVA's from non-parametric tests (Spearman's rho and Kruskal-Wallis H, respectively).



Table 6-1. continued

Constituent	Offshore Stations					
	B1	B2	B3	B4	B5	B6
SIMPSON DIVERSITY						
Repl. 1	0.96	0.94	0.97	0.91	0.94	0.96
Repl. 2	0.97	0.93	0.96	0.97	0.96	0.95
Repl. 3	0.97	0.93	0.95	0.96	0.94	0.96
Repl. 4	0.96	0.96	0.96	0.97	0.97	0.94
Repl. 5	0.94	0.95	0.96	0.97	0.96	0.93
Mean =	0.96	0.94	0.96	0.95	0.95	0.95
Std. Dev. =	0.01	0.01	0.01	0.03	0.01	0.01
Lower Conf. Int. =	19.20	22.03	20.29	19.88	24.65	21.34
Upper Conf. Int. =	26.20	24.95	25.23	21.70	27.68	22.39
Overall Mean = 0.952	r (outfall) = -0.10		r (point) = -0.20		r (prt.sz.) = 0.27	
Overall S.D. = 0.015	F = 7.69		Comp. of means = NA			
SCHWARTZ DOMINANCE						
Repl. 1	26	26	32	20	22	33
Repl. 2	39	23	27	28	30	25
Repl. 3	40	30	22	28	24	27
Repl. 4	25	27	28	32	34	23
Repl. 5	24	30	32	25	28	26
Mean =	31	27	28	27	28	27
Std. Dev. =	8	3	4	4	5	4
Lower Conf. Int. =	19	22	20	20	25	21
Upper Conf. Int. =	26	25	25	22	28	22
Overall Mean = 27.87	r (outfall) = 0.03		r (point) = -0.21		r (prt.sz.) = 0.16	
Overall S.D. = 4.72	F = 0.49		Comp. of means = NA			
INFAUNAL INDEX						
Repl. 1	70	72	73	74	75	74
Repl. 2	71	70	73	73	71	75
Repl. 3	74	71	72	75	68	74
Repl. 4	71	66	73	76	78	78
Repl. 5	70	71	71	73	78	73
Mean =	71	70	72	74	74	74
Std. Dev. =	1.5	2.3	1.2	1.4	4.5	2.1
Lower Conf. Int. =	19	22	20	20	25	21
Upper Conf. Int. =	26	25	25	22	28	22
Overall Mean = 72.60	r (outfall) = 0.10		r (point) = 0.38		r (prt.sz.) = 0.09	
Overall S.D. = 2.85	F = 3.26		Comp. of means = B5, B4, B6 > B2			
BENTHIC RESPONSE INDEX						
Repl. 1	29	30	25	28	33	25
Repl. 2	30	30	29	29	34	27
Repl. 3	27	28	27	30	33	26
Repl. 4	32	31	30	30	31	25
Repl. 5	30	27	26	29	31	25
Mean =	30	29	28	29	32	26
Std. Dev. =	1.7	1.6	2.2	0.9	1.2	1.2
Lower Conf. Int. =	19	22	20	20	25	21
Upper Conf. Int. =	26	25	25	22	28	22
Overall Mean = 28.93	r (outfall) = -0.56		r (point) = -0.51		r (prt.sz.) = -0.15	
Overall S.D. = 2.50	F = 11.50		Comp. of means = B5 > B1, B2, B3, B4, B6			

Bold = Marginally Significant (0.05 < p < 0.10)

Bold & Gray = Significant (p < 0.05)

1. The van Veen Grab collects samples one tenth of one square meter in area. To determine individuals per meter, multiply by ten.
2. Non-normal data: correlation coefficients and ANOVA's from non-parametric tests (Spearman's rho and Kruskal-Wallis H, respectively).



Figure 6-1. Infauna community variables, station (n = 6) means and standard deviations since 1990.




Figure 6-1. (continued).




Table 6-2. Comparison of Goleta infauna variables with results from other studies (per 0.1 m²).

Variable	Goleta 2012		SCBRMP 1998		SCBRMP 2003 Inner Shelf		SCBRMP 2008 Inner Shelf	
	Mean	Range	Mean	Range	Mean	±95% CI	Mean	SE
Number of Individuals	1025	405 - 2480	385	35 - 1696	283	30	346	22
Number of Species	130	82 - 191	85	18 - 162	62	5	85	4
Shannon Diversity Index	3.7	3.4 - 4.0	3.60	2.00 - 4.40	3.48	0.09	3.63	0.06
Dominance	27.9	20.0 - 40.0	--	-- - --	23	2	27	1




Figure 6-2. Station dendrogram based on cluster analysis (UPGMA, Sneath and Sokal 1973). The Bray-Curtis dissimilarity index was used to calculate the distances among stations and species (Gauch 1982, Jongman et. al. 1995).



Table 6-3. Average abundances of the top twenty species for each cluster group in 2013.

Species	Cluster Group			
	1	2	3	4
Mediomastus sp	239	106	109	33
Idarcturus allelomorphus	122	58		
Levinsenia gracilis	119	66	37	
Caprella californica	103			
Cossura sp A	55	33	73	28
Photis brevipes	43	30	26	
Oligochaeta	39	29	23	
Spiochaetopterus costarum Cmplx	37	50	48	20
Podocerus cristatus	35			
Idarcturus sp	34			
Spiophanes duplex	32	34	65	9
Aoroides intermedia	31	42		
Columbaora cyclocoxa	30	32		
Tellina modesta	27	52	85	24
Owenia collaris	27			
Gammaropsis thompsoni	26	74		
Pista agassizi	25			
Leptochelia dubia Cmplx	24	34	52	13
Euclymeninae sp A	24			
Aoroides sp	23			
Euphilomedes carcharodonta		187	267	87
Prionospio (Prionospio) jubata		52	39	13
Platynereis bicanaliculata		46		
Foxiphalus obtusidens		35		8
Alia tuberosa		34		
Foxiphalus golfensis		30	20	
Magelona berkeleyi		29	33	9
Macoma yoldiformis			74	12
Glottidia albida			49	19
Rhepoxynius stenodes			29	11
Photis californica			28	10
Pectinaria californiensis			23	
Ampelisca brevisimulata			19	12
Photis sp			18	7
Ampelisciphotis podophthalma				13
Amphideutopus oculatus				12
Caecognathia crenulatifrons				7
Westwoodilla tone				12



Table 6-4. Biological metrics for each station averaged by cluster group.

Station	Cluster Group	Number of Species	Total Abundance	BRI	ITI	Evenness	Margalef Richness	Schwartz Dominance	Shannon Diversity	Simpson Diversity
B5	1	300	1796	32	73	0.68	39.90	33	3.85	0.96
B1	2	277	956	29	71	0.77	40.22	43	4.3	0.97
B4	2	266	1047	29	74	0.75	38.11	36	4.2	0.97
	Average	272	1002	29	73	0.76	39.16	40	4.25	0.97
B2	3	250	936	28	69	0.71	36.40	32	3.93	0.95
B3	3	244	844	28	72	0.74	36.07	35	4.09	0.96
	Average	247	890	28	71	0.73	36.23	34	4.01	0.96
B6	4	203	572	26	74	0.75	31.82	32	3.99	0.96




Figure 6-3. Plot of ordination scores for infauna communities at stations measured from 2004 to 2013.



Table 6-5. Biological metrics for each station for each year individually from 2004 thru 2013 and averaged by cluster group.

Station/Year	Cluster Group	Number of Species	Total Abundance	BRI	ITI	Shannon Diversity
B1 05	1	320	1246	29.94	78.3	4.25
B1 06	1	308	1386	30.05	74.3	4.11
B2 05	1	249	1302	32.94	71.6	3.41
B2 06	1	302	1580	31.59	72.7	3.65
B3 05	1	289	1499	30.5	74.3	3.77
B3 06	1	302	1775	31.05	73.6	3.77
B4 05	1	265	1112	31.51	71	3.4
B4 06	1	293	1580	32.24	71.5	3.58
B5 05	1	306	1221	29.73	73.5	3.75
B5 06	1	318	1804	29.95	72.1	3.83
B6 05	1	271	1124	30.7	74.9	3.67
B6 06	1	293	1270	30.47	73.9	3.7
Average		293	1408	30.9	73.5	3.74
B1 04	2	369	2159	32.86	72.8	3.62
B2 04	2	331	1616	30.61	76.6	3.95
B3 04	2	249	1430	30.71	74	3.49
B4 04	2	242	1132	31.31	71.5	3.2
B5 04	2	262	1220	27.89	74.9	3.77
B6 04	2	260	945	26.56	78.9	3.88
Average		286	1417	30	74.8	3.65
B1 07	3	318	1022	31.05	75.9	4.38
B1 08	3	254	582	26.13	80.6	4.56
B2 07	3	251	729	31.66	79.8	4.37
B2 08	3	226	677	30.64	80.5	4.39
B3 07	3	264	1400	32.83	73.4	3.85
B3 08	3	262	1093	30.21	77.2	4.08
B4 07	3	249	1023	31.98	71.2	3.89
B4 08	3	238	854	32.26	63.4	3.97
B5 07	3	281	1220	31.18	75.2	4.08
B5 08	3	247	741	29.43	76.6	4.08
B6 07	3	321	1349	30.53	78.3	4.22
B6 08	3	259	910	27.79	76.1	4.02
Average		264	967	30.5	75.7	4.16
B1 09	4	315	1203	28.13	77.9	4.29
B1 10	4	300	1210	28.09	74.5	4.29
B2 09	4	289	1024	29.82	76.4	4.39
B2 10	4	295	920	27.35	75	4.55
B3 09	4	278	1124	29.24	79.9	4.15
B3 10	4	276	985	28.33	75.3	4.41
B4 09	4	251	950	27.17	80.5	4.07
B4 10	4	272	995	30.39	71.5	4.21
B5 09	4	296	1154	27.16	81.4	4.17
B5 10	4	355	1973	28.6	79.6	4.18
B6 09	4	269	1037	26.75	80.8	4.5
B6 10	4	268	855	26.27	79.3	4.37
Average		289	1119	28.1	77.7	4.3
B4 11	5	243	735	29.42	72.7	4.17
B4 13	5	268	1047	28.92	74.1	4.29
B5 11	5	247	738	31.75	75.1	4.28
B5 13	5	314	1796	32.15	73.1	4.22
Average		268	1079	30.6	73.8	4.24
B1 11	6	328	1343	28.7	72.8	4.36
B1 12	6	335	1457	28.69	72.8	4.42
B1 13	6	280	956	28.84	71.1	4.43
B2 11	6	243	973	28.02	76.7	3.97
B2 12	6	263	1179	28.82	73.7	4.07
B2 13	6	254	936	28.38	69.2	4.07
B3 11	6	228	816	26.5	82.1	3.81
B3 12	6	259	1114	27.96	75.3	3.98
B3 13	6	248	844	27.7	72.3	4.19
B4 12	6	215	773	25.48	80.9	3.71
B5 12	6	222	830	24.38	78.9	3.65
B6 11	6	245	617	26.24	77.3	4.32
B6 12	6	228	842	28.77	73.5	3.68
B6 13	6	205	572	25.83	74.2	4.09
Average		254	947	27.5	75.1	4.05



CHAPTER 7

Trawled Fish and Invertebrate Populations

7.1. Background

Demersal fishes and megabenthic invertebrates (species living closely associated with the seafloor) are widely distributed on the soft-bottom habitats along the southern California shelf. This diverse community is composed of approximately 100 species of fish and several hundred species of invertebrates (Allen 1982, Allen et al. 1998, Moore and Mearns 1978). Since these populations are generally sedentary, they can act as indicators of human impacts on the soft bottom habitat. As a result, trawl programs have been part of the monitoring activities of both large and small municipal dischargers for nearly thirty years. The goal of the Goleta Sanitary District's trawl program is to look for population changes in the vicinity of the ocean outfall.

7.2. Materials and Methods

Trawl sampling was conducted in accordance with *Use of Small Otter Trawls in Coastal Biological Surveys*, EPA 600/3-78/083, August 1978; *Quality Assurance and Quality Control (QA/QC) for 301(h) Monitoring Programs: Guidance on Field and Laboratory Methods*, Tetra Tech 1986; and the *Southern California Bight Project Field Operations Manual*, 2008. Duplicate ten-minute trawls were taken at a uniform speed of 2.0 - 2.5 knots with a 7.6 m Marinovich otter trawl. Care was taken to not trawl over previous transects or grab sampling sites. For each trawl, all fish and macroinvertebrates were identified, counted, measured, and weighed. Collection observations, such as algae or cobble in the trawl, were recorded. Fish abnormalities, such as fin rot, parasites, or tumors, were also noted. Species abundance lists were compiled for all trawl samples. All fish and invertebrates were identified by Jim Mann and Karin Patrick. All animals collected for tissue dissection were placed in plastic zip-lock bags in coolers over ice during transit.

Following enumeration of trawl organisms by species, the total and animal group biomasses, numbers of individuals, and numbers of separate species were compiled for each station replicate. In addition, several required biological indices were calculated: Shannon-Weiner species diversity (H'), Margalef's richness index (d), Simpson's species diversity (SI), and Schwartz's dominance (D). These indices are described in detail in Chapter 6, in Section 6.2, Materials and Methods. Since there were only two stations sampled, no clustering or numerical classification analyses could be calculated. Stations were compared by t-test (see Materials and Methods section above).

7.3. Results

The demersal fish and macrobenthic invertebrate community was compared among two trawl stations using measures of population abundance and diversity. These included numbers of individuals, numbers of species, species diversity, and species dominance. In addition, ranges of these variables were compared to surveys conducted in past years. Duplicate trawls were taken at two locations, one near Station B3 (TB3) and the other near Station B6 (TB6) (Figure 6-1).

7.3.1. Trawled Fish

7.3.1.1. Fish Community Metrics



The averaged fish community metrics and biomass for replicate trawls are presented in Table 7-1, with results by replicate presented in Appendix 10.7 (Tables 10-9 and 10-10). A total of 500 individual fish were collected from both stations combined during the 2013 survey, with the average numbers of individuals at TB3 (69) less than half the average numbers collected at TB6 (181) (Table 7-1). There was, however, no statistically significant difference in average abundances between sites ($p > 0.05$; Table 7-1). Similarly, the average numbers of species collected at Station TB3 (9) was nearly half that collected at TB6 (17) with a marginally significant difference between sites ($0.05 < p < 0.10$). Average biomass was nearly the same (3.09 and 3.28 Kg at station TB3 and TB6, respectively) and there was no significant difference between sites. Shannon Diversity, Simpsons Diversity and Dominance were low at each site and were not significantly different between sites. Margalef's Richness was significantly greater at TB6 (2.98) compared to TB3 (1.77).

7.3.1.2. Species Composition

As with past years, the fish caught in the 2013 trawls were typical of those found on most southern California near shore soft bottom habitats (Table 7-2). A total of 13 and 20 unique taxa, were collected at stations TB3 and TB6, respectively. The most abundant species collected in the Goleta survey area was the speckled sanddab (*Citharichthys stigmaeus*) followed by the Pacific sanddab (*Citharichthys sordidus*). At TB3 vermillion rockfish (*Sebastes miniatus*) were also abundant, while at TB6 darkblotched rockfish (*Sebastes crameri*) and California lizardfish (*Synodus lucioceps*) were abundant.

7.3.1.3. Fish Community Metrics Compared to Past Surveys

Fish assemblage community metrics for 2013 were compared to previous Goleta area surveys starting in 1991 (Figure 7-1). The numbers of individuals collected in 2013 was within the range of past surveys. Fish biomass was again very low during 2013 and similar to the past 20 years. Numbers of species was slightly greater in 2013 compared to 2012, and was far greater than 2008, when taxa richness reached an all-time low. In 2013, Shannon Diversity and dominance were low and similar to past surveys.

7.3.1.4. Fish Community Metrics Compared to Reference Surveys

Fish community metrics for the 2013 Goleta survey were compared to fish assemblage data collected in the northern region on the inner continental shelf in the southern California bight during the 2008 Southern California Bight Regional Monitoring Survey (SCBRMP) (SCCWRP 2011; Table 7-3). Number of individuals, number of species, Shannon Diversity and biomass were all well within the range fish assemblages found in the vicinity of the northern region inner shelf.

7.3.1.5. Fish Length

Fish size class distributions. The size frequency distributions for all fish collected from trawl samples are presented in Appendix 10.7 (Table 10.7-1). The size frequency distributions for one of the historically most abundant species in the survey area (speckled sanddabs, *Citharichthys stigmaeus*) are presented in Figure 7-2. Across years, sanddab lengths ranged from 3 to 13 cm at both stations, with 2013 having slightly more individuals in the 6 and 7 size class. At TB3, near the outfall, the numbers of fish collected were relatively evenly spread across size classes for all years, except in 2007 and 2012 when large numbers of individuals in the 7 and 8 cm size classes were captured. The majority of sanddabs collected 2004, 2007, 2009 and 2012 at TB6 were in 6 to 8 cm size classes.



Table 7-1. Trawled fish - Summary of biological metrics of fish collected at Stations TB3 and TB6. Comparison between sites by two sample T-test ($p < 0.05$).

Fish							
Metric	Station	TB3		TB6		T-test	
		Avg	SD	Avg	SD	t score	p =
Individuals		69	57	181	21	-2.62	0.12
Species		9	2	17	2	-3.77	0.06
Biomass (kg)		3.09	3.72	3.28	0.22	-0.07	0.95
Shannon Diversity		1.50	0.16	1.78	0.19	-1.58	0.26
Simpson Diversity		0.71	0.07	0.74	0.09	-0.43	0.71
Margalef Richness		1.77	0.10	2.98	0.34	-4.52	0.05
Schwartz Dominance		3	1	4	1	-1.41	0.29

Bold - Marginally Significant ($0.05 < p < 0.10$)

Bold - Significant ($p < 0.05$)

1. Non-normal data: T-test by Mann-Whitney U test.

Table 7-2. Trawled fish abundance and biomass sorted from most to least abundant.

Trawl TB3				Trawl TB6			
Scientific Name	Common Name	Mean Abundance	Mean Weight (kg)	Scientific Name	Common Name	Mean Abundance	Mean Weight (kg)
<i>Citharichthys stigmaeus</i>	speckled sanddab	28	0.14	<i>Citharichthys stigmaeus</i>	speckled sanddab	76	0.46
<i>Citharichthys sordidus</i>	Pacific sanddab	23	0.57	<i>Citharichthys sordidus</i>	Pacific sanddab	31	0.63
<i>Sebastes miniatus</i>	vermillion rockfish	8	0.09	<i>Sebastes crameri</i>	darkblotched rockfish	24	0.17
<i>Pleuronichthys decurrens</i>	curfin sole	3	0.19	<i>Synodus lucioceps</i>	California lizardfish	24	0.44
<i>Sebastes crameri</i>	darkblotched rockfish	2	<0.1	<i>Sebastes miniatus</i>	vermillion rockfish	5	0.07
<i>Sebastes dallii</i>	calico rockfish	2	<0.1	<i>Odontopyxis trispinosa</i>	pygmy poacher	4	<0.1
<i>Citharichthys xanthostigma</i>	longfin sanddab	1	<0.1	<i>Pleuronichthys verticalis</i>	homyhead turbot	4	0.45
<i>Hypsurus caryi</i>	rainbow seaperch	1	<0.1	<i>Syngnathus californiensis</i>	kelp pipefish	4	<0.1
<i>Neoclinus blanchardi</i>	sarcastic fringehead	1	0.05	<i>Icelinus quadriseriatus</i>	yellowchin sculpin	2	<0.1
<i>Odontopyxis trispinosa</i>	pygmy poacher	1	<0.1	<i>Pleuronichthys decurrens</i>	curfin sole	2	0.06
<i>Ophiodon elongatus</i>	lingcod	1	0.12	<i>Zaniolepis latipinnis</i>	longspine combfish	2	0.08
<i>Paralichthys californicus</i>	California halibut	1	1.75	<i>Ophiodon elongatus</i>	lingcod	1	<0.1
<i>Synodus lucioceps</i>	California lizardfish	1	<0.1	<i>Paralabrax nebulifer</i>	barred sand bass	1	<0.1
	composite weight*		0.19	<i>Parophrys vetulus</i>	English sole	1	<0.1
				<i>Porichthys notatus</i>	plainfin midshipman	1	<0.1
				<i>Raja inornata</i>	California skate	1	0.16
				<i>Scorpaenichthys marmoratus</i>	cabezon	1	<0.1
				<i>Sebastes saxicola</i>	stripetail rockfish	1	<0.1
				<i>Ulvicola sanctaerosae</i>	kelp gunnel	1	<0.1
				<i>Xystreurys liolepis</i>	fantail sole	1	0.30
					composite weight*		0.48

*Species <0.1 kg are weighed together as a composite weight.






Figure 7-1. Fish community metric annual averages (\pm SD) for Goleta trawl transect data (n=2) since 1991.



Table 7-3. Comparison of trawl fish metrics with results from the Southern California Regional Survey, Bight 2008 (SCCWRP 2011).

Metric	Trawl Fish		
	Goleta Range	Bight '08 Northern Region Inner Shelf	Below Range?
Biomass (kg)	3.09 - 3.28	0.7 - 4.7	No
Individuals	69 - 181	24 - 467	No
Species	9 - 17	5 - 22	No
Shannon Diversity	1.50 - 1.78	0.5 - 2.31	No






Figure 7-2. Length (cm) frequency distributions for speckled sanddabs (*Citharichthys stigmaeus*) collected from 2003 to 2012 from stations TB3 and TB6 in the Goleta survey area.



7.3.2. Trawl Macroinvertebrates

7.3.2.1. Macroinvertebrate Community Metrics

The averaged macroinvertebrate community metrics and biomass for replicate trawls are presented in Table 7-4, with results by replicate presented in Appendix 10.7 (Tables 10-11 and 10-12). A total of 24 individual invertebrates were collected from both stations combined during the 2013 survey. An average of 4 macroinvertebrates was collected at station TB3 compared to 8 at TB6, but there was no significant difference between sites (Table 7-4). Numbers of species collected averaged 3 at station TB3 and 5 at station TB6, with no significant difference between sites. Biomass was similar at the two sites and there was no significant difference. Shannon Diversity, Simpson Diversity and Margalef Richness were low at both station and there were no significant differences between sites.

7.3.2.2. Species Composition

As with past years, the invertebrates in the 2013 trawls were typical of those found on most southern California near shore soft bottom habitats (Table 7-5). A total of 9 individual taxa were collected in the survey area. The most abundant species collected in the survey area were the orange bigeye octopus (*Octopus californicus*) and the graceful rock crab (*Cancer gracilis*).

7.3.2.3 Macroinvertebrate Community Metrics Compared to Past Surveys

Macroinvertebrate community metrics for 2013 were compared to previous Goleta area surveys starting in 1991 (Figure 7-2). The numbers of individuals and average biomass in 2013 was similar to the previous ten years. Numbers of species, Shannon Diversity and Dominance were also similar to the previous decade. These three metrics declined in 1998 from historic highs and have been relatively stable since. The reasons for these reductions are unclear.

7.3.2.4. Macroinvertebrate Community Metrics Compared to Reference Surveys

Macroinvertebrate community metrics for the 2013 Goleta survey were compared to invertebrate assemblage data collected in the northern region on the inner continental shelf in the southern California bight during the 2008 Southern California Bight Regional Monitoring Survey (SCBRMP) (SCCWRP 2011; Table 7-6). Biomass, numbers of individuals, numbers of species and Shannon Diversity were all within the range of fish assemblages found in the northern region inner shelf.



Table 7-4. Trawled inverts - Summary of biological metrics of invertebrates collected at Stations TB3 and TB6. Comparison between sites by two sample T-test ($p > 0.05$).

Metric	Station	Invertebrates				T-test	
		TB3		TB6		t score	p =
		Avg	SD	Avg	SD		
Individuals		4	3	8	1	-1.79	0.22
Species		3	1	5	1	-1.41	0.29
Biomass (kg)		0.62	0.57	0.74	0.28	-0.27	0.81
Shannon Diversity		0.97	0.39	1.44	0.41	-1.19	0.36
Simpson Diversity		0.58	0.12	0.71	0.14	-0.95	0.44
Margalef Richness		1.44	0.16	1.92	0.52	-0.92	0.46
Schwartz Dominance ¹		3	1	4	1	-1.41	0.29

Bold - Marginally Significant ($0.05 < p < 0.10$)

Bold - Significant ($p < 0.05$)

1. Non-normal data: T-test by Mann-Whitney U test.

Table 7-5. Trawled invertebrate abundance and biomass sorted from most to least abundant.

Trawl TB3				Trawl TB6			
Scientific Name	Common Name	Mean Abundance	Mean Weight (kg)	Scientific Name	Common Name	Mean Abundance	Mean Weight (kg)
<i>Cancer gracilis</i>	graceful rock crab	2	0.39	<i>Octopus californicus</i>	orange bigeye octopus	3	0.12
<i>Octopus californicus</i>	orange bigeye octopus	1	<0.1	<i>Cancer gracilis</i>	graceful rock crab	2	0.13
<i>Pisaster brevispinus</i>	shortspined sea star	1	0.23	<i>Astropecten californicus</i>	California sand star	1	<0.1
<i>Sicyonia ingentis</i>	ridgeback rock shrimp	1	<0.1	<i>Loxorhynchus grandis</i>	sheep crab	1	0.44
	composite weight*		<0.1	<i>Octopus rubescens</i>	red octopus	1	<0.1
				<i>Ophiothrix spiculata</i>	Pacific spiny brittlestar	1	<0.1
				<i>Pyromaia tuberculata</i>	tuberculate pear crab	1	<0.1
					composite weight*		0.05

*Species <0.1 kg are weighed together as a composite weight.






Figure 7-2. Invertebrate community metric annual averages (\pm SD) for Goleta trawl transect data (n=2) since 1991.



Table 7-6. Comparison of trawl invertebrate metrics with results from the Southern California Regional Survey, Bight 2008 (from SCCWRP, 2011).

Metric	Trawl Invertebrate		
	Goleta Range	Bight '08 Northern Region Inner Shelf	Below Range?
Biomass (kg)	0.62 - 0.74	0.0 - 3.0	No
Individuals	4 - 8	3 - 135	No
Species	3 - 5	2 - 20	No
Shannon Diversity	0.97 - 1.44	0.64 - 2.30	No

7.4. Discussion

Results from this trawl survey support past studies that indicated that the discharge from the Goleta Sanitary District's ocean outfall does not appear to be impacting the resident fish or macroinvertebrate communities. This was confirmed by comparing results among stations both near and far from the diffuser, comparing results with historical surveys, and comparing results with other studies being performed in southern California.

A total of 500 individual fish and 24 individual invertebrates were collected from both stations combined during the 2013 survey. There were no statistically significant differences ($p < 0.05$) between stations near to and far from the outfall when metrics for fish or invertebrates total abundance, biomass, and numbers of taxa, diversity and dominance were compared. The only exception to this was for Margalef's Richness which was significantly greater at station TB6 compared to TB3. In addition, both fish and invertebrate population indices measured in 2013 (including abundance, numbers of species and biomass) were within the range of reference sites sampled during the 2008 Southern California Bight Regional Monitoring Program.

As with past years, the fishes and macroinvertebrates caught in the 2013 trawls were typical of those found on most southern California near shore soft bottom habitats. A total of 9 and 17 individual fish taxa were collected at stations TB3 and TB6, respectively. The most abundant species collected at station TB3 and TB6 was the speckled sanddab (*Citharichthys stigmaeus*) followed by the Pacific sanddab (*Citharichthys sordidus*).

A total of 9 unique invertebrate taxa were collected in the trawl area. The most abundant species collected in the survey area were the orange bigeye octopus (*Octopus californicus*) and the graceful rock crab (*Cancer gracilis*).

When the 2013 trawled fish and invertebrate results were compared against past surveys, average abundances, numbers of species, biomass, diversity and dominance were within the ranges of the previous twenty years. This was especially true of the trawled fish community. In contrast, the trawled invertebrate community has been very similar for each biological metric over the past ten years, but prior to 2001 the numbers of invertebrate taxa and diversity were much greater. The reasons for the decrease in trawled invertebrate diversity



are unclear. Since an outfall related impact has never been detected, it is probable that some larger oceanographic condition has influenced this community. Frequent cold water upwelling events which are typical of this coastal region, coupled with warm water El Nino events over the past 15 years may be playing a significant role in the recruitment to and stability of this community.

Although there are no specific numerical limitations regarding trawl animals, the California Ocean Plan (1997) states that:

- *The rate of deposition of inert solids and the characteristics of inert solids in the ocean shall not be changed such that benthic communities are degraded.*
- *The concentration of substances set forth in Chapter IV, Table B, in marine sediments shall not be increased to levels which would degrade indigenous biota.*
- *The concentration of organic materials in marine sediments shall not be increased to levels which would degrade marine life.*
- *Nutrient materials shall not cause objectionable aquatic growths or degrade indigenous biota.*
- *Marine communities, including vertebrate, invertebrate, and plant species, shall not be degraded.*
- *Waste management systems that discharge to the ocean must be designed and operated in a manner that will maintain the indigenous marine life and a healthy and diverse marine community.*
- *Waste discharged to the ocean must be essentially free of: "2) Settleable material or substances that may form sediments which will degrade benthic communities or other aquatic life."*

Based upon spatial and temporal comparisons and analogies with other studies, results of the trawl survey indicate that the discharge is in compliance with the general limitations and that it causes no adverse impact.



CHAPTER 8

Fish and Bivalve Tissue Bioaccumulation

8.1. Background

Outfall discharges can potentially increase contaminant concentrations in sediments and the water column to the extent that marine plant and animal communities are altered, reduced, or eliminated. Harvested fish or invertebrate flesh may become contaminated and unfit for human consumption. Bioaccumulation is a process whereby contaminants are assimilated by organisms, retained and bioconcentrated over time. The degree of bioconcentration is different among species and among toxicants. Biomagnification may also occur when predators eat organisms, resulting in the concentration of contaminants in higher levels of the food chain. In this way, higher-level predators, such as large fish, birds, and mammals can experience chronic toxicity, reproductive failure, or even mortality.

8.2. Materials and Methods

The measure of contaminants in animal tissues was performed with both fish (speckled sanddabs, *Citharichthys stigmaeus*) and invertebrates (California bivalves, *Mytilus californianus*) using two completely different collection procedures. Speckled sanddabs were collected by otter trawl procedures, which are described in Section 7 above. Sanddabs collected in the population trawls were kept, and additional trawls were continued until sufficient total biomass for tissue analysis had been collected. Animals from each of two stations (TB3 between the diffuser and Goleta Point and TB6 at the down coast field control) were placed in plastic zip-lock bags and covered with ice in coolers. Immediately upon return to the laboratory, dorsal muscle and livers were removed from each animal, using standard clean room techniques, and placed in new pre-cleaned glass jars with Teflon-lined caps. All tissue samples were then stored in a freezer until ready to be shipped to the chemistry laboratory (PHYSIS Laboratories in Anaheim, California). Analytical methods were similar to sediments, except that special extraction and clean-up techniques were used to eliminate lipid interferences commonly found in marine animal tissues.

Bivalves were collected from Anacapa Island, California, an area anticipated to be very low in anthropogenic contamination. Prior to deployment these bivalves were cleaned of all debris and growth and held in a pre-cleaned seawater tank at 15° C until use. Bivalves were deployed using three arrays, each composed of a float, line, and anchor. Bivalve cages, made of plastic mesh netting, were attached to the middle of the arrays, so that the bivalves could be suspended at about mid-depth (16 m). The arrays were deployed in duplicate at Stations B3, B4, and B6; located 250, 25, and 3000 m (respectively) from the diffuser. The duplicate array at each station was suspended on a sub-surface buoy and attached to the first array with a 100 meter long line that was weighted to the bottom. Prior to deployment of the arrays in July, laboratory control bivalves were randomly selected and tissues were resected and frozen. In October, each of the three bivalve arrays was successfully retrieved.

Once bivalves were removed from the array, they were placed on ice and returned to the laboratory. Exposed bivalves, as well as bivalves from the original population were cleaned, measured, and weighed. Their tissues were resected, stored, and analyzed, as above.

For the purposes of statistical analysis, all analytes from each of four groups (DDT and its derivatives (i.e. DDD and DDE), PCB's, PAH's, and non-DDT chlorinated pesticides) were combined. Results for individual analytes are presented in Appendix 10-16 and 10-17. All



data were converted to mg/Kg or µg/Kg, dry weight and statistically compared among stations using either t-test for two stations or analysis of variance (ANOVA) for three or more stations (see Section 3.4). When assumptions of parametric statistics could not be met (such as non-normality or excessive variability), the tests were replaced with nonparametric analogues (Aspin-Welch Unequal Variance Test, Mann-Whitney U, and Kruskal-Wallis Rank Test, respectively). Significance was noted when $p \leq 0.05$ and marginal significance was noted when $0.05 < p \leq 0.10$. *A posteriori* tests were utilized for significant ANOVA results to determine which stations were significantly different (see Zar 1996 or Sokal and Rohlf 1981 for a general description of statistical testing).

To compare tissue concentrations to the Office of Environmental Health Hazard Assessment (OEHHA) thresholds (OEHAA 2008) and NOAA Status and Trends mussel watch historical surveys (Kimbrough et al. 2008), Goleta tissue data were converted to wet weight units.

8.3. Results

Table 8-1 lists the physical and general descriptions of the animals utilized in the Goleta bioaccumulation study. Appendix Tables 10-13 and 10-14 lists lengths and weights of organisms, as well as tissue weights. Tables 8-2 to 8-4 and Figures 8-1 and 8-2 present average concentrations for each chemical constituent measured in the three types of animal tissues at each Station. Appendix Table 10-15 lists each constituent by replicate and averages by stations. Figures 8-3 through 8-5 compare historical contamination trends in the three tissue types. Tables 8-5 to 8-6 compare the Goleta tissue chemistry results with reference surveys and state OEHHA thresholds and NOAA status and trends tissue levels. Appendix 10-16 and 10-17 lists the concentrations of the derivatives of total DDT, non-DDT chlorinated hydrocarbons, total PCBs, and total PAHs. General descriptions of all chemical constituents have been presented earlier in Chapter 5, and so will not be repeated here.

8.3.1. Spatial contaminant patterns in tissues

Speckled Sanddabs

A total of 190 speckled sanddabs (*Citharichthys stigmaeus*) were collected for tissue dissections from trawl transects TB3 (n = 82) and TB6 (n = 108), respectively. Average standard lengths (88 and 74 mm, respectively) were similar between sites, while average weight was nearly double at TB3 (13.4 g) compared to TB6 (7.5 g). Dissected tissue weights were greater for muscle tissue (2.5 and 1.3 wet g, respectively) compared to liver (0.3 and 0.2 wet g, respectively).

Of the ten metals measured in sanddab muscle tissue all were above detection except nickel and silver (Table 8-2 and Figure 8-1). Arsenic, cadmium, copper, selenium and zinc were each slightly, but significantly greater by t-test ($p < 0.05$) at station TB3 nearest the outfall, compared to concentrations TB6. Mercury was significantly greater at TB6. Of the groups of complex organic compounds measured in sanddab muscle tissue, total chlordane, total PCBs, arochlors, total HCHs and total PAHs were below detection at both stations. Total DDTs concentrations were doubled and significantly greater at TB6 (21.8 ug/L) compared to TB3 (10.9 ug/L).

Of the ten metals measured in sanddab liver, all were above detection (Table 8-3 and Figure 8-1). Cadmium, lead, mercury, selenium, silver and zinc were slightly, but significantly, greater at TB6 compared to TB3 by t-test ($p < 0.05$). Copper and arsenic were slightly, but significantly greatest at TB3. HCHs, PCBs (at TB3) and arochlors were below detection. Total



DDT, chlordane, total PAHs, benz[a]anthracene, and perylene were significantly greater by t-test at TB3 compared to TB6. Total PCBs and biphenyl were significantly greater at TB6. It should be noted that where significant differences in muscle or liver tissue concentrations were detected by t-test, the differences in average metal concentrations between stations was extremely small.

Bivalves

Of the ten metals measured in bivalve (*Mytilus californianus*) tissue, all were above detection (Table 8-4, Figure 8-1). Chromium and nickel were significantly greater at B3 and B4 compared to B6 by ANOVA ($p < 0.05$). Of the complex organic compounds measured in bivalve tissue, chlordane, total HCHs, total PCBs and arochlors were below detection in each replicate for all stations. Total DDT and total PAHs were detected at each station, but only PAHs were significantly greater at B4 and B6.

8.3.2 Tissue contaminant concentrations compared with past years

Sanddabs

The average concentration of contaminants in sanddab muscle and liver tissues remained within range of previous years (Figures 8-3 and 8-4). Increases in sanddab muscle concentrations of chromium, nickel and silver reported for the 2009 survey returned to lower concentrations in 2010 and remained low thru 2013. Arsenic concentrations increased seven fold in muscle tissue from 2010 (2 mg/dry Kg) to 2011 (15 mg/dry Kg), but dropped to 6 mg/dry Kg thru 2013. Increases in liver DDT and PCB concentrations, which had increased between 2010 and 2011, dropped to lower concentrations from 2012 to 2013.

Bivalves

The average concentration of each contaminant in bivalve tissues remained the same in 2013 (Figure 8-5). Similar to sanddab muscle and liver, silver concentrations which had increased to the greatest concentrations of all past surveys in 2009, decreased in 2010 and remained low thru 2013.

8.3.3 Tissue contaminant concentrations compared with other surveys & State Thresholds & EPA Ranges

The concentrations of the contaminants measured in sanddab and bivalve tissues during the 2013 survey were compared to the concentrations measured at other sites throughout southern California (Table 8-5 and 8-6). Where comparisons were available, sanddab muscle and liver tissues, and mussel tissues were below or within the range of contaminant concentrations reported from other surveys (see references in Table 8-5 and 8-6 footnotes). Sanddab and muscle tissue concentrations of metals and organic constituents did not exceed OEHHA consumption thresholds. Finally, mussel tissue concentrations were in the 'low' range reported by the NOAA Status and Trends Mussel Watch program.

8.4. Discussion

Results from this survey support past studies showing that the Goleta outfall discharge appears not to effect the concentrations of contaminants in the tissues of fish and invertebrates residing in the survey area. Results from the chemical analysis of tissues were compared among stations, compared to past surveys in the area, compared to other studies performed in



southern California, and compared to State thresholds and Federal ranges for concentrations of contaminants in animal tissue. Results for each variable were statistically compared among stations by either t-test or analysis of variance (ANOVA).

The sampling design for fish differed from the design for bivalve arrays. The bivalve sampling plan included a laboratory control (unexposed bivalves from Anacapa Island, CA) and bivalves exposed at three site locations: one station down coast (field control), one station nearest the outfall, and one station up coast and nearest Goleta Point. For fish, there was no laboratory control, and fish were collected from only two locations: one station down coast of the outfall corresponding to the field control, and one up coast of the outfall corresponding to the station nearest Goleta Point.

A total of 15 chemical compounds or groups of compounds were analyzed in speckled sanddab muscle tissue from the two trawl locations. Sanddab muscle tissues had two metals (nickel and silver), as well as total chlordane, total PCBs, arochlors, total HCHs and total PAHs that were each below method detection. Among the remaining compounds, arsenic, cadmium, copper, selenium and zinc were each slightly, but significantly greater at station TB3 nearest the outfall, compared to concentrations measured at TB6. In sanddab liver tissues each metal was above detection at each site, while HCHs, PCBs and arochlors were below detection. Cadmium, lead, mercury, selenium, silver and zinc were slightly, but significantly, greater at TB6 compared to TB3 by t-test, while copper and selenium were significantly greatest at outfall station TB3. Total DDT, chlordane, total PAHs, benz[a]anthracene, and perylene were significantly greater by t-test at TB3 compared to TB6. Total PCBs and biphenyls were significantly greater at TB6. It should be noted that where significant differences in muscle or liver tissue concentrations were detected by t-test, the differences in average metal concentrations between stations was extremely small.

A total of 15 chemical compounds or groups of compounds were analyzed in the whole body tissues of bivalves. Each of the metals was above method detection limits, and chromium and nickel were significantly greater at B3 and B4 compared to B6 by ANOVA. Of the complex organic compounds measured in bivalve tissue, chlordane, total HCHs, total PCBs and arochlors were below detection in each replicate for all stations. Total DDT and total PAHs were detected at each station, but only PAHs were significantly greater at B4 and B6.

Comparison of the 2013 tissue concentrations from the Goleta survey area against results from the past nineteen years revealed that in all cases contaminant concentrations were similar to or less than in past years. Increases in sanddab muscle chromium, nickel and silver reported for the 2009 survey returned to lower concentrations in 2010 and remained low thru 2013. Arsenic concentrations increased seven fold in sanddab muscle tissue from 2010 (2 mg/dry Kg) to 2011 (15 mg/dry Kg), and then dropped to 6 mg/dry Kg thru 2013.

The concentrations of the contaminants measured in sanddab and bivalve tissues during the 2013 survey were compared to the concentrations measured at other sites throughout southern California. Where comparisons were available, sanddab muscle and liver tissues, and mussel tissues were below or within the range of contaminant concentrations reported from other surveys. Sanddab and bivalve tissue concentrations of metals and organic constituents did not exceed OEHHA consumption thresholds. Since the speckled sanddab is not caught for human consumption due to its small size, comparison of its tissue burdens against the OEHHA standard is included to provide context. Finally, bivalve tissue concentrations were in the 'low' range reported by the NOAA Status and Trends Mussel Watch program (Kimbrough et al. 2008).



Although there are no specific numerical limitations regarding trawl animals, the California Ocean Plan (1997) states that:

The natural taste, odor, and color of fish, shellfish, or other marine resources used for human consumption shall not be altered.

The concentration of organic materials in fish, shellfish or other marine resources used for human consumption shall not bioaccumulate to levels that are harmful to human health.

Based upon spatial and temporal patterns and comparisons with other studies, results of the bioaccumulation survey indicate that the discharge is in compliance with the general limitations that it causes no adverse impact.



Table 8-1. Numbers of animals, length (mm), weight (g) and tissues weight (g) in fish and bivalve tissue collected in the Goleta survey area.

Constituent	Replicate	Fish Muscle		Fish Liver		Bivalves			
		T3	T6	T3	T6	Control	B3	B4	B6
Number of Animals		82	108	82	108	45	45	45	45
Average Standard Length (mm)	Mean = S.D. =	88.1 10.6	74.2 8.7	88.1 10.6	74.2 8.7	69.2 7.4	73.6 7.3	75.7 7.1	76.9 8.5
Average Weight/Animal (g)	Mean = S.D. =	13.4 4.9	7.5 3.3	13.4 4.9	7.5 3.3	30.6 8.8	41.3 8.6	40.0 8.3	42.0 10
Average Tissue Weight (g)	Mean = S.D. =	2.5 1.0	1.3 0.6	0.3 0.2	0.2 0.1	10.0 3.7	13.4 3.5	13.1 3.1	14 3.9



Table 8-2. Mean concentrations of speckled sanddab (*Citharichthys stigmaeus*) muscle collected in the Goleta survey area. Comparisons of means determined by T-test ($p < 0.05$).

Constituent	Fish Muscle						T-Test		
	TB3			TB6			n	t	p
	mean	±	SD	mean	±	SD			
Metals (µg/dry g)									
Arsenic	7.323	±	0.216	4.133	±	0.216	3	18.11	<0.01
Cadmium	0.043	±	0.002	0.033	±	0.003	3	4.95	0.01
Chromium	0.063	±	0.027	0.058	±	0.005	3	0.34	0.75
Copper ²	2.025	±	0.820	1.192	±	0.063	3	1.96	0.05
Lead	0.028	±	0.003	0.035	±	0.011	3	-1.21	0.29
Mercury	0.162	±	0.005	0.187	±	0.003	3	-0.70	<0.01
Nickel ²	0.025	±	0.000	0.042	±	0.029	3	-1.00	0.32
Selenium	1.301	±	0.065	0.992	±	0.071	3	6.57	0.01
Silver	0.025	±	0.000	0.025	±	0.000	3	NA	NA
Zinc	19.106	±	0.527	16.365	±	0.993	3	4.22	0.01
Complex Organics (ng/dry Kg)									
DDTs ^{1,2}	10.9	±	0.4	21.8	±	2.5	3	-0.20	0.05
Chlordane ¹	0.0	±	0.0	0.0	±	0.0	3	NA	NA
HCHs ¹	0.0	±	0.0	0.0	±	0.0	3	NA	NA
Aldrin	1.0	±	0.0	1.0	±	0.0	3	NA	NA
Dieldrin	1.0	±	0.0	1.0	±	0.0	3	NA	NA
Heptachlor	1.0	±	0.0	1.0	±	0.0	3	NA	NA
Hexachlorobenzene	1.0	±	0.0	1.0	±	0.0	3	NA	NA
Mirex	1.0	±	0.0	1.0	±	0.0	3	NA	NA
PCBs ¹	0.0	±	0.0	0.0	±	0.0	3	NA	NA
Arochlors ¹	0.0	±	0.0	0.0	±	0.0	3	NA	NA
PAHs ¹	30.7	±	3.2	29.6	±	1.9	3	0.55	0.61
1-Methylnaphthalene	2.4	±	0.2	1.5	±	0.3	3	4.80	0.01
1-Methylphenanthrene ²	2.0	±	0.3	1.9	±	0.0	3	-0.71	0.48
2-Methylnaphthalene	4.4	±	0.5	2.4	±	1.2	3	2.74	0.05
2,3,5-Trimethylnaphthalene	1.0	±	0.0	1.0	±	0.0	3	NA	NA
2,6-Dimethylnaphthalene	3.9	±	0.9	1.7	±	0.4	3	3.80	0.02
Acenaphthene ²	1.0	±	0.0	2.3	±	0.8	3	-0.21	0.04
Biphenyl	2.3	±	0.3	1.7	±	0.5	3	1.75	0.15
Benz[a]anthracene	1.0	±	0.0	1.0	±	0.0	3	NA	NA
Benzo[b]fluoranthene	1.0	±	0.0	1.0	±	0.0	3	NA	NA
Benzo[e]pyrene	1.0	±	0.0	1.0	±	0.0	3	NA	NA
Benzo[g,h,i]perylene	1.0	±	0.0	1.0	±	0.0	3	NA	NA
Fluoranthene	5.3	±	2.7	4.6	±	0.6	3	0.43	0.69
Napthalene	13.6	±	1.5	6.3	±	1.1	3	6.78	0.00
Perylene	1.0	±	0.0	1.0	±	0.0	3	NA	NA

1. Complex Organic derivatives are listed in Table 10-16.
2. Non-normal data. Statistics by Mann-Whitney U Test.



Table 8-3. Mean concentrations of speckled sandab (*Citharichthys stigmaeus*) liver collected in the Goleta survey area. Comparisons of means determined by T-test ($p < 0.05$).

Constituent	Fish Liver						T-Test		
	TB3			TB6			n	t	p
	mean	±	SD	mean	±	SD			
Metals (µg/dry g)									
Arsenic	8.434	±	0.383	6.113	±	0.101	3	10.14	<0.01
Cadmium	8.056	±	0.099	9.852	±	0.232	3	-12.32	<0.01
Chromium ²	0.095	±	0.064	0.113	±	0.009	3	-0.65	0.15
Copper	10.845	±	0.145	10.233	±	0.216	3	4.07	0.02
Lead	0.442	±	0.041	0.693	±	0.027	3	-8.84	<0.01
Mercury	0.100	±	0.003	0.123	±	0.003	3	-9.31	<0.01
Nickel	0.089	±	0.009	0.098	±	0.017	3	-0.83	0.45
Selenium	5.325	±	0.431	7.521	±	0.179	3	-8.15	<0.01
Silver	0.098	±	0.004	0.291	±	0.008	3	-37.56	<0.01
Zinc	66.349	±	1.681	72.619	±	1.032	3	-5.51	0.01
Complex Organics (ng/dry Kg)									
DDTs ^{1,2}	912.9	±	52.5	423.9	±	15.0	3	15.52	<0.01
Chlordane ^{1,2}	6.6	±	5.8	0.0	±	0.0	3	1.55	0.12
HCHs ¹	0.0	±	0.0	0.0	±	0.0	3	NA	NA
Aldrin	1.0	±	0.0	1.0	±	0.0	3	NA	NA
Dieldrin	1.0	±	0.0	1.0	±	0.0	3	NA	NA
Heptachlor	1.0	±	0.0	1.0	±	0.0	3	NA	NA
Hexachlorobenzene	1.0	±	0.0	1.0	±	0.0	3	NA	NA
Mirex	1.0	±	0.0	1.0	±	0.0	3	NA	NA
PCBs ^{1,2}	0.0	±	0.0	34.6	±	4.2	3	-2.09	0.04
Arochlors	0.0	±	0.0	0.0	±	0.0	3	NA	NA
PAHs ¹	219.2	±	6.5	127.3	±	23.0	3	6.67	<0.01
1-Methylnaphthalene	6.2	±	1.7	8.6	±	2.6	3	-1.32	0.26
1-Methylphenanthrene	13.1	±	2.1	9.9	±	3.7	3	1.30	0.26
2-Methylnaphthalene	7.5	±	9.8	16.9	±	2.6	3	-1.61	0.02
2,3,5-Trimethylnaphthalene ²	1.0	±	0.0	5.2	±	7.3	3	-1.00	0.32
2,6-Dimethylnaphthalene	11.3	±	2.7	8.6	±	1.4	3	1.58	0.19
Acenaphthene	1.0	±	0.0	1.0	±	0.0	3	NA	NA
Biphenyl	6.6	±	0.7	10.8	±	2.2	3	-3.21	0.03
Benz[a]anthracene ²	73.2	±	3.1	1.0	±	0.0	3	2.09	0.04
Benzo[b]fluoranthene	1.0	±	0.0	1.0	±	0.0	3	NA	NA
Benzo[e]pyrene	1.0	±	0.0	1.0	±	0.0	3	NA	NA
Benzo[g,h,i]perylene	1.0	±	0.0	1.0	±	0.0	3	NA	NA
Fluoranthene ²	24.1	±	10.5	21.1	±	0.2	3	0.66	0.51
Napthalene	37.9	±	2.7	38.2	±	9.8	3	-0.06	0.96
Perylene	35.5	±	16.1	1.0	±	0.0	3	2.09	0.04

1. Complex Organic derivatives are listed in Table 10-16.
2. Non-normal data. Statistics by Mann-Whitney U Test.



Table 8-4. Heavy metals and complex organics in California bivalve (*Mytilus californianus*) tissues. Comparisons of means by ANOVA ($p < 0.05$).

Constituent	Bivalve Tissue											
	B3			B4			B6			ANOVA		
	mean	±	SD	mean	±	SD	mean	±	SD	n	F	p
Metals (µg/dry g)												
Arsenic	9.285	±	0.403	9.426	±	0.264	9.448	±	0.312	3	0.21	0.81
Cadmium	5.260	±	0.515	5.864	±	0.203	5.256	±	0.248	3	3.00	0.13
Chromium	0.891	±	0.061	0.931	±	0.031	0.749	±	0.030	3	14.97	<0.01
Copper	6.818	±	0.283	7.012	±	0.062	7.088	±	0.154	3	1.61	0.27
Lead	1.419	±	0.130	1.285	±	0.077	1.225	±	0.062	3	0.75	0.51
Mercury	0.041	±	0.002	0.043	±	0.001	0.040	±	0.003	3	0.75	0.51
Nickel	0.809	±	0.024	0.895	±	0.047	0.734	±	0.036	3	14.11	0.01
Selenium	2.842	±	0.182	3.166	±	0.105	3.095	±	0.166	3	3.66	0.09
Silver	0.237	±	0.046	0.257	±	0.043	0.241	±	0.048	3	0.16	0.86
Zinc	123.339	±	3.395	124.759	±	3.000	130.636	±	3.851	3	3.81	0.09
Complex Organics (ng/dry Kg)												
DDTs ¹	18.7	±	0.8	21.3	±	4.0	19.3	±	3.7	3	0.56	0.60
Chlordane ¹	0.0	±	0.0	0.0	±	0.0	0.0	±	0.0	3	NA	NA
HCHs ¹	0.0	±	0.0	0.0	±	0.0	0.0	±	0.0	3	NA	NA
Aldrin	1.0	±	0.0	1.0	±	0.0	1.0	±	0.0	3	NA	NA
Dieldrin	1.0	±	0.0	1.0	±	0.0	1.0	±	0.0	3	NA	NA
Heptachlor	1.0	±	0.0	1.0	±	0.0	1.0	±	0.0	3	NA	NA
Hexachlorobenzene	1.0	±	0.0	1.0	±	0.0	1.0	±	0.0	3	NA	NA
Mirex	1.0	±	0.0	1.0	±	0.0	1.0	±	0.0	3	NA	NA
PCBs ¹	0.0	±	0.0	0.0	±	0.0	0.0	±	0.0	3	NA	NA
Arochlors ¹	0.0	±	0.0	0.0	±	0.0	0.0	±	0.0	3	NA	NA
PAHs ¹	61.1	±	3.2	72.0	±	4.9	74.9	±	5.9	3	6.97	0.03
1-Methylnaphthalene	2.4	±	0.4	3.2	±	0.6	3.5	±	0.5	3	4.48	0.06
1-Methylphenanthrene	6.4	±	0.4	6.7	±	0.7	5.8	±	0.1	3	3.12	0.12
2-Methylnaphthalene	5.8	±	2.1	7.9	±	0.3	7.6	±	2.2	3	1.30	0.34
2,3,5-Trimethylnaphthalene	1.3	±	0.4	1.2	±	0.2	1.2	±	0.3	3	0.08	0.92
2,6-Dimethylnaphthalene	2.4	±	0.6	3.4	±	0.4	3.0	±	0.2	3	4.18	0.07
Acenaphthene	1.0	±	0.0	1.0	±	0.0	1.0	±	0.0	3	NA	NA
Biphenyl	2.3	±	0.1	3.2	±	0.4	3.2	±	0.3	3	8.14	0.02
Benz[a]anthracene	1.0	±	0.0	1.0	±	0.0	1.0	±	0.0	3	NA	NA
Benzo[b]fluoranthene	1.0	±	0.0	1.0	±	0.0	1.0	±	0.0	3	NA	NA
Benzo[e]pyrene	1.0	±	0.0	1.0	±	0.0	1.0	±	0.0	3	NA	NA
Benzo[g,h,i]perylene	1.0	±	0.0	1.0	±	0.0	1.0	±	0.0	3	NA	NA
Fluoranthene	14.6	±	2.6	13.9	±	6.4	15.1	±	1.1	3	0.07	0.94
Naphthalene	10.7	±	1.6	13.5	±	2.4	13.5	±	1.8	3	2.00	0.22
Perylene	1.0	±	0.0	1.0	±	0.0	1.0	±	0.0	3	NA	NA

1. Complex Organic derivatives are listed in Table 10-17.
2. Non-normal data. Statistics by Kruskal-Wallis Test.



Figure 8-1. Metal concentrations (mg/dry Kg) measured in fish muscle and liver tissues (Stations TB3 and TB6), and bivalves (Stations B3, B4, B6 and lab control).




Figure 8-2. Organic concentrations ($\mu\text{g}/\text{dry Kg}$) measured in fish muscle and liver tissues (Stations TB3 and TB6), and mussels (B3, B4, B6 and lab control).




Figure 8-3. Contaminants (mg/dry Kg) measured in speckled sanddab muscle (*Citharichthys stigmaeus*) from Goleta since 1991 (mean \pm SD, n=6).




Figure 8-4. Contaminants (mg/dry Kg) measured in speckled sanddab liver (Citharichthys stigmaeus) from Goleta since 1991 (mean \pm SD).




Figure 8-5. Contaminants (mg/dry Kg) measured in whole bivalves (*Mytilus californianus*) from Goleta since 1991 (mean \pm SD, n = 3).




Table 8-5. Comparison of Goleta tissue chemistry with results from other studies (ug/wet g) and state and federal limits.

Constituent	GOLETA S.D.		Reference µg/g Wet Weight Stations ¹	OEHHA²	
	Means	Ranges		µg/g Wet Weight FCG³	ATL⁴
<u>Fish Muscle</u>					
Arsenic	1.123	0.761 - 1.481	42.2 - 57.8	---	---
Cadmium	0.007	0.006 - 0.009	<0.01 - 0.045	---	---
Chromium	0.012	0.008 - 0.018	0.08 - 2.8	---	---
Copper	0.315	0.226 - 0.582	0.45 - 2.4	---	---
Lead	0.006	0.005 - 0.009	1.2	---	---
Mercury	0.034	0.031 - 0.037	0.36 - 0.78	0.22	≤0.07 ⁵
Nickel	0.007	0.005 - 0.015	0.4 - 5.1	---	---
Selenium	0.225	0.180 - 0.269	2.8 - 3.95	7.4	≤2.5
Silver	0.005	0.005 - 0.005	<0.005 - 1.4	---	---
Zinc	3.476	2.994 - 3.836	12.4 - 30.5	---	---
DDTs	0.003	0.002 - 0.005	0.005 - 2.15	0.021	≤0.52
Chlordane	0.000	0.000 - 0.000	---	0.0056	≤0.052
PCBs	0.000	0.000 - 0.000	0.005 - 2.7	0.0036	≤0.021
PAHs	0.006	0.005 - 0.007	---	---	---
<u>Fish Liver</u>					
Arsenic	2.17	1.80 - 2.62	---	---	---
Cadmium	2.67	2.37 - 3.00	---	---	---
Chromium	0.03	0.01 - 0.05	0.5	---	---
Copper	3.14	3.00 - 3.28	---	---	---
Lead	0.17	0.12 - 0.21	---	---	---
Mercury	0.03	0.03 - 0.04	---	---	---
Nickel	0.03	0.03 - 0.04	---	---	---
Selenium	1.91	1.48 - 2.28	---	---	---
Silver	0.06	0.03 - 0.09	---	---	---
Zinc	20.71	19.22 - 21.92	---	---	---
DDTs	0.199	0.121 - 0.285	28	---	---
Chlordane	0.001	0.000 - 0.003	---	---	---
PCBs	0.005	0.000 - 0.011	4	---	---
PAHs	0.052	0.030 - 0.067	---	---	---

1. Sources: SWRCB 1978, 1988 (EDL 85); SCCWRP 1975, 1976, 1977, 1982, 1998c; Short & Harris 1996; Brown & Caldwell 1997; NOAA 1991, OEHHA 1991

2. OEHHA, 2008

3. Fish Contamination Goal (FCG)

4. Advisory Tissue Levels (ATLs) , most conservative tissue consumption threshold based on cancer or non-cancer risk.

5. Mercury ATL for women aged 18-45 years & children aged 1-17 years (OEHHA 2008).



Table 8-6. Comparison of Goleta tissue chemistry with results from other studies (ug/wet g).

Constituent	GOLETA S.D.		Reference µg/g Wet Weight Stations ¹	OEHHA ²		NOAA Status & Trends, 1986 to 2005		
	Means	µg/g Wet Weight Ranges		µg/g Wet Weight FCG ³	ATL ⁴	low	medium	high
<u>Mussel Tissue</u>								
Arsenic	1.69	1.35 - 1.85	16.0 - 23.8	---	---	5 - 11	12 - 22	23 - 41
Cadmium	0.99	0.79 - 1.16	1.9 - 54	---	---	0 - 3	4 - 9	10 - 20
Chromium	0.15	0.10 - 0.18	1.23 - 3.9	---	---	---	---	---
Copper	1.28	1.08 - 1.38	4.0 - 21.8	---	---	5 - 16	17 - 39	40 - 857
Lead	0.24	0.21 - 0.30	1.09 - 11	---	---	0 - 3	4 - 6	7 - 13
Mercury	0.01	0.01 - 0.01	0.01 - 0.4	0.22	≤0.07 ⁵	0.00 - 0.17	0.18 - 0.35	0.36 - 1.28
Nickel	0.14	0.09 - 0.18	3.2 - 5.3	---	---	0 - 5	6 - 14	15 - 44
Selenium	0.54	0.42 - 0.62	2.70 - 4.57	7.4	≤2.5	---	---	---
Silver	0.04	0.03 - 0.06	0.36 - 0.7	---	---	---	---	---
Zinc	22.20	15.91 - 25.67	133 - 336	---	---	48 - 139	140 - 320	321 - 11500
DDTs	0.0052	0.0030 - 0.0100	0.017 - 0.35	0.21	≤0.52	0 - 0.112	0.113 - 0.286	0.287 - 0.520
Chlordane	0.0000	0.0000 - 0.0000	---	0.0056	≤0.19	0 - 0.008	0.009 - 0.020	0.021 - 0.049
PCBs	0.0000	0.0000 - 0.0000	0.017 - 0.35	0.0036	≤0.021	0.003 - 0.153	0.154 - 0.478	0.479 - 1.413
PAHs	0.0133	0.0112 - 0.0152	0.81	---	---	0.063 - 1.187	1.118 - 4.434	4.435 - 7.561

¹ Sources: SWRCB 1978, 1988 (EDL 85); SCCWRP 1975, 1976, 1977, 1982, 1998c; Short & Harris 1996; Brown & Caldwell 1997; NOAA 1991, OEHHA 1991

² OEHHA, 2008

³ Fish contaminant goals; based on cancer and non-cancer risk using an 8 oz/week consumption rate.

⁴ Advisory tissue levels; based on cancer and non-cancer risk using an 8 oz/week consumption rate

1. Sources: SWRCB 1978, 1988 (EDL 85); SCCWRP 1975, 1976, 1977, 1982, 1998c; Short & Harris 1996; Brown & Caldwell 1997; NOAA 1991, OEHHA 1991

2. OEHHA, 2008

3. Fish contaminant goals; based on cancer and non-cancer risk using an 8 oz/week consumption rate.

4. Advisory tissue levels; based on cancer and non-cancer risk using an 8 oz/week consumption rate (OEHHA 2008).



CHAPTER 9

COLLECTION SYSTEM ANNUAL SUMMARY

Background

Sanitary sewer overflows associated with the Goleta Sanitary District's collection system are subject to the online reporting and notification requirements set forth in the Statewide General Waste Discharge Requirements for Sanitary Sewer Systems Order NO. 2006-0003-DWQ. The Goleta Sanitary District has enrolled under the statewide waste discharge requirement for sanitary sewer systems.

GSD completed the Sanitary Sewer Management Plan (SSMP) in December 2006 and reviews and revises the SSMP annually, as needed. The District's SSMP was updated in September of 2013 in accordance with SWRCB Order No. WQ 2013-0058 – EXEC MRP.

This annual report summarizes all lift station and collection system overflows that occurred during 2013 and includes, if any, the cause, corrective actions taken and corrective actions planned. In conjunction with the annual report the District will conduct the annual SSMP update. The update is a part of the wastewater collection system management plan and requires the District to conduct an internal audit to evaluate the wastewater collection system management plan and delineate steps the District will take to correct any deficiencies that are found.

Annual Reporting Requirement

This chapter is included as part of the wastewater treatment plant annual report.

Summary of 2013 Spills

Lift Station Overflows

There were no lift station overflows that occurred within the Goleta Sanitary District service area during 2013.

Collection System Overflows

There was one collection system overflow that occurred within the Goleta Sanitary District service area during 2013.

This overflow occurred on November 15 as a result of dislodged roots and debris from a private lateral entering into the District's sewer main line. Approximately 100 gallons were calculated to have spilled into a dirt area around a District manhole in an easement area. Corrective actions taken include the cleaning and CCTV inspection of the District sewer line, cleaning and removal of debris from around the spill site and verification that there were no other readily apparent issues that would contribute to a repeat spill at this location.

Corrective planned action measures include the increased monitoring of this sewer line and informing the contractor who that dislodged the roots and debris to contact the District whenever similar work would be performed in the future, in accordance with District procedures. This sewer line was rehabilitated in 2006.

Discussion

The Goleta Sanitary District's wastewater collection system management plan has been completed and complies with all of the requirements of MRP No. R3-2010-0012. All detailed tasks have been addressed in a timely manner and the collection system has complied with all requirements of the monitoring and reporting program.

OUTFALL DIVE SURVEY

Aquatic Bioassay biologists conducted underwater dive surveys and underwater videos of the outfall pipe and diffuser from the Goleta Sanitary District Wastewater Treatment Plant on October 30th, 2013. The purposes of the survey were to inspect the physical integrity of the outfall pipe and associated armor rock and note any impediments to flow from the 36 diffuser ports. Aquatic Bioassay biologists also assessed the presence of attached and mobile marine organisms that were associated with the outfall and the diffuser.

Materials and Methods

Five divers, using Sony 2100 Camcorders enclosed in Gates underwater housings with attached NiteRider underwater lights, conducted the survey. Once the outfall had been located by global positioning (GPS) and bottom finder, a buoy, attached to a line and a weight, was deployed over the side. Divers entered the water, descended down the line, swam to the diffuser terminus, and began filming. At the end of each dive, a lift float was deployed as a marker for the subsequent dive. On deck between dives, the camera was removed from the housing, the footage was inspected, batteries were replaced, and the housing was reassembled. A total of seven dives were completed for the video: diffuser, west and east ports (100 ft. to 70 ft.); deep outfall (70 ft. to 40 ft.); middle outfall (40 ft. to 20 ft.), and shallow outfall (20 ft. to surf zone).

The footage was downloaded to computer files, edited using *Adobe Premiere* software, and then transferred to DVD. DVDs were then reviewed by the survey team to assess conditions of the outfall. The video is arranged from the deepest part of the dives (outfall terminus) to the shallowest part of the dives (outfall beginning).

Results

Outfall dive surveys were conducted between approximately 0800 and 1530 hours on October 30th, 2013 aboard the research vessel *Hey Jude*. Weather conditions were clear with a 10 knot wind from the west, northwest (315°) and a 4 to 6 ft swell from the southwest (245°). Water color was green with high turbidity. Visibility at the terminus of the diffuser (100 feet) was ≤ 1 meter. There was a thermocline at approximately 14 m.

Diffuser Section (Depth: 100 TO 70 ft)

Physical Description

The pipe survey was conducted in October in hopes that water quality would be optimal for taking video footage of the pipe. This year's visibility was poor, ranging from 0 to 1 meter. The diffuser section contains 34 lateral and two terminal discharge ports. The lateral ports are alternately arranged 17 on each side of the diffuser. The end of the pipe is closed except for the two terminal ports, which are situated one above the other. Divers cleared an obstruction on the upper port of the terminus cap, after which the flow from both the upper and lower terminal ports was strong.

Lateral ports were observed and videotaped, starting at the terminus and moving shoreward on the east side of the pipe, then from the terminus down the west side until the most shoreward east port was identified at the beginning of the diffuser. This year, nine ports on the western side of the diffuser were partially blocked with debris. The debris was successfully cleared by the divers so that the ports were flowing freely. All other lateral



ports were flowing freely. Along the length of the diffuser pipe, no evidence of leaks, damage, erosion, holes, or cracks were observed.

An approximately one meter high bed of armor rock supports the diffuser section. Intermittent observations of the supporting armor rock revealed a stable bed of rock with little displacement throughout the diffuser section. Probably during initial construction, the diffuser section appears to have been rotated counter-clockwise (as if one were facing the terminus). Thus, the line across east and west diffuser ports is not parallel to the sea floor, and west ports are about 30 cm lower than east ports. Armor rock covers the outfall from the shoreward beginning of the diffuser to the shoreward beginning of the outfall in very shallow water. The thickness of the armor rock is about one meter.

Biological Description

Because of the depth and relative low light at the diffuser (100 ft), algal species are typically scarce. Algae that were present included the kelp *Desmarestia ligulata* a tubular and leafy red alga (Rhodophyta) and the Turkish Towel (*Gigartina sp.*). Among invertebrates; brown cup coral (*Paracyathus sternsi*), colonial strawberry anemones (*Corynactis californica*), red gorgonian (*Lophogorgia chilensis*) and various species of colonial hydroids and bryozoans dominated. Tube worms and especially the strawberry anemones were commonly observed surrounding the diffuser ports. Sea stars (*Pisaster sp.*), batstars (*Patiria miniata*), blacksmith (*Chromis punctipinnis*) and rockfish (*Sebastes sp.*) were observed either on the pipe, or in its immediate vicinity.

Deep Outfall Section (Depth: 70 TO 40 ft)

Physical Description

Throughout the dive survey, the outfall was completely covered by approximately one-meter layer of armor rock. Visibility was very poor in this section. The rock covered pipe extended vertically from the sea floor for about 2 to 3 meters and laterally for about 6 to 7 meters. The armor rock bed appeared stable with little displacement throughout this section. No obvious leaks or discoloration were observed from the armor rock covering the top or sides of the outfall pipe.

Biological Description

On this section, crustose coralline alga (Rhodophyta), foliose red algae (*Gigartina sp.*) dominated the algal community. Among invertebrates, the most abundant were the red gorgonian (*Lophogorgia chilensis*), colonial strawberry anemones (*Corynactis californica*), several species of bryozoans, bat stars (*Patria miniata*), giant sea stars (*Pisaster giganteus*), the giant keyhole limpet (*Megathura crenulata*), red urchin (*Strongylocentrotus franciscanus*) and the wavy top turban (*Megastraea undosa*). Due to the low visibility, fish species were difficult to identify, however several juvenile fishes and adult fish species were observed including sargo (*Anisotremus davidsoni*), sheephead (*Orthopristis cantharinus*) and blacksmith (*Chromis punctipinnis*).

Middle Outfall Section (Depth: 40 TO 20 ft)

Physical Description

As with the previous section, this outfall section was covered by about one meter of armor rock. The armor rock covered pipe extended horizontally and laterally as above. The armor



rock bed appeared stable with little displacement throughout this section. No obvious leaks or discoloration were observed from the armor rock covering the top or sides of the outfall pipe.

Biological Description

This section supported a giant kelp forest (*Macrocystis pyrifera*) in past years, but similar to 2009, 2010, 2011 and 2012 the density of the kelp was less than in past surveys (MBC 1997, 1998; Aquatic Bioassay 1999 to 2008). The armor rock on this section was populated by large densities of purple sea urchins (*Strongylocentrotus purpuratus*) whose favorite food source is giant kelp. It appears that during the preceding three years the purple urchin population had thinned the kelp forest residing on the outfall pipe. This growth and predation cycle is typical on California rocky reefs and it is probable that during the next two to three years the kelp forests will reestablish on the outfall pipe.

Other dominant algae in this pipe section included foliose red algae (*Gigartina sp.*) and crustose coralline algae. Among the macroinvertebrates, the giant keyhole limpets (*Megathura crenulata*), giant sea stars (*Pisaster brevispinus*), purple sea urchins (*Strongylocentrotus purpuratus*) and the feather boa hydroid (*Aglaophenia struthionides*) were most dominant. Fish species observed at this depth included blacksmith (*Chromis punctipinnis*), kelp bass (*Paralabrax clathratus*), female sheephead (*Semicossyphus pulcher*), lingcod (*Ophiodon elongates*) and opaleye (*Girella nigricans*).

Shallow Outfall Section (Depth: 20 TO 12 ft)

Physical Description

Similar to the deeper portions of the outfall, this section was similarly covered by about one meter of armor rock. Visibility in this section was extremely poor. The armor rock covered pipe extended vertically and laterally as above. As with other sections, the armor rock bed appeared stable with little displacement throughout this section. No obvious leaks or discoloration were observed from the armor rock covering the top or sides of the outfall or from the pipe itself where it was exposed.

Biological Description

Giant kelp, leafy red alga, and coralline crustose alga were present among algae, but as discussed in the previous section, were not as dense as in previous surveys. Among invertebrates, purple urchins (*Strongylocentrotus purpuratus*) were observed in large numbers. Other invertebrates included giant sea stars (*Pisaster brevispinus*), red sea urchins (*Strongylocentrotus franciscanus*), and giant keyhole limpets (*Megathura crenulata*). A few different species of fish were observed in low abundances, including kelp bass (*Paralabrax clathratus*) painted greenling (*Oxylebius pictus*)



Discussion

During the diffuser dive survey, 36 diffuser ports were carefully inspected for flow and general efficiency. This year, nine of the ports were obstructed with debris. The debris was successfully cleared and all of the ports were flowing freely. The remainder of the outfall pipe was inspected for damage, leaks or evidence of leaks and general stability of the pipe and armor rock. Inspection of the outfall yielded no evidence of damage, holes, cracks, or erosion. The pipe and associated armor rock appeared stable with little or no displacement.

The outfall continues to support a rocky reef community typical of other areas on the central California coast. A visual survey yielded numerous different species of plants, macroinvertebrates, and fishes. A number of species of fish were represented by juvenile or larval forms, which indicates that recruitment has been occurring. Fish appeared healthy, with no evidence of deformities, tumors, fin rot, or lesions.

During past surveys the 40 to 20 foot outfall section supported a giant kelp forest (*Macrocystis pyrifera*) that was extremely dense (MBC 1997, 1998; Aquatic Bioassay 1999 to 2008). As in 2009, 2010, 2011, and 2012 this year the density of the kelp was less than in past surveys. The armor rock on this section was populated by large densities of purple sea urchins (*Strongylocentrotus purpuratus*) whose favorite food source is giant kelp. The figure to the right shows purple urchins eating a giant kelp holdfast. During the preceding years the purple urchin population had thinned the kelp forest residing on the outfall pipe through predation (Tegner et al. 1995). Once the kelp plant holdfast is weakened, storms act to break the plant free. This growth and predation cycle is typical on California rocky reefs. Recovery of the kelp forest on the Goleta outfall pipe will be assessed during the next several years.



10.0 APPENDICES



10.1. References



- Allen, M.J. 1982. Functional structure of soft-bottom fish communities of the southern California shelf. Ph.D. dissertation. University of California, San Diego. La Jolla, CA. 577 p.
- American Public Health Association. See APHA.
- Anderson, B., J. Hunt, B. Phillips, R. Fairey, C. Roberts, J. Oakden, H. Puckett, M. Stevenson, R. Tjeerderma, E. Long, C. Wilson, and J.M. Lyons. 2001. Sediment quality in Los Angeles Harbor, USA: a triad Assessment. *Env. Tox. & Chem.* Vol. 20, No. 2, 359-70.
- Anikouchine, W.A., and R.W. Sternberg. *The World Ocean, an Introduction to Oceanography*. Prentice-Hall, Inc. Englewood Cliffs, NJ. 338 p.
- APHA, 2012. *Standard methods for the Examination of Water and Wastewater*. Am. Publ. Health Assn., Am. Water Work Assn., Water Poll. Control Fed. 22nd ed.
- Aquatic Bioassay. 1997. City of Oxnard Receiving Water Monitoring Report, Summary Report 1996.
- Aquatic Bioassay. 1999 to 2010. Goleta Sanitary District Annual Receiving Water Monitoring Report. Prepared for the Goleta Sanitary District, Goleta, CA.
- Aquatic Bioassay. 2007a. Goleta Sanitary District Quarterly Receiving Water Monitoring Report. January 2007. Prepared for the Goleta Sanitary District, Goleta, CA.
- Aquatic Bioassay. 2007b. Goleta Sanitary District Annual Receiving Water Monitoring Report. April 2007. Prepared for the Goleta Sanitary District, Goleta, CA.
- Aquatic Bioassay. 2007c. Goleta Sanitary District Annual Receiving Water Monitoring Report. July 2007. Prepared for the Goleta Sanitary District, Goleta, CA.
- Aquatic Bioassay. 2007d. Goleta Sanitary District Annual Receiving Water Monitoring Report. October 2007. Prepared for the Goleta Sanitary District, Goleta, CA.
- Battelle Ocean Sciences. 1980. Gonadal index and histopathology for the east and west coasts used in the national status and trends project. R.E. Hillman, Battelle Ocean Science, Duxbury, MA.
- Boesch, D. 1977. Application of numerical classification in ecological investigations of water pollution. EPA600/377/033.
- Bray, J.R., and J.T. Curtis. 1957. An ordination of the upland forest communities of southern Wisconsin. *Ecol. Monog.* 27:325-49.
- Brillouin, L. 1962. *Science and Information Theory*. 2nd Ed. Academic Press. NY.
- Brown and Caldwell. 1997. Annual analyses reports, 301 (h) monitoring and reporting program. Prepared for Goleta Sanitary District. Brown and Caldwell, Irvine, Calif.



- Clifford, H.T. and W. Stephenson. 1975. *An Introduction to Numerical Classification*. Academic Press. NY. 229 p.
- Daily, M.D., D.J. Reish, and J.W. Anderson. 1993. *Ecology of the Southern California Bight, A Synthesis and Interpretation*. Univ. Calif. Press, Berkeley. 926 p.
- Dugdale, R.C., Wilkerson, F.P., 1989. New production in the upwelling center at Point Conception, California – temporal and spatial patterns. *Deep-Sea Research Part A – Oceanographic Research Papers* 36 (7), 985–1007.
- Emery, K.O. 1960. *The Sea off Southern California*. J. Wiley & Sons, Inc. New York. 366 p.
- Gray, J.S. 1981. *The Ecology of Marine Sediments, An Introduction to the Structure and Function of Benthic Communities*. Cambridge Univ. Press, Cambridge. 185 p.
- Hanninen, J., I. Vuorinen and P. Hjelt. 2001. Climatic factors in the Atlantic control the oceanographic and ecological changes in the Baltic Sea. *Limnol. Oceanogr.*, 46(7), 2001, 1774–1780.
- Hickey, B.M., 1998. Coastal oceanography of western North America from the tip of Baja California to Vancouver Island. In: Robinson, A.R., Brink, K.H. (Eds.), *The Sea*, vol. 11. John Wiley & Sons, Inc, New York, pp. 345–393.
- Horne, R.A. 1969. *Marine Chemistry, The Structure of Water and the Chemistry of the Hydrosphere*. Wiley-Interscience. John Wiley & Sons. New York. 568 p.
- Kimbrough, K. L., W. E. Johnson, G. G. Lauenstein, J. D. Christensen and D. A. Apeti. 2008. An Assessment of Two Decades of Contaminant Monitoring in the Nation's Coastal Zone. Silver Spring, MD. NOAA Technical Memorandum NOS NCCOS 74. 105 pp.
- Laws, E.A. 1981. *Aquatic Pollution*. John Wiley & Sons, Inc. New York. 482 p.
- Long, E.R., D.D. MacDonald, S.L. Smith, and F.D. Calder. 1995. Incidence of adverse biological effects within ranges of chemical concentrations in marine and estuarine sediments. *Environ. Mgmt.* 19:81-97.
- Long, E.R. and L.G. Morgan. 1990. The Potential for Biological Effects of Sediment-Sorbed Contaminants Tested in the National Status and Trends Program. NOAA Tech. Mem. NOS OMA 52. 175p.
- Margalef, R. 1958. Information theory in ecology. *Gen. Syst.* 3:36-71.
- Mearns, A.J., M. Matta, G. Shigenaka, D. MacDonald, M. Bucbman, H. Harris, J. Golas, and G. Lauenstein. 1991. Contaminant trends in the Southern California Bight: inventory and Assessment. NOAA Technical Memorandum NOS/ORCA 62. 398p.
- Moore, M., and A.J. Mearns. 1978. Coastal fish data base of the Southern California Coastal Water Research Project. Southern California Coastal Water Research Project. El Segundo, CA. 13 p.



- NOAA. 1991. The Potential for Biological Effects of Sediment-Sorbed Contaminants Tested in the National Status and Trends Program. NOAA Tech. Mem. NOS OMA 52. 175 p.
- NOAA. 1999. National status and trends program. <http://seaserver.nos.noaa.gov/projects/nsandt/rawdata.html>.
- OEHHA. 1991. A study of chemical contamination of marine fish from southern California. II. Comprehensive Study. G.A. Pollock, I.J. Uhaa, A.M. Fan, J.A. Wisniewski, I. Witherell.
- OEHHA (Office of Environmental Health Hazard Assessment) Klasing, S. and R. Brodberg). 2008. Development of fish contaminant goals and advisory tissue levels for common contaminants in California sport fish: chlordane, DDTs, dieldrin, methylmercury, PCBs, selenium, and toxaphene. Pesticide and Environmental Toxicology Branch, Office of Environmental Health Hazard Assessment, California Environmental Protection Agency. Sacramento, CA.
- Office of Environmental Health Hazard Assessment, California EPA. See OEHHA.
- Ottersen, G. and N.C. Stenseth. 2001. Atlantic climate governs oceanographic and ecological variability in the Barents Sea. *Limnol. Oceanogr.* 46(7), 1774-1780.
- Pearson, T., and R. Rosenberg. 1978. Macrobenthic succession in relation to organic enrichment and pollution of the marine environment. *Oceanogr. Mar. Biol.* 16:229-311.
- Plumb, R.H. 1981. Procedures for Handling and Chemical Analysis of Sediment and Water Samples. USEPA Contract No. 48-05-572010.
- SCCWRP. 1973. The Ecology of the Southern California Bight: Implications for Water Quality Management. SCCWRP TR104.
- SCCWRP. 1975. Decreases of DDT and PCB in mussels. In: SCCWRP Annual Report 1995. D.R. Young, I.S. Szpila.
- SCCWRP. 1976. Uptake and effects of chromium on marine fish. In: SCCWRP Annual Report 1996. M.J. Sherwood, J.L. Wright.
- SCCWRP. 1977. Metals in seafoods near outfalls. In: SCCWRP Annual Report 1977. T.K. Jan, M.D. Moore, D. Young.
- SWRCB. 1978. Trace elements in seafood organisms around southern California municipal wastewater outfalls. Publication No. 60. D.R. Young, M.D. Moore.
- SCCWRP. 1979. 60-Meter Control Survey Off Southern California. J.Q. Word. and A.J. Mearns. TM 229.
- SCCWRP. 1982. Seasonal changes in mussels. In: SCCWRP Biennial Report 1981-82. D.A. Brown, E.M. Perkins, K.D. Jenkins, P.S. Oshida, S.M. Bay, J.F. Alfafara, V. Raco.
- SCCWRP. 1987. 1985 Reference Site Survey. B.E. Thompson, J.D. Laughlin, and D.T. Tsukada. TR 221.



- SCCWRP. 1998a. Southern California bight 1994 pilot project: III. Sediment chemistry. K.C. Schiff, R.W. Gossett.
- SCCWRP. 1998b. Southern California bight 1994 pilot project: IV. Benthic infauna. M. Bergen, S.B. Weisberg, D. Cadien, A. Dalkey, D. Montagne, R.W. Smith, J.K. Stull, R.G. Velarde.
- SCCWRP. 1998c. Southern California bight 1994 pilot project: V. Demersal fish and megabenthic invertebrates. M.J. Allen, S.L. Moore, K.C. Schiff, S.B. Weisberg, D. Diener, J.K. Stull, A. Groce, J. Mubarak, C.L. Tang, R. Gartman.
- SCCWRP, 2003. Nobelt, J.A., E.Y. Zeng, R. Barid, R.W. Gossett, R.J. Ozretich, and C.R. Phillips. Southern California Bight 1998 Regional Monitoring Program: VI. Sediment Chemistry. Southern California Coastal Water Research Project, Westminster, CA.
- SCCWRP. 2008. The Southern California Bight Regional Monitoring Project Field Operations Manual.
- SCCWRP, 2006. Schiff, K. K. Maruya and K. Christensen. Southern California Bight 2003 Regional Monitoring Program: II. Sediment Chemistry. Southern California Coastal Water Research Project. Westminster, CA.
- SCCWRP. 2011. Allen, M.J., D. Cadien, E. Miller, D.W. Diehl, K. Ritter, S.L. Moore, C. Cash, D.J. Pondella, V. Raco-Rands, C. Thomas, R. Gartman, W. Power, A.K. Latker, J. Williams, J.L. Armstrong, and K. Schiff. Southern California Bight 2008 Regional Monitoring Program: Volume IV. Demersal Fishes and Megabenthic Invertebrates. Southern California Coastal Water Research Project, Costa Mesa, CA.
- SCCWRP. 2012. Schiff, K., R. Gossett, K. Ritter, L. Tiefenthaler, N. Dodder, W. Lao, and K. Maruya. Southern California Bight 2008 Regional Monitoring Program: III. Sediment Chemistry. Southern California Coastal Water Research Project, Costa Mesa, CA.
- Schwartz, R.C. 1978. Techniques for sampling and analyzing the marine macrobenthos. EPA/600/3 78/030.
- Shannon, C.E. and W. Weaver. 1963. *The Mathematical Theory of Communication*. Univ. Ill. Press, Urbana. 117 p.
- Short, J.W., and P.M. Harris. 1996. Petroleum hydrocarbons in caged mussels deployed in Prince William Sound after the *Exxon Valdez* oil spill. *Am. Fisheries Soc. Symp.* 18:29-39.
- Simpson, E.H. 1949. Measurement of diversity. *Nature (London)*. 163:668.



- Smith, Robert, M. Bergen, S.B. Weisberg, D. Cadien, A. Dalkey, D. Montagne, J. Stull and R. Velarde. 2001. Benthic response index for assessing infaunal communities on the Southern California Mainland Shelf. *Ecological Applications* 11:1073-1087.
- Sokal, R.R., and F.J. Rohlf. 1981. *Biometry*. 2nd Ed. W.H. Freeman Co. San Francisco. 859 pp.
- Soule D.F. and M. Oguri. 1987. The marine environment of Marina del Rey, 1986. *Marine Studies of SanPedroBay, California*. Part 20C. Harbors Environmental Projects. Univ. So. Calif., Los Angeles. 360 p.
- Soule D.F. and M. Oguri. 1988. The marine environment of Marina del Rey, 1987. *Marine Studies of SanPedroBay, California*. Part 20D. Harbors Environmental Projects. Univ. So Calif., Los Angeles. 289 p.
- Soule D.F. and M. Oguri. 1996. The marine environment of Marina del Rey, 1996, *Marine Studies of San Pedro Bay, Part 23, Harbors Environmental Project, University of Southern California*, 377 p.
- Southern California Coastal Water Research Project. See SCCWRP.
- SWQCB. 1965. An Oceanographic and Biological Survey of the Southern California Mainland Shelf. State of California Resource Agency. Publ. No. 27.
- SWRCB. 1973. Water quality criteria. 2nd Ed. Pub. 3-A. Calif. SWRCB. J.E. McKee, H.W. Wolf.
- SWRCB. 1978. Trace elements in seafood organisms around southern California municipal wastewater outfalls. Publication No. 60. D.R. Young and M.D. Moore.
- SWRCB. 1988. California State Mussel Watch. Ten year data summary 1977-1988. State Water Resources Control Board, Division of Water Quality, Sacramento, CA.
- SWRCB. 2009. Water Quality Control Plan, Ocean Waters of California, California Ocean Plan. State of California, State Water Resources Control Board.
- Tegner M. J., P. K. Dayton, P. B. Edwards and K. L. Riser. 1995. Sea urchin cavitation of giant kelp (*Macrocystis pyrifera* C. Agardh) holdfasts and its effects on kelp mortality across a large California forest. *Journal of Experimental Marine Biology and Ecology* Volume 191, Issue 1, 22 August 1995, Pages 83-99.
- Tetra Tech. 1986. Quality Assurance and Quality Control (QA/QC) for 301(h) Monitoring Programs: Guidance on Field and Laboratory Methods. EPA Contract No. 68-01-6938. TC-3953-04.
- United States Environmental Protection Agency. See USEPA.
- USEPA. 1978a. Techniques for Sampling and Analyzing the Marine Macrobenthos. EPA 600/3-78/030.



USEPA. 1978b. Use of Small Otter Trawls in Coastal Biological Surveys. EPA/600/3-78/083.

USEPA. 1983. Methods for the Chemical Analysis of Water and Wastes. EPA/600/4-79/020.

USEPA. 1986. Quality Criteria for Water, 1986. EPA/440/5-86/001.

USEPA and USCOE. 1991. Evaluation of dredged material proposed for ocean disposal. EPA/503/8-91/001.

USEPA. 1993. Guidance for assessing chemical contaminant data for use in fish advisories. Vol. 1. Fish sampling and analysis. EPA/823/R-93/002.

Word, J.Q. 1980. Classification of benthic invertebrates into infaunal trophic index feeding groups. In SCCWRP biennial report 1979-80, pp 103-22.

Zar, J.H. 1996. *Biostatistical Analysis*. 3rd Ed. Prentice-Hall, Inc. Englewood Cliffs, NJ. 820 pp.



10.2. Water Quality Correlation Data






Figure 10-1. Correlations between CTD probes and analysis of discrete water samples measured using field probes.






Figure 10-1. (continued)






Figure 10-1. (continued)






Figure 10-1. (continued)






Figure 10-1. (continued)



10.3. Particle Size



Table 10-2. Particle sizes by channel sizes in phi and microns for each Goleta sediment station.

Sample ID	phi Size																											
	<-1	-0.5	0	0.5	1	1.5	2	2.5	3	3.5	4	4.5	5	5.5	6	6.5	7	7.5	8	8.5	9	9.5	10	10.5	11	11.5	>12	
	Microns																											
	>2000	1410	1000	710	500	354	250	177	125	88.4	62.5	44.2	31.3	22.1	15.6	11.1	7.8	5.5	3.9	2.8	1.95	1.38	0.98	0.69	0.49	0.35	<0.24	
	coarse sand	coarse sand	med sand	med sand	med sand	med sand	fine sand	very fine sand	course silt	course silt	course silt		fine silt	very fine silt	very fine silt													
B1	0.00	0.00	0.00	0.08	1.76	9.79	21.15	19.85	14.08	8.29	4.66	2.99	2.59	2.64	2.69	2.63	2.10	1.49	0.95	0.76	0.51	0.37	0.35	0.28	0.00	0.00	0.00	
B2	0.00	0.00	0.00	0.03	0.41	1.49	6.08	18.67	25.68	17.28	7.74	3.65	2.65	2.61	2.76	2.84	2.36	1.70	1.09	0.88	0.61	0.45	0.43	0.39	0.20	0.00	0.00	
B3	0.00	0.00	0.00	0.03	0.42	1.79	8.38	25.17	27.56	13.78	5.23	2.51	2.04	2.18	2.34	2.34	1.86	1.31	0.84	0.68	0.48	0.36	0.34	0.29	0.07	0.00	0.00	
B4	0.00	0.00	0.00	0.03	0.51	2.59	13.05	32.03	23.63	8.48	3.17	1.84	1.75	1.99	2.19	2.25	1.88	1.38	0.91	0.73	0.50	0.36	0.35	0.31	0.07	0.00	0.00	
B5	0.00	0.00	0.00	0.04	0.61	2.59	9.56	20.24	20.90	13.25	6.83	3.81	2.98	2.98	3.17	3.30	2.82	2.11	1.37	1.09	0.72	0.49	0.48	0.45	0.22	0.00	0.00	
B6	0.00	0.00	0.00	0.04	0.77	3.67	12.95	23.71	22.74	12.75	5.47	2.66	2.07	2.15	2.32	2.35	1.89	1.34	0.85	0.69	0.49	0.38	0.35	0.29	0.07	0.00	0.00	

Table 10-3. Summary of particle sizes by fraction, percentiles, dispersion, sorting index and distribution.

Sample ID	Summary (Percent)					Percentile (microns)					Percentile (phi)					Microns			phi			Dispersion or Sorting Index	Distribution (phi)	
	Gravel*	Sand	Silt	Clay	Silt-Clay	5%	16%	50%	84%	95%	5%	16%	50%	84%	95%	Mean	Median	Mode	Mean	Median	Mode		Skewness	Kurtosis
B1	0.00	79.65	18.08	2.27	20.35	5.77	26.07	130.88	232.75	316.33	7.45	5.26	2.93	2.10	1.65	137.55	130.88	193.77	2.86	2.93	2.36	1.58	-0.05	-2.83
B2	0.00	77.37	19.66	2.97	22.63	4.71	21.15	91.25	152.50	209.92	7.74	5.56	3.45	2.71	2.24	94.25	91.25	106.11	3.40	3.45	3.23	1.43	-0.03	-2.92
B3	0.00	82.36	15.41	2.23	17.64	6.18	35.28	104.56	164.32	223.12	7.35	4.83	3.25	2.60	2.16	106.94	104.56	112.92	3.22	3.25	3.14	1.11	-0.03	-3.33
B4	0.00	83.50	14.18	2.33	16.50	5.91	40.21	121.77	177.88	237.95	7.41	4.64	3.03	2.48	2.06	120.21	121.77	140.44	3.05	3.03	2.83	1.08	0.02	-3.49
B5	0.00	74.01	22.54	3.44	25.99	4.02	15.24	94.35	167.49	234.53	7.97	6.04	3.40	2.57	2.08	99.95	94.35	111.39	3.32	3.40	3.16	1.73	-0.05	-2.70
B6	0.00	82.09	15.63	2.28	17.91	6.06	34.50	109.22	183.91	246.61	7.37	4.86	3.19	2.44	2.01	115.88	109.22	138.44	3.10	3.19	2.85	1.21	-0.07	-3.21

*Percentage of the sample retained on a 2 mm sieve.



10.4 Sediment Chemistry



Appendix

10-4. Sediment contaminant concentrations normalized to percent fine sediments in the Goleta survey area. Correlations by nonparametric Spearman's rho.

Constituent	Sediment Stations						Mean	S.D.	Correlations	
	B1	B2	B3	B4	B5	B6			Outfall	Point
Undifferentiated Organics										
Oil and Grease	40	31	25	28	9	10	23.8	12.2	0.29	-0.89
TKN	15	19	21	22	28	17	20.3	4.5	-0.93	0.43
TOC	123	150	11	242	196	73	133	83	-0.64	0.09
AVS	8.46	0.50	0.25	0.25	4.32	0.20	2.33	3.41	-0.16	-0.64
Heavy Metals										
Aluminum	397	388	444	515	406	377	421	52	-0.84	-0.09
Antimony	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.00	-0.58	-0.03
Arsenic	0.25	0.27	0.31	0.33	0.21	0.27	0.27	0.04	-0.12	0.14
Cadmium	0.02	0.02	0.03	0.02	0.02	0.02	0.02	0.00	0.00	-0.26
Chromium	1.42	1.29	1.47	1.66	1.29	1.29	1.40	0.15	-0.26	-0.03
Copper	0.20	0.26	0.24	0.31	0.26	0.20	0.25	0.04	-0.78	-0.14
Iron	413	482	518	585	457	461	486	59	-0.43	0.14
Lead	0.16	0.18	0.20	0.23	0.17	0.18	0.19	0.03	-0.32	0.37
Mercury	0.001	0.001	0.005	0.002	0.001	0.001	0.002	0.0017	-0.14	-0.14
Nickel	0.62	0.76	0.76	0.85	0.69	0.67	0.72	0.08	-0.70	0.09
Selenium	0.01	0.02	0.01	0.02	0.01	0.01	0.01	0.00	-0.14	0.26
Silver	0.002	0.003	0.002	0.004	0.003	0.002	0.003	0.001	-0.81	0.23
Tin	0.07	0.03	0.03	0.04	0.03	0.03	0.04	0.01	-0.17	-0.77
Zinc	1.02	1.24	1.27	1.44	1.17	1.12	1.21	0.14	-0.70	0.09
Complex Organics										
DDTs	0.25	0.10	0.07	0.10	0.12	0.06	0.12	0.07	-0.35	-0.49
HCH	ND	ND	ND	ND	ND	ND	ND	ND	0.00	0.00
Chlordane	ND	ND	ND	ND	ND	ND	ND	ND	0.00	0.00
Aldrin	ND	ND	ND	ND	ND	ND	ND	ND	0.00	0.00
Dieldrin	ND	ND	ND	ND	ND	ND	ND	ND	0.00	0.00
Heptachlor	ND	ND	ND	ND	ND	ND	ND	ND	0.00	0.00
Heptachlor epoxide	ND	ND	ND	ND	ND	ND	ND	ND	0.00	0.00
Mirex	ND	ND	ND	ND	ND	ND	ND	ND	0.00	0.00
Hexachlorobenzene	ND	ND	ND	ND	ND	ND	ND	ND	0.00	0.00
PCBs	ND	ND	ND	ND	ND	ND	ND	ND	0.00	0.00
Aroclors	ND	ND	ND	ND	ND	ND	ND	ND	0.00	0.00
PAHs	3.65	1.61	1.64	8.97	1.89	1.39	3.19	2.95	-0.55	-0.37
1-Methylnaphthalene	0.06	ND	ND	ND	ND	ND	0.01	0.02	0.40	-0.65
1-Methylphenanthrene	ND	0.05	0.07	0.11	0.07	ND	0.05	0.04	-0.97	0.29
2-Methylnaphthalene	0.07	0.05	0.06	ND	0.04	0.07	0.06	0.01	0.83	-0.21
2,3,5-Trimethylnaphthalene	ND	ND	ND	ND	ND	ND	ND	ND	0.00	0.00
2,6-Dimethylnaphthalene	0.06	0.06	0.06	ND	0.05	ND	0.04	0.03	0.00	-0.62
Acenaphthene	ND	ND	ND	0.07	ND	ND	0.01	0.03	-0.53	0.13
Biphenyl	ND	ND	ND	ND	ND	ND	ND	ND	0.00	0.00
Benz[a]anthracene	0.26	0.12	0.15	0.71	0.20	0.09	0.26	0.23	-0.66	-0.26
Benzo[b]fluoranthene	0.30	0.16	0.16	0.70	0.15	0.11	0.26	0.22	-0.55	-0.37
Benzo[e]pyrene	0.30	0.14	0.15	0.52	0.19	0.11	0.23	0.16	-0.55	-0.37
Benzo[g,h,i]perylene	ND	ND	ND	1.01	ND	ND	0.17	0.41	-0.53	0.13
Fluoranthene	0.78	0.31	0.31	1.35	0.31	0.27	0.56	0.43	-0.55	-0.37
Naphthalene	0.12	0.09	0.11	0.14	0.10	0.14	0.12	0.02	0.19	0.58
Perylene	3.54	1.34	0.99	1.17	1.29	0.76	1.51	1.02	-0.20	-0.60

Bold = marginally significant (0.05 < p < 0.10)

Bold = significant (p < 0.05)



Appendix

10-5 Sediment contaminant concentrations normalized to % total organic carbon (TOC) in the Goleta survey area. Correlations by nonparametric Spearman's rho.

Constituent	Sediment Stations						Mean	S.D.	Correlations	
	B1	B2	B3	B4	B5	B6			Outfall	Point
Undifferentiated Organics										
Oil and Grease	3257	2070	21813	1152	451	1417	5026.9	8277.9	0.46	-0.66
TKN	1232	1274	18850	913	1410	2277	4325.8	7130.1	0.23	0.37
AVS	688.92	33.12	220.00	10.28	219.94	27.92	200.03	258.18	0.17	-0.60
Heavy Metals										
Aluminum	32289	25803	391590	21263	20666	51962	90595	147909	0.64	-0.14
Antimony	0.60	0.41	5.50	0.32	0.34	0.87	1.34	2.05	0.64	-0.09
Arsenic	20.41	17.73	270.95	13.68	10.87	37.08	61.79	102.88	0.64	-0.14
Cadmium	1.63	1.65	23.57	0.88	0.77	2.78	5.21	9.02	0.58	-0.09
Chromium	115.50	85.69	1293.74	68.33	65.78	178.41	301.24	488.01	0.64	-0.14
Copper	16.43	17.49	214.61	12.71	13.23	27.13	50.27	80.68	0.58	-0.03
Iron	33633	32107	457100	24124	23294	63485	105624	172812	0.64	-0.14
Lead	12.83	11.79	175.45	9.56	8.87	24.65	40.52	66.35	0.64	-0.14
Mercury	0.093	0.087	4.720	0.065	0.052	0.159	0.863	1.8898	0.64	-0.14
Nickel	50.60	50.50	672.50	34.95	35.16	91.92	155.94	253.92	0.64	-0.09
Selenium	0.90	1.10	10.05	0.64	0.63	2.01	2.56	3.71	0.58	-0.09
Silver	0.160	0.176	2.000	0.150	0.137	0.308	0.489	0.743	0.58	-0.09
Tin	5.31	2.07	28.45	1.75	1.39	3.58	7.09	10.56	0.58	-0.43
Zinc	83.24	82.77	1122.30	59.53	59.57	154.69	260.35	423.71	0.64	-0.09
Complex Organics										
DDTs	20.40	2.20	1.20	1.70	3.20	1.10	4.97	7.60	0.58	-0.37
HCH	ND	ND	ND	ND	ND	ND	ND	ND	0.00	0.00
Chlordane	ND	ND	ND	ND	ND	ND	ND	ND	0.00	0.00
Aldrin	ND	ND	ND	ND	ND	ND	ND	ND	0.00	0.00
Dieldrin	ND	ND	ND	ND	ND	ND	ND	ND	0.00	0.00
Heptachlor	ND	ND	ND	ND	ND	ND	ND	ND	0.00	0.00
Heptachlor epoxide	ND	ND	ND	ND	ND	ND	ND	ND	0.00	0.00
Mirex	ND	ND	ND	ND	ND	ND	ND	ND	0.00	0.00
Hexachlorobenzene	ND	ND	ND	ND	ND	ND	ND	ND	0.00	0.00
PCBs	ND	ND	ND	ND	ND	ND	ND	ND	0.00	0.00
Aroclors	ND	ND	ND	ND	ND	ND	ND	ND	0.00	0.00
Total PAHs	297.20	36.50	29.00	148.10	49.00	24.90	97.45	108.14	-0.03	-0.26
1-Methylnaphthalene	5.20	ND	ND	ND	ND	ND	0.87	2.12	0.64	-0.14
1-Methylphenanthrene	ND	1.20	1.30	1.80	1.90	ND	1.03	0.85	0.14	0.31
2-Methylnaphthalene	6.00	1.10	1.10	ND	1.10	1.20	1.75	2.13	0.64	-0.14
2,3,5-Trimethylnaphthalene	ND	ND	ND	ND	ND	ND	ND	ND	0.00	0.00
2,6-Dimethylnaphthalene	4.80	1.30	1.10	ND	1.20	ND	1.40	1.77	0.64	-0.14
Acenaphthene	ND	ND	ND	1.10	ND	ND	0.18	0.45	0.64	-0.14
Biphenyl	ND	ND	ND	ND	ND	ND	ND	ND	0.00	0.00
Benz[a]anthracene	21.20	2.80	2.70	11.70	5.30	1.60	7.55	7.62	-0.174	-0.086
Benzo[b]fluoranthene	24.40	3.60	2.80	11.60	3.90	2.00	8.05	8.73	-0.029	-0.257
Benzo[e]pyrene	24.40	3.20	2.70	8.60	4.90	1.90	7.62	8.56	0.029	-0.257
Benzo[g,h,i]perylene	ND	ND	ND	16.70	ND	ND	2.78	6.82	0.029	0.029
Fluoranthene	63.60	7.10	5.40	22.30	8.10	4.90	18.57	22.99	0.174	-0.429
Naphthalene	9.60	2.10	1.90	2.30	2.50	2.50	3.48	3.01	0.638	-0.143
Perylene	288.40	30.30	17.40	19.30	33.40	13.60	67.07	108.70	0.580	-0.371

Bold = marginally significant (0.05 < p < 0.10)
Bold = significant (p < 0.05)



Appendix

10-6. Sediment chemistry minimum detection limits (MDL) and reporting limits (RL) and methods.

Parameter	MDL	RL	Units	Method	Parameter	MDL	RL	Units	Method
General Chemistry					Polynuclear Aromatic Hydrocarbons (Continued)				
Acid Volatile Sulfides	0.05	0.1	µg/g	Plumb, 1981 and TERL	Fluorene	1	5	ng/g	EPA 8270C
Oil & Grease	100	200	µg/g	SM 5520 E	Indeno[1,2,3-c,d]pyrene	1	5	ng/g	EPA 8270C
Percent Solids	0.1	0.1	%	SM 2540B	Naphthalene	1	5	ng/g	EPA 8270C
TKN	0.6	5	µg/g	EPA 351.3	Perylene	1	5	ng/g	EPA 8270C
Total Organic Carbon	100	200	µg/g	GC-01-111	Phenanthrene	1	5	ng/g	EPA 8270C
Trace Metals					Pyrene	1	5	ng/g	EPA 8270C
Aluminum	1	5	µg/g	EPA 6020	Polychlorinated Biphenyls (PCB's)				
Antimony	0.025	0.05	µg/g	EPA 6020	PCB003	1	5	ng/g	EPA 8270C
Arsenic	0.025	0.05	µg/g	EPA 6020	PCB008	1	5	ng/g	EPA 8270C
Cadmium	0.0025	0.005	µg/g	EPA 6020	PCB018	1	5	ng/g	EPA 8270C
Chromium	0.0025	0.005	µg/g	EPA 6020	PCB028	1	5	ng/g	EPA 8270C
Copper	0.0025	0.005	µg/g	EPA 6020	PCB031	1	5	ng/g	EPA 8270C
Iron	1	5	µg/g	EPA 6020	PCB033	1	5	ng/g	EPA 8270C
Lead	0.0025	0.005	µg/g	EPA 6020	PCB037	1	5	ng/g	EPA 8270C
Mercury	0.00001	0.00002	µg/g	EPA 245.7	PCB044	1	5	ng/g	EPA 8270C
Nickel	0.01	0.02	µg/g	EPA 6020	PCB049	1	5	ng/g	EPA 8270C
Selenium	0.025	0.05	µg/g	EPA 6020	PCB052	1	5	ng/g	EPA 8270C
Silver	0.01	0.02	µg/g	EPA 6020	PCB056(060)	1	5	ng/g	EPA 8270C
Tin	0.025	0.05	µg/g	EPA 6020	PCB066	1	5	ng/g	EPA 8270C
Zinc	0.025	0.05	µg/g	EPA 6020	PCB070	1	5	ng/g	EPA 8270C
Chlorinated Pesticides					PCB074	1	5	ng/g	EPA 8270C
2,4'-DDD	1	5	ng/g	EPA 8270C	PCB077	1	5	ng/g	EPA 8270C
2,4'-DDE	1	5	ng/g	EPA 8270C	PCB081	1	5	ng/g	EPA 8270C
2,4'-DDT	1	5	ng/g	EPA 8270C	PCB087	1	5	ng/g	EPA 8270C
4,4'-DDD	1	5	ng/g	EPA 8270C	PCB095	1	5	ng/g	EPA 8270C
4,4'-DDE	1	5	ng/g	EPA 8270C	PCB097	1	5	ng/g	EPA 8270C
4,4'-DDT	1	5	ng/g	EPA 8270C	PCB099	1	5	ng/g	EPA 8270C
Aldrin	1	5	ng/g	EPA 8270C	PCB101	1	5	ng/g	EPA 8270C
BHC-alpha	1	5	ng/g	EPA 8270C	PCB105	1	5	ng/g	EPA 8270C
BHC-beta	1	5	ng/g	EPA 8270C	PCB110	1	5	ng/g	EPA 8270C
BHC-delta	1	5	ng/g	EPA 8270C	PCB114	1	5	ng/g	EPA 8270C
BHC-gamma	1	5	ng/g	EPA 8270C	PCB118	1	5	ng/g	EPA 8270C
Chlordane-alpha	1	5	ng/g	EPA 8270C	PCB119	1	5	ng/g	EPA 8270C
Chlordane-gamma	1	5	ng/g	EPA 8270C	PCB123	1	5	ng/g	EPA 8270C
cis-Nonachlor	1	5	ng/g	EPA 8270C	PCB126	1	5	ng/g	EPA 8270C
Dieldrin	1	5	ng/g	EPA 8270C	PCB128	1	5	ng/g	EPA 8270C
Endosulfan sulfate	1	5	ng/g	EPA 8270C	PCB138	1	5	ng/g	EPA 8270C
Endosulfan-I	1	5	ng/g	EPA 8270C	PCB141	1	5	ng/g	EPA 8270C
Endosulfan-II	1	5	ng/g	EPA 8270C	PCB149	1	5	ng/g	EPA 8270C
Endrin	1	5	ng/g	EPA 8270C	PCB151	1	5	ng/g	EPA 8270C
Endrin aldehyde	1	5	ng/g	EPA 8270C	PCB153	1	5	ng/g	EPA 8270C
Endrin ketone	1	5	ng/g	EPA 8270C	PCB156	1	5	ng/g	EPA 8270C
Heptachlor	1	5	ng/g	EPA 8270C	PCB157	1	5	ng/g	EPA 8270C
Heptachlor epoxide	1	5	ng/g	EPA 8270C	PCB158	1	5	ng/g	EPA 8270C
Methoxychlor	1	5	ng/g	EPA 8270C	PCB167	1	5	ng/g	EPA 8270C
Mirex	1	5	ng/g	EPA 8270C	PCB168/132	1	5	ng/g	EPA 8270C
Oxychlordane	1	5	ng/g	EPA 8270C	PCB169	1	5	ng/g	EPA 8270C
Perthane	5	10	ng/g	EPA 8270C	PCB170	1	5	ng/g	EPA 8270C
trans-Nonachlor	1	5	ng/g	EPA 8270C	PCB174	1	5	ng/g	EPA 8270C
Polynuclear Aromatic Hydrocarbons (PAHs)					PCB177	1	5	ng/g	EPA 8270C
1-Methylnaphthalene	1	5	ng/g	EPA 8270C	PCB180	1	5	ng/g	EPA 8270C
1-Methylphenanthrene	1	5	ng/g	EPA 8270C	PCB183	1	5	ng/g	EPA 8270C
2,3,5-Trimethylnaphthalene	1	5	ng/g	EPA 8270C	PCB187	1	5	ng/g	EPA 8270C
2,6-Dimethylnaphthalene	1	5	ng/g	EPA 8270C	PCB189	1	5	ng/g	EPA 8270C
2-Methylnaphthalene	1	5	ng/g	EPA 8270C	PCB194	1	5	ng/g	EPA 8270C
Acenaphthene	1	5	ng/g	EPA 8270C	PCB195	1	5	ng/g	EPA 8270C
Acenaphthylene	1	5	ng/g	EPA 8270C	PCB199(200)	1	5	ng/g	EPA 8270C
Anthracene	1	5	ng/g	EPA 8270C	PCB201	1	5	ng/g	EPA 8270C
Benz[a]anthracene	1	5	ng/g	EPA 8270C	PCB206	1	5	ng/g	EPA 8270C
Benzo[a]pyrene	1	5	ng/g	EPA 8270C	PCB209	1	5	ng/g	EPA 8270C
Benzo[b]fluoranthene	1	5	ng/g	EPA 8270C	Polychlorinated Biphenyls (PCB's)				
Benzo[e]pyrene	1	5	ng/g	EPA 8270C	Aroclor 1016	1	10	ng/g	EPA 8270C
Benzo[g,h,i]perylene	1	5	ng/g	EPA 8270C	Aroclor 1221	1	10	ng/g	EPA 8270C
Benzo[k]fluoranthene	1	5	ng/g	EPA 8270C	Aroclor 1232	1	10	ng/g	EPA 8270C
Biphenyl	1	5	ng/g	EPA 8270C	Aroclor 1242	1	10	ng/g	EPA 8270C
Chrysene	1	5	ng/g	EPA 8270C	Aroclor 1248	1	10	ng/g	EPA 8270C
Dibenz[a,h]anthracene	1	5	ng/g	EPA 8270C	Aroclor 1254	1	10	ng/g	EPA 8270C
Dibenzothiophene	1	5	ng/g	EPA 8270C	Aroclor 1260	1	10	ng/g	EPA 8270C
Fluoranthene	1	5	ng/g	EPA 8270C					



10-7. Sediment chemistry complex organic derivatives.

Sediment Stations	B1	B2	B3	B4	B5	B6
DDTs (ng/g)						
2,4'-DDD	0.0	0.0	0.0	0.0	0.0	0.0
2,4'-DDE	0.0	0.0	0.0	0.0	0.0	0.0
2,4'-DDT	0.0	0.0	0.0	0.0	0.0	0.0
4,4'-DDD	0.0	0.0	0.0	0.0	0.0	0.0
4,4'-DDE	5.1	2.2	1.2	1.7	3.2	1.1
<u>4,4'-DDT</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>
Sum =	5.1	2.2	1.2	1.7	3.2	1.1
Chlordane (ng/g)						
Chlordane-alpha	0.0	0.0	0.0	0.0	0.0	0.0
Chlordane-gamma	0.0	0.0	0.0	0.0	0.0	0.0
cis-Nonachlor	0.0	0.0	0.0	0.0	0.0	0.0
trans-Nonachlor	0.0	0.0	0.0	0.0	0.0	0.0
<u>None</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>
Sum =	0.0	0.0	0.0	0.0	0.0	0.0
HCH (ng/g)						
HCH-alpha	0.0	0.0	0.0	0.0	0.0	0.0
HCH-beta	0.0	0.0	0.0	0.0	0.0	0.0
HCH-delta	0.0	0.0	0.0	0.0	0.0	0.0
<u>HCH-gamma</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>
Sum =	0.0	0.0	0.0	0.0	0.0	0.0
Polychlorinated Biphenyls (PCB's, ng/g)						
PCB003	0.0	0.0	0.0	0.0	0.0	0.0
PCB008	0.0	0.0	0.0	0.0	0.0	0.0
PCB018	0.0	0.0	0.0	0.0	0.0	0.0
PCB028	0.0	0.0	0.0	0.0	0.0	0.0
PCB031	0.0	0.0	0.0	0.0	0.0	0.0
PCB033	0.0	0.0	0.0	0.0	0.0	0.0
PCB037	0.0	0.0	0.0	0.0	0.0	0.0
PCB044	0.0	0.0	0.0	0.0	0.0	0.0
PCB049	0.0	0.0	0.0	0.0	0.0	0.0
PCB052	0.0	0.0	0.0	0.0	0.0	0.0
PCB056(060)	0.0	0.0	0.0	0.0	0.0	0.0
PCB066	0.0	0.0	0.0	0.0	0.0	0.0
PCB070	0.0	0.0	0.0	0.0	0.0	0.0
PCB074	0.0	0.0	0.0	0.0	0.0	0.0
PCB077	0.0	0.0	0.0	0.0	0.0	0.0
PCB081	0.0	0.0	0.0	0.0	0.0	0.0
PCB087	0.0	0.0	0.0	0.0	0.0	0.0
PCB095	0.0	0.0	0.0	0.0	0.0	0.0
PCB097	0.0	0.0	0.0	0.0	0.0	0.0
PCB099	0.0	0.0	0.0	0.0	0.0	0.0
PCB101	0.0	0.0	0.0	0.0	0.0	0.0
PCB105	0.0	0.0	0.0	0.0	0.0	0.0
PCB110	0.0	0.0	0.0	0.0	0.0	0.0
PCB114	0.0	0.0	0.0	0.0	0.0	0.0
PCB118	0.0	0.0	0.0	0.0	0.0	0.0
PCB119	0.0	0.0	0.0	0.0	0.0	0.0
PCB123	0.0	0.0	0.0	0.0	0.0	0.0
PCB126	0.0	0.0	0.0	0.0	0.0	0.0
Polychlorinated Biphenyls (PCB's, ng/g)						
PCB128	0.0	0.0	0.0	0.0	0.0	0.0
PCB138	0.0	0.0	0.0	0.0	0.0	0.0
PCB141	0.0	0.0	0.0	0.0	0.0	0.0
PCB149	0.0	0.0	0.0	0.0	0.0	0.0
PCB151	0.0	0.0	0.0	0.0	0.0	0.0
PCB153	0.0	0.0	0.0	0.0	0.0	0.0
PCB156	0.0	0.0	0.0	0.0	0.0	0.0
PCB157	0.0	0.0	0.0	0.0	0.0	0.0
PCB158	0.0	0.0	0.0	0.0	0.0	0.0
PCB167	0.0	0.0	0.0	0.0	0.0	0.0
PCB168/132	0.0	0.0	0.0	0.0	0.0	0.0
PCB169	0.0	0.0	0.0	0.0	0.0	0.0
PCB170	0.0	0.0	0.0	0.0	0.0	0.0
PCB174	0.0	0.0	0.0	0.0	0.0	0.0
PCB177	0.0	0.0	0.0	0.0	0.0	0.0
PCB180	0.0	0.0	0.0	0.0	0.0	0.0
PCB183	0.0	0.0	0.0	0.0	0.0	0.0
PCB187	0.0	0.0	0.0	0.0	0.0	0.0
PCB189	0.0	0.0	0.0	0.0	0.0	0.0
PCB194	0.0	0.0	0.0	0.0	0.0	0.0
PCB195	0.0	0.0	0.0	0.0	0.0	0.0
PCB199(200)	0.0	0.0	0.0	0.0	0.0	0.0
PCB201	0.0	0.0	0.0	0.0	0.0	0.0
PCB206	0.0	0.0	0.0	0.0	0.0	0.0
<u>PCB209</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>
Sum =	0.0	0.0	0.0	0.0	0.2	0.0
Aroclors						
Aroclor 1016	0.0	0.0	0.0	0.0	0.0	0.0
Aroclor 1221	0.0	0.0	0.0	0.0	0.0	0.0
Aroclor 1232	0.0	0.0	0.0	0.0	0.0	0.0
Aroclor 1242	0.0	0.0	0.0	0.0	0.0	0.0
Aroclor 1248	0.0	0.0	0.0	0.0	0.0	0.0
Aroclor 1254	0.0	0.0	0.0	0.0	0.0	0.0
<u>Aroclor 1260</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>
Sum =	0.0	0.0	0.0	0.0	0.0	0.0
Polynuclear Aromatic Hydrocarbons (PAH's, ng/g)						
Acenaphthylene	0.0	0.0	0.0	0.0	0.0	0.0
Anthracene	4.4	1.1	0.0	2.7	1.6	1.4
Benz[a]anthracene	5.3	2.8	2.7	11.7	5.3	1.6
Benzo[a]pyrene	3.8	2.5	3.1	11.1	4.2	1.5
Benzo[b]fluoranthene	6.1	3.6	2.8	11.6	3.9	2.0
Benzo[g,h,i]perylene	0.0	0.0	0.0	16.7	0.0	0.0
Benzo[k]fluoranthene	4.1	2.8	3.0	12.5	3.9	1.3
Chrysene	9.7	4.7	3.9	14.6	8.4	2.3
Dibenz[a,h]anthracene	0.0	0.0	0.0	6.1	0.0	0.0
Fluorene	2.0	1.5	0.0	2.1	1.6	1.2
Indeno[1,2,3-c,d]pyrene	0.0	0.0	0.0	19.0	0.0	0.0
Phenanthrene	20.9	11.4	8.8	20.1	10.2	9.7
<u>Pyrene</u>	<u>18.0</u>	<u>6.1</u>	<u>4.7</u>	<u>19.9</u>	<u>9.9</u>	<u>3.9</u>
Sum =	74.3	36.5	29.0	148.1	49.0	24.9



10.6. Benthic Infauna



10-8. Benthic infauna taxonomic abundances.

Phylum	Class	Species	Station & Replicate																													
			B1					B2					B3					B4					B5					B6				
			1	2	3	4	5	1	2	3	4	5	1	2	3	4	5	1	2	3	4	5	1	2	3	4	5	1	2	3	4	5
Annelida	Oligochaeta	Oligochaeta	21	19	20	17	29	12	9	7	51	12	11	1	3	5	6	1	18	8	2	9	34	30	112	7	12	2	1			
	Polychaeta	Amaeana occidentalis	1								1														1	2		1				
		Ampharete finmarchica																									1	2				
	Total Annelids 12,789	Ampharete labrops	1	3	2		2	2		1	4	2	1	3			2	1							1							
	% of Population 41.58	Amphiteis scaphobranchiata	2	4	1	1			1	1	4	1	1			1													2	1		
		Ancistrosyllis hamata																													1	
		Anobothrus gracilis	2	1																												
		Anotomastus gordiodes	1																													
		Aphelochaeta elongata	11	12	6	6	5	4	5	4	3	3	8	3	4	2		2	8	3		1	5	2	2							
		Aphelochaeta glandaria Cmplx	6	7		1	2	3	1	4	10	17	1	6	1	1						1	1	2	4	3	4	3				
		Aphelochaeta petersenae		1			1		1	1	2			1						1	1				2	2						
		Aphelochaeta sp	1	5					1	1	3	1											1	1	5	1	2					
		Aphelochaeta sp HYP2						1					1																			
		Apopriopio pygmaea	6		1	2	2		1	2			3		2	1	1		1	3	3		6	3			1	2	1	1	1	1
		Arabella iricolor																1	5			1	7	7	4	2						
		Arabella pectinata																					2	1								
		Arcteoobia cf anticostiensis																										1		1		
		Arctonoe pulchra																					1		1							
		Aricidea (Acmira) catherinae	1	2			1			1	1		1	1			1					2	2		1	2						1
		Aricidea (Acmira) horikoshii		5	1	1	4	3			1		2				1	1		1	1		1	1								1
		Aricidea (Acmira) simplex	1					1																			1					
		Aricidea (Allia) hartleyi																														
		Aricidea (Aricidea) wassi		1														1	2		1							1		1		1
		Armandia brevis			2	2	4					1					1		9	2	2	3	3	9	6							
		Artacameila hancocki																						1								
		Asabellides lineata			1												1						4		1							
		Bipalponophytys cornuta	1		1			2	1	2	2	1	1	1	1		1			1	3		2		3	1	1	1				
		Brada pilosa																	1													
		Brania californiensis		1										1		1	1		1				4	1	2	5	1					
		Brania mediodentata					2				1						1															
		Capitella capitata Cmplx				3			1		4				2	2		4	15	4	5	4	5	3	47	26	24	1				
		Capitellidae	1																													
		Carazziella sp A																		3					1							
		Caulierella pacifica																								1						
		Chaetopterus variopedatus Cmplx	1																	1	1	1		1								
		Chaetozone acuta	1			1						1											2	1	1							
		Chaetozone columbiana										1				2		1	3		2			2	5	1						
		Chaetozone hedgpethi													2	1									3	1						
		Chaetozone sp																							1							
		Cossura sp A	37	20	19	18	30	55	47	23	68	19	45	36	21	34	18	9	13	8	11		64	59	92	36	26	50	67	12	6	5
		Dialychone albocincta										2											2	1	8	3						
		Dialychone trilineata																	1		1											
		Dialychone veleronis	2	1		1				1	2	1		1				2		1		1										
		Diopatra ornata	1	3	10	4	4			2	1		3	1		4	6	4	8	5	2	4	8	17	13	4	14	1	1			1
		Diopatra sp									1																					
		Diopatra tridentata		1	2																							1		1		
		Dipolydora socialis		1			1	2	5	6	8	1	8	4	2	4	4		1								1	4	5	4		1
		Drilonereis mexicana			1								1		1																	
		Drilonereis sp							1										1						1							
		Epigamia-Myrianida Cmplx				1																1										
		Eteone dilatata	1			1			1		1						1								4		1					
		Euclymeninae	2	2							1	1								2			6	7	11	5	4			1		



Appendix

10-8. Continued.

Phylum	Class	Species	Station & Replicate																														
			B1					B2					B3					B4					B5					B6					
			1	2	3	4	5	1	2	3	4	5	1	2	3	4	5	1	2	3	4	5	1	2	3	4	5	1	2	3	4	5	
		Euclymeninae sp A	1	1	2			1	2		4	5	1		3	2		5	4	8	1		23	20	35	16	26	5	7	3	2	2	
		Eulalia levicornuta Cmplx																					1										
		Eulalia quadrioculata																					1	1									
		Eumida longicornuta	3	2	3			1		1			1	1						3	2									1			
		Eupolyornia heterobranchia	2							3															3					1			
		Eusyllinae																															
		Eusyllis sp		1		4									2			3	9	2	2		7		3	6	1	2	2	1		2	
		Exogone dwisula									1				1													1					
		Exogone lourei	8	1							1											1		1	4								
		Glycera americana	1				1							1					2	1	1		2	2		1					1		
		Glycera macrobranchia		2											1				1			2						1					
		Glycera nana	6	8	7	4	8	2	3		1		2	7	3		1	3	12	3	1	1	2	4	2	2		1		1	1	1	
		Glycine armigera	8	1	2	3	1	1	2	1	2	1	2	3	2	3	3	3	3	3	1	1	3	6	3	4	3	4	2	3	5	3	
		Goniada maculata			2						1			1																2	2		
		Halosydna johnsoni		1			1	1			2				1	3		5			1		4	1	3	5	2						
		Harmothoe hirsuta																					3	3	1		2						
		Hesionidae																															
		Hyalopomatus biformis		1											2										1	1							
		Lanice conchilega		1														1	1									1					
		Laonice cirrata	1	1		1	1					1													1		1					1	
		Leitoscoloplos pugettensis	1	4			2		2		2			1		1		2	2	3	3			1	5	2				3	2	2	
		Levinsenia gracilis	32	106	17	48	94	11	17	7	23	12	58	19	16	15	8	4	12	5	5	5	119	134	217	51	74			2		1	
		Loimia sp A	1				1												1			1			1			1					
		Lumbrineris cruzensis	3	6	1	6	9		1				1	1				1	5	1	5	1	7	4	2	2	2			1		1	
		Lumbrineris japonica	3	1	4		2	1	1	1		1	1	1		2		1		1		1			2		1	1					
		Lumbrineris ligulata	3		2	1	1		2									2	4	1	1		11	18	8	6	4						
		Lumbrineris limicola		1																													
		Lumbrineris sp	4	3	1							1	1											2	4	3	1	1					
		Lysippe sp A	1	4				1															9	3	7		4	2	1	1			
		Magelona berkeleyi	11	5	9	5	4	11	33	8	18	4	22	15	13	19	20	23	27	34	5	22	8	5	6	14	3	25	3	9	3	5	
		Magelona sacculata																4	5	1												1	
		Magelona sp	2		1			1	4		2	1	2	1	1	3	2						1										
		Malmgreniella macginitiei		1						1	2	4	1	1		2	1	5						1	2						1		
		Malmgreniella scriptoria			1																												
		Malmgreniella sp					1	1	1		2						1					1		1	4		1	2	1	1			
		Malmgreniella sp A	4	3	2		2	1	1				3				1						2		4	1	1						
		Marphysa disjuncta				1																											
		Mediomastus sp	56	82	29	73	44	29	36	39	138	38	72	56	35	45	57	37	95	24	34	58	341	201	415	116	123	82	24	44	6	8	
		Megalomma pigmentum									1														1								
		Melinna oculata		1		1	1				1	2	1														1	1			1	1	
		Metasychis disparidentatus	1	1	1		1	3	1	2	3	7	4	3	1	2	1		1	1			2		1		1	2	1	1	2		
		Micropodarke dubia		1								1																					
		Monticellina cryptica	1	1	1	1	1	1		2	1		2	3		1	3	3		1	1		2	11	5	3	7	8	1	1			
		Monticellina serratiseta		1																													
		Monticellina siblina	1	25	4	8	4	1		1	43	15		9		1	5	21	2	1	1			1	17	2	1	2	1		1		
		Mooreonuphis nebulosa	1	1	1			1	1	1	1	1	2		1							1	2	1	2			4	2		14	1	
		Neosabellaria cementarium			1							1				1		1	5	8	1	3	14	6	6	2	3						
		Nephtys caecoides	1	1	3		1	8	7	9	9	9	6	6	6	4	4	3	1		1	2	1	1	6	5	1	6	7	7	5	4	
		Nephtys ferruginea	3	1		1		9	6	6	3	3	1	4	1	5	5		1	1	10	1	4	1	6		3	3	1			3	
		Nereiphylla sp SD1			1															1													
		Nereis latescens																2		2						1							
		Nereis sp A	2	5	1	2	1	1			3	1	2		2			1	5		1	5	12	12	12	2	17			1			



Appendix

10-8. Continued.

Phylum	Class	Species	Station & Replicate																														
			B1					B2					B3					B4					B5					B6					
			1	2	3	4	5	1	2	3	4	5	1	2	3	4	5	1	2	3	4	5	1	2	3	4	5	1	2	3	4	5	
		<i>Sigambra setosa</i>		2			2																									1	
		<i>Sphaerosyllis californiensis</i>							1	2							1		1	1				1	1	1	1					1	
		<i>Spiochaetopterus costarum</i> Cmplx	18	7	26	29	30	46	8	20	48	21	21	30	9	24	15	13	35	33	30	31	45	36	53	33	17	31	22	16	26	6	
		Spionidae						1	1	4	1						1	1	1	1	2		3	1	2	1	1					1	
		<i>Spiophanes berkeleyorum</i>	5	2	3			3	1	1	4	1	3	7	1	7	6		8	4	5	2	10	3	7	4	5	6	4	3	3		
		<i>Spiophanes duplex</i>	23	38	37	22	14	20	14	38	42	23	35	56	16	22	60	7	9	5	6	10	28	21	51	26	32	22	6	8	3	8	
		<i>Spiophanes norrisi</i>	3	4	4	5	3	5	2	6	14	7	12	9	5	7	6	7	6	5	4	2	7	10	5	7		5	2	3	1	1	
		<i>Spiophanes</i> sp	4	1	3	5	2						1																				
		<i>Sternaspis affinis</i>	2	4	2		2		1	2	3	3	4	3	3	1	3					1	1		1	3	2		7	2		1	
		<i>Sthenelais berkeleyi</i>																														1	
		<i>Sthenelais tertialabra</i>	5		1		1	12	4	9	7	5	3	2	5	8	10	2	1		3	1	1	1	5		3	5	8	4	2	8	
		<i>Sthenelanella uniformis</i>		4	3	1		3	1	1	3	6	4			1		1						1	1			1	3				
		<i>Streblosoma crassibranchia</i>						2	1	1				2	2											1			1	1			
		<i>Streblosoma</i> sp																						1								1	
		<i>Streblosoma</i> sp B																							2								
		<i>Tenonia priops</i>					1	1	2		3	1	2	1		1	1	1	3		5	1		4	2	1		1	1		2		
		<i>Terebellides californica</i>		1				1		1		1	1	1					1	1	1	1	1	1					2				
		<i>Terebellides reishi</i>					1				1															1							
		<i>Travisia</i> sp					1																										
		<i>Typosyllis farallonensis</i>																										3					
		<i>Typosyllis heterochaeta</i>	1	2						2	1		1			1															1		
		<i>Typosyllis</i> sp																2		1													
Arthropoda	Cirripedia	<i>Hamatoscalpellum californicum</i>																					1								5		
		<i>Megabalanus californicus</i>		2			12				1						1							1					1	1		1	
	Malacostraca	<i>Acidostoma hancocki</i>		1													1																
		<i>Alpheus bellimanus</i>																	1					2		1			4	2	2		
		<i>Americhelidium shoemakeri</i>	4	1	1	3	6	2		2		1	1	1	2	1	4	6	14	8	7	11	5	2	4	6	4	4	2	2			
		<i>Ampelisca agassizi</i>	3	2	1	1	1					9	1	2	1	1	1				1	1		1				3	7	3		3	
		<i>Ampelisca brachycladus</i>					1				8					3	6											1					
		<i>Ampelisca brevisimulata</i>	17	14	19	17	2	17	11	6	14	3	10	12	10	9	3	3		2								13	15	12	14	6	
		<i>Ampelisca careyi</i>															1																
		<i>Ampelisca cf. brevisimulata</i>																										1					
		<i>Ampelisca cristata cristata</i>			3	3	2	8	3	6	2	17	5	11	1	6	7	26	14	7	9	3	6	6	19	11		1	2	3	4		
		<i>Ampelisca cristata microdentata</i>	1	3	1	1																											
		<i>Ampelisca milleri</i>	1	1	1	1	41			1	2		1		1	2		1	1				5	2	1						1	2	
		<i>Ampelisca pugetica</i>	1	1	3			1		1	5	1	1	2	2	1			1									5	4	1	1		
		<i>Ampelisca</i> sp						1					1																				
		<i>Ampelisciphotis podophthalma</i>	5	12		1		1		6	3	4	2	5		1	2							2				13	16	15	17	3	
		<i>Amphideutopus oculatus</i>	3	13	4	10		7	1	2	1	3	2	10	2	1										2		19	21	8	5	5	
		<i>Ampithoe plumulosa</i>																						3	1		1						
		<i>Ampithoe simulans</i>																	2	1	2												
		<i>Ampithoe valida</i>																													1		
		Anthuridae							1																								
		Aoridae																										18	20	16	2	2	
		<i>Aoroides exilis</i>												2		1		17		2	1		8	11	16	3	5						
		<i>Aoroides inermis</i>																			2	3											
		<i>Aoroides intermedia</i>	6	8		21						2				7		4	95	14	32	31	30	37	49	19	22					4	
		<i>Aoroides</i> sp					5				1	1				1							30	41	27	5	11						
		<i>Aoroides</i> sp A																					12	2	3	4	3	3					
		<i>Aoroides spinosa</i>								7	1				1	4		20	4	5	3		4	3	1	3		1					
		<i>Apolochus picadurus</i>					15																1	1									
		Arcturidae					21				2					2		6	5	7	2		4	3	4	5	3						



Appendix

10-8. Continued.

Phylum	Class	Species	Station & Replicate																														
			B1					B2					B3					B4					B5					B6					
			1	2	3	4	5	1	2	3	4	5	1	2	3	4	5	1	2	3	4	5	1	2	3	4	5	1	2	3	4	5	
		<i>Argissa hamatipes</i>	3	2	1		2	3	2	6	1	1	3	4	3	2	4	4				1					1	1	2	2	2	2	
		<i>Aruga oculata</i>	1				1						1	1				2	1	1							2	2		1		1	
		<i>Bemlos concavus</i>																					22	9	8	1	5						
		<i>Byblis millsii</i>			1	1	1	1					1	1				1										1	1				
		<i>Caecognathia crenulatifrons</i>	2	2	4	2	1	2	2	6	6	21	7	9	7	3	1	1	1									8	5	16		6	
		<i>Califanthura squamosissima</i>															1																
		<i>Campylaspis canaliculata</i>											1																				
		<i>Campylaspis hartae</i>			1																						1						
		<i>Campylaspis rubromaculata</i>		2		2		1					2					1										1					
		<i>Cancer productus</i>	1				1											1	1			1	1	1	2	1							
		<i>Cancridae</i>	1		1																		1										
		<i>Caprella californica</i>			1		9										6	28	37	39	16		185	14	140	81	97					1	
		<i>Caprella gracilior</i>					1																				4						
		<i>Caprella sp</i>																6	3	4	3		17	3	15	14	16						
		<i>Caprellidae</i>																2	15	1	13		15	6	11	21	32						
		<i>Chevalia inaequalis</i>										1				2																	
		<i>Columbaora cyclocoxa</i>			1		38																43	45	7	28	46	41	52	2	8		
		<i>Crangon nigricauda</i>																				1											
		<i>Cyclaspis nubila</i>										1																					
		<i>Deutella californica</i>					12																				8						
		<i>Diastylis californica</i>	1	4	3		1	1	2	3			4	3	1		1	1	1	1			1	1	1			2	3	1	2		
		<i>Discorsopagurus schmitti</i>					1																										
		<i>Edotia sp B</i>	11	6	7	7	3				5	2	2	1	1	4		2					1						1				
		<i>Erichthonius brasiliensis</i>			1							6					3		2	1	2		1		8		1						
		<i>Eualus lineatus</i>		2		1	71				2	1			2			18	22	17	4		17	27	2	6	10					1	
		<i>Foxiphalus golfensis</i>	34	14	20	26	8	21	7	22	6	3	4	12	6	11	6	2	13	17	10	7	2	6	6	19	5	8	8	5	9	2	
		<i>Foxiphalus obtusidens</i>	19	24	26	12	6	4	5	2	2	5	10	22	12	6	5	20	26	18	8	17	11	4	3	7	8	4	7	12	10	9	
		<i>Foxiphalus sp</i>	1		1						1			2				1	2			1					1						
		<i>Gammaropsis shoemakeri</i>																					8	8									
		<i>Gammaropsis sp</i>					23					2					3	11	16	5	12		3	2	1								
		<i>Gammaropsis thompsoni</i>			3		248					13	1		1		2	23	33	31	23	30		28	41	15	21	23					1
		<i>Haigia diegensis</i>					3																										
		<i>Haliophasma geminatum</i>	3	2	4	2	1	5	6	2		1	3	9	1	7	1	2		1	3	1		2	4		3	2	5	3	2	2	
		<i>Hartmanodes hartmanae</i>	2	6	3	5		1	1	1		1	6	6	2	3		5	1													1	
		<i>Hemilamprops californicus</i>	1										1	1			1														1	3	
		<i>Heterophoxus oculatus</i>	3		3	3	3	5	3	1	2	3	1	2	1	1		1	3	6	5		4	8	8	3	8	2	1				
		<i>Hippomedon zetesimus</i>	1	1					1	2		1							1	6					1			2					
		<i>Idarcturus allelomorphus</i>	3	5		109		2		3	6	3		2			7	46	52	50	26		276	49	98	52	135	8	1	1	2	10	
		<i>Idarcturus sp</i>	1	3		23			3		2						4		13	27	14	13	67	21	27	21	33						
		<i>Ischyroceridae</i>																					3	4	4	1	3						
		<i>Ischyrocerus pelagops</i>					4												47	2	6	1	2	7	4	2	4						
		<i>Isocheles pilosus</i>																									1						
		<i>Latulambus occidentalis</i>					2					1	1	1	1		2	1	1		1	1	1	1	1	1	2						
		<i>Lepidepecreum serraculum</i>										1	1	1									1	1									
		<i>Leptocheilia dubia Cmplx</i>						28	41	16	51	14	11	53	7	9	31	6	62	22	18	60	27	31	22	29	13	10	19	20	9	9	
		<i>Leptostyliis sp</i>																									1						
		<i>Leucon subnasica</i>											1																				
		<i>Listriella diffusa</i>	8	8				3		2	4		3	1	1		1		2						1		1	2	2	1			
		<i>Listriella goleta</i>				2		4	2	2	4	3		5		1	2		1			2			1			6	4	2			
		<i>Listriella melanica</i>			2												1		1		1		2		2	2	1						
		<i>Lophopanopeus bellus</i>					13												5	2	1	1	27	18	14	3	12						
		<i>Lophopanopeus frontalis</i>																				1	1	2	4	1	2						



Appendix

10-8. Continued.

Phylum	Class	Species	Station & Replicate																													
			B1					B2					B3					B4					B5					B6				
			1	2	3	4	5	1	2	3	4	5	1	2	3	4	5	1	2	3	4	5	1	2	3	4	5	1	2	3	4	5
		Euphlymedes carcharodonta	120	53	87	52	50	183	167	193	126	149	70	155	86	102	102	176	94	147	88	67	5	4	12	12	7	82	93	89	70	99
		Eusarsiella thominx	3		1																		1				1					
		Leuroleberis sharpei																					1				1					
		Postasterope barnesi	4	4	3		2	2					2			1		1	5		1		3	1	4		2					
		Rutiderma rostratum		2	1	1					1							1	1	1	1	1	1	1	1	2						
		Xenoleberis californica	1	6	4	2	4	7		1	1		1	1		3	1	1	4	4	3	3				2		3	2	5	1	11
	Pycnogonida	Anoplodactylus pacificus																1					3	1	1	1						
Echinodermata	Aphiuroidea	Amphioplus sp A																														1
	Asteroidea	Astropecten californicus		7	5	4				1			2	1	1	2				1						1		1	3			
	Echinoidea	Echinoidea																					1									
		Strongylocentrotus purpuratus																						1								
	Holothuroidea	Chiridota sp																1					1									
		Leptosynapta sp	18	16	6	5	11	6	4	5	2	3	8	1		1	4						4	1	4		1	2	1	1	1	5
		Pentamera lissoplaca																													1	
		Pentamera pseudopoplifera			1																											
	Ophiuroidea	Amphiodia digitata	1		3		1		1		3		2	2		3		2	2	1	3		1		1	1			3	1	2	
		Amphiodia psara									1															1						
		Amphiodia sp	2	1	1	1		11	5	5	5	2	6	9	1	9	6	6	10	4	6	10	3	2	3	5	2	11	4	10	2	2
		Amphiodia urtica	1	4	1		3	2	3	5	4	1	3	3	1	7	2	6	6	5			2	1	7	3	5	2	1			
		Amphioplus sp								1																						
		Amphipholis sp	1																													
		Amphipholis squamata			3			5		2	5	2		1	2	1	2		1	2			2	1	7	2	2	3				1
		Amphiuroidae	3	6	2	1		4	6	6	5	1	5	2	1	3	3	4	1	4	1		1		1	4	2	4		1	2	1
		Ophiothrix spiculata					1									1				1												
		Ophiura luetkenii																					1									
		Ophiuroconis bispinosa							1					2	1			1		1												1
Mollusca	Bivalvia	Axinopsida serricata									1																	1	1	1		
		Compsomyx subdiaphana	1	3				2	1	2	1	2	2	1															2	1		
		Cooperella subdiaphana								1		1	1	2	1	2				1											1	
		Ensis myrae	1					2	1	1		1	2	2									1						2			
		Gari fucata						3	1	3	6	1	2	1		1		1		1						2		2		1		1
		Kurtiella compressa									1													1	1	1						
		Kurtiella grippi		1			3	1										1							2						1	4
		Kurtiella tumida	5	9	3	16	5	7	8	7	7	1	4	8	7	9	1	16	15	12	4	12	8	6	30	21	2	2	5	1		
		Leptopecten latiauratus					5					1							1	1			1		1			1				
		Luciniscia nuttalli			1						1	2														1						
		Lucinoma annulatum	1	3	2		3				1			4	2					1			2	1								
		Macoma nasuta						7	2	5	2	4	2	3	4	3			4	1		2	2			1				1	2	
		Macoma sp					1																	2								
		Macoma yoldiformis	11	17	8	5	7	56	33	38	36	23	30	62	38	29	25	11	24	9	10	20	5	1	15	11	8	24	15	3	7	12
		Modiolatus neglectus																													1	
		Modiolinae						1			2				2	3		1	11	3	2	4	16	7	8	5	6					1
		Neaeromya compressa										1																				
		Neolepton salmoneum						1										4	1	4			35	29	14	5	8					
		Nuculana taphria	5	2		1		10	4	10	6	5	5	9	7	6	3		1		1		1	1	4	1	2	7	7	2	2	1
		Parvilucina tenuisculpta	18	16	11	10	13	12	7	6	18	16	2	6	5	6	5	1	3	7	4	4	4	4	7	2	6	9	4	5	5	5
		Periploma discus			1	1	2	6	2	5		4	1	6		1							1	1				1	1	1		
		Pododesmus sp					2																1									
		Saccella penderi																						1	4							
		Saxicavella nybakkeni																					1									
		Saxidomus nuttalli																							1							
		Solamen columbianum	1	1	1			1			1	1												2	1			2	1		1	



Appendix

10-8. Continued.

Phylum	Class	Species	Station & Replicate																															
			B1					B2					B3					B4					B5					B6						
			1	2	3	4	5	1	2	3	4	5	1	2	3	4	5	1	2	3	4	5	1	2	3	4	5	1	2	3	4	5		
		<i>Solemya pervernica</i>	22	6	4	5																					9	15	3					
		<i>Solen sicarius</i>					4	2	3	2	6	2	2	1	1	1	1	2	1	1	1	3	1	5	1	3	1							
		<i>Tellina carpenteri</i>																		1														
		<i>Tellina idae</i>																																
		<i>Tellina modesta</i>	12	12	12	5	37	39	29	39	68	40	24	41	39	55	49	34	47	35	52	14	21	27	42	25	20	15	25	19	40	19		
		<i>Tellina sp</i>														2				1														
		<i>Tellina sp B</i>	1	3		1			2			3			1	1		3	1	2						1		1						
		Thracioidea								1																								
		<i>Thyasira flexuosa</i>					1			1																		1						
	Caudofoveata	<i>Chaetoderma pacificum</i>	1													1																		
		<i>Falcidens longus</i>														1																1		
	Gastropoda	<i>Aglaja ocelligera</i>														1																		
		<i>Alia carinata</i>																1	2			1	3	1										
		<i>Alia sp</i>												1		14	14	1			11	38	4	2	3									
		<i>Alia tuberosa</i>				6								7	5	2	144	3	12	5	34	28	10	15	2									
		<i>Amphissa versicolor</i>		3																														
		Balcis oldroydae										1																						
		<i>Barleeia haliotiphila</i>																2	2	1				1										
		<i>Bullomorpha sp ABC1</i>																					1											
		<i>Conus californicus</i>								2								2		1	3	15	5	2	2									
		<i>Crepidula glottidiarum</i>													6			1		1					4	1	18	6						
		<i>Crepidula sp</i>							1			1	8	1	4			2																
		<i>Cylchna diegensis</i>	6	4	2	2	4	2	2	2	5	3	8	3	2	2	2				1	1	2	1	2		1	1						
		<i>Dirona picta</i>																		2														
		<i>Epitonium sawinae</i>	1			1	1				1					2			1				1	1	1									
		Eulimidae							1																									
		<i>Eulithidium pulloides</i>																7	1	2	1	1	4		1									
		<i>Glossaulax reclusianus</i>									1	2		1	1					1		1		1		1								
		<i>Hermisenda crassicornis</i>																					1											
		<i>Kelletia kellei</i>																										1						
		<i>Kurtzia arteaga</i>		2	3	1	4							2	1			1	1				1	2										
		<i>Kurtziella plumbea</i>									2				1			2																
		<i>Kurtzia beta</i>								1	1																							
		<i>Lacuna unifasciata</i>																3	1			4	2	1										
		<i>Lirularia acuticostata</i>																1																
		<i>Melanella rosa</i>				1	1			1											1						1							
		<i>Melibe leonina</i>																					4											
		<i>Odostomia sp</i>	1	2	5		12		8	9	9	3	3	1	27	7		1		1	1	3			2	7	3	4	6					
		<i>Ophiidermella inermis</i>	1	2				1	3	1	2											2	2	2		1								
		<i>Philina sp A</i>																																
		<i>Pleurobranchaea californica</i>																	1															
		Polyceridae													1																			
		<i>Polygireulima rutila</i>				1				1		4	1	1													1							
		<i>Rictaxis punctocaelatus</i>					1			3	1		2				3																	
		<i>Sinum scopulosum</i>								1		1											1											
		<i>Skenea coronadoensis</i>	1			2																	1											
		<i>Turbonilla sp A</i>	1			1					1																							
		<i>Turbonilla sp SD1</i>				1					1				1												1							
		<i>Turbonilla sp SD2</i>	2							1	2	1			1																			
		<i>Turbonilla sp SD5</i>				1							1		1											1								
		<i>Volvulella panamica</i>								1		2																						
	Scaphopoda	<i>Gadila aberrans</i>	1	4		1	1	2	3		1	3	2	2		1	1							1	1	1	3			1				



Appendix

10-8. Continued.

Phylum	Class	Species	Station & Replicate																													
			B1					B2					B3					B4					B5					B6				
			1	2	3	4	5	1	2	3	4	5	1	2	3	4	5	1	2	3	4	5	1	2	3	4	5	1	2	3	4	5
Sipuncula	Phascolosomatidea Sipunculidea	Turbellaria sp A											1																			
		Apionsoma misakianum																														
		Sipunculus nudus	1															2														
		Thysanocardia nigra										1					1															1



10.7. Fish and Invertebrate Abundance and Biomass



10-9. Fish abundance by size class (cm) for each replicate trawl.

Scientific Name	Common Name	Size Class (cm)	Abundance				
			TB3		TB6		
			1	2	1	2	
<i>Citharichthys sordidus</i>	Pacific sanddab	6		1		1	
		7		3		1	2
		8	2	4		5	5
		9	1	4		12	2
		10		2		2	4
		11	1	4		5	2
		12	1	6		6	5
		13		3		3	3
		14		5		1	
		15		3		1	1
		16		2			
		17	1	1			
		18		1			
		<i>Citharichthys stigmaeus</i>	speckled sanddab	4			
5	1			1		10	1
6	4			21		36	16
7				16		16	12
8	1			8		28	13
9				3		6	6
10				1		5	1
<i>Citharichthys xanthostigma</i>	longfin sanddab	7		1			
		9		1			
<i>Hypsurus caryi</i>	rainbow seaperch	11		1			
<i>Icelinus quadriseriatus</i>	yellowchin sculpin	7					2
		8					1
<i>Neoclinus blanchardi</i>	sarcastic fringehead	16		1			
<i>Odontopyxis trispinosa</i>	pygmy poacher	7					1
		8					4
		9	1				1
<i>Ophiodon elongatus</i>	lingcod	17					1
		19		1			1
		25		1			
<i>Paralabrax nebulifer</i>	barred sand bass	13					1
<i>Paralichthys californicus</i>	California halibut	62		1			
<i>Parophrys vetulus</i>	English sole	17					1
<i>Pleuronichthys decurrens</i>	curffin sole	8	1				2
		9		1			
		11					1
		14		1			
		15		1			1
		18		1			
<i>Pleuronichthys verticalis</i>	homyhead turbot	8					1
		9					1
		16					2
		17					1
		18					1
		23					1
<i>Porichthys notatus</i>	plainfin midshipman	12					1
<i>Raja inornata</i>	California skate	23.5					1
<i>Scorpaenichthys marmoratus</i>	cabezon	11					1
		12					1
<i>Sebastes crameri</i>	darkblotched rockfish	6					2
		7		1			10
		8		2			15
<i>Sebastes dallii</i>	calico rockfish	7		1			
		8		2			
<i>Sebastes miniatus</i>	vermilion rockfish	6	1				
		7	2	2			6
		8	7	3			1
		9	1				2
<i>Sebastes saxicola</i>	stripetail rockfish	6					1
<i>Syngnathus californiensis</i>	kelp pipefish	19					1
		20					2
		22					2
		23					1
		27					1
<i>Synodus lucioceps</i>	California lizardfish	10					1
		11					4
		12	2				12
		13					9
		15					10
		16					2
		17					2
		18					1
		22					1
		23					2
<i>Ulvicola sanctaerosae</i>	kelp gunnel	9					1
<i>Xystreurus liolepis</i>	fantail sole	31					1
<i>Zaniolepis latipinnis</i>	longspine combfish	15					1
		16					2



10-10. Fish biomass (Kg) by replicate.

Scientific Name	Common Name	Weight (kg)			
		T3		T6	
		1	2	1	2
<i>Citharichthys sordidus</i>	Pacific sanddab	0.16	0.98	0.71	0.54
<i>Citharichthys stigmaeus</i>	speckled sanddab	<0.1	0.28	0.59	0.33
<i>Citharichthys xanthostigma</i>	longfin sanddab	<0.1			
<i>Hypsurus caryi</i>	rainbow seaperch		<0.1		
<i>Icelinus quadriseriatus</i>	yellowchin sculpin			<0.1	<0.1
<i>Neoclinus blanchardi</i>	sarcastic fringehead		0.1		
<i>Odontopyxis trispinosa</i>	pygmy poacher	<0.1		<0.1	<0.1
<i>Ophiodon elongatus</i>	lingcod		0.24	<0.1	<0.1
<i>Paralabrax nebulifer</i>	barred sand bass			<0.1	<0.1
<i>Paralichthys californicus</i>	California halibut		3.5		
<i>Parophrys vetulus</i>	English sole				<0.1
<i>Pleuronichthys decurrens</i>	curffin sole	<0.1	0.37	<0.1	0.12
<i>Pleuronichthys verticalis</i>	hornyhead turbot			0.4	0.49
<i>Porichthys notatus</i>	plainfin midshipman			<0.1	
<i>Raja inornata</i>	California skate				0.31
<i>Scorpaenichthys marmoratus</i>	cabezon			<0.1	<0.1
<i>Sebastes crameri</i>	darkblotched rockfish		<0.1	0.11	0.23
<i>Sebastes dallii</i>	calico rockfish		<0.1		
<i>Sebastes miniatus</i>	vermilion rockfish	0.17	<0.1	<0.1	0.13
<i>Sebastes saxicola</i>	stripetail rockfish			<0.1	
<i>Syngnathus californiensis</i>	kelp pipefish			<0.1	<0.1
<i>Synodus lucioceps</i>	California lizardfish	<0.1		0.36	0.52
<i>Ulvicola sanctaerosae</i>	kelp gunnel			<0.1	
<i>Xystreurus liolepis</i>	fantail sole			0.6	
<i>Zaniolepis latipinnis</i>	longspine combfish			0.16	
	composite	0.13	0.25	0.5	0.45
	Sum	0.46	5.72	3.43	3.12



Appendix

10-11. Invertebrate abundances by replicate.

Scientific Name	Common Name	Abundance			
		TB3		TB6	
		1	2	1	2
<i>Astropecten californicus</i>	California sand star				2
<i>Cancer gracilis</i>	graceful rock crab	1	3	1	2
<i>Loxorhynchus grandis</i>	sheep crab			1	1
<i>Octopus californicus</i>	orange bigeye octopus	1	1	4	2
<i>Octopus rubescens</i>	red octopus			1	
<i>Ophiothrix spiculata</i>	Pacific spiny brittlestar				1
<i>Pisaster brevispinus</i>	shortspined sea star		1		
<i>Pyromaia tuberculata</i>	tuberculate pear crab				1
<i>Sicyonia ingentis</i>	ridgeback rock shrimp		1		
Sum		2	6	7	9

10-12. Invertebrate biomass (Kg) by replicate.

Scientific Name	Common Name	Weight (kg)			
		TB3		TB6	
		1	2	1	2
<i>Astropecten californicus</i>	California sand star				<0.1
<i>Cancer gracilis</i>	graceful rock crab	0.22	0.56	<0.1	0.26
<i>Loxorhynchus grandis</i>	sheep crab			0.32	0.56
<i>Octopus californicus</i>	orange bigeye octopus	<0.1	<0.1	0.12	0.12
<i>Octopus rubescens</i>	red octopus			<0.1	
<i>Ophiothrix spiculata</i>	Pacific spiny brittlestar				<0.1
<i>Pisaster brevispinus</i>	shortspined sea star		0.46		
<i>Pyromaia tuberculata</i>	tuberculate pear crab				<0.1
<i>Sicyonia ingentis</i>	ridgeback rock shrimp		<0.1		
	composite		<0.1	0.1	<0.1
Sum		0.22	1.02	0.54	0.94



10.8. Fish and Bivalve Bioaccumulation Data



10-13. Whole weight, tissue weight and standard length of fish.

STATION TB3				STATION TB6			
Standard Length (mm)	Total Weight (g)	Muscle Weight (g)	Liver Weight (g)	Standard Length (mm)	Total Weight (g)	Muscle Weight (g)	Liver Weight (g)
100	121	3.6	0.29	95	115	2.9	0.21
90	118	2.6	0.27	76	90	1.6	0.29
90	113	3.6	0.28	70	82	0.9	0.13
87	105	2.2	0.20	64	77	0.7	0.18
85	114	2.5	0.24	58	70	0.5	0.09
110	115	4.6	0.43	61	75	0.7	0.18
89	103	2.3	0.14	70	82	0.7	0.18
82	111	1.4	0.12	65	77	0.9	0.15
81	99	2.0	0.18	72	87	0.9	0.26
95	111	1.7	0.16	58	70	0.6	0.13
100	118	2.9	0.08	62	74	0.7	0.24
90	111	3.0	0.22	73	93	1.4	0.22
75	92	1.5	0.20	77	94	1.9	0.30
67	81	1.2	0.09	70	84	0.9	0.25
94	115	3.8	0.44	68	83	0.3	0.06
73	90	1.2	0.15	76	92	1.8	0.32
91	107	2.3	0.40	70	92	1.4	0.36
78	91	1.4	0.17	86	100	2.4	0.31
75	93	2.4	0.09	68	82	1.2	0.19
76	91	1.9	0.19	78	95	0.9	0.11
87	103	2.3	0.28	93	110	2.7	0.42
79	92	1.4	0.10	75	88	1.4	0.33
91	115	3.3	0.17	75	91	1.2	0.32
75	84	1.4	0.19	72	85	1.1	0.23
90	107	2.3	0.46	70	84	1.5	0.23
81	100	1.6	0.14	67	77	0.3	0.06
90	115	2.5	0.22	70	85	1.2	0.22
97	113	2.1	0.11	72	86	1.0	0.24
92	111	2.5	0.33	72	85	1.3	0.17
92	109	1.9	0.36	75	85	1.2	0.28
86	103	2.4	0.33	70	83	1.1	0.29
88	103	2.0	0.18	95	113	2.7	0.64
90	100	1.9	0.21	78	90	0.8	0.08
102	125	4.5	0.51	70	84	1.2	0.24
85	102	2.7	0.32	65	77	0.9	0.23
71	86	1.3	0.20	70	83	1.1	0.15
100	120	3.8	0.48	74	89	1.5	0.29
87	104	2.3	0.17	60	70	0.6	0.06
84	102	2.5	0.33	66	78	1.0	0.23
114	125	3.3	0.73	70	82	1.2	0.18
90	110	2.0	0.36	69	81	1.0	0.29
79	96	1.8	0.21	70	85	0.7	0.11
70	85	1.2	0.16	65	79	1.0	0.28
90	107	2.9	0.17	67	80	1.0	0.20
110	130	6.0	0.57	68	80	0.8	0.25
89	105	1.8	0.22	73	89	1.1	0.24
80	97	1.4	0.25	75	82	1.1	0.15
71	88	1.7	0.20	62	87	0.9	0.05
103	121	4.5	0.40	85	103	2.0	0.34
98	115	3.0	0.23	80	95	1.4	0.30
90	102	1.7	0.20	75	88	1.2	0.16
82	102	1.8	0.19	96	114	2.9	0.53
101	120	3.9	0.43	80	95	1.8	0.42
78	95	1.9	0.18	96	114	2.8	0.37
75	91	1.6	0.18	87	100	2.0	0.04
87	103	2.7	0.25	85	100	1.7	0.21
100	120	3.7	0.42	77	90	1.3	0.35
92	111	3.3	0.54	73	86	1.5	0.32
86	104	2.1	0.19	70	83	1.2	0.22
90	108	2.1	0.60	71	84	1.3	0.35
82	102	2.2	0.24	75	90	1.0	0.06
95	112	3.0	0.24	68	82	1.3	0.28
103	123	3.4	0.22	68	80	1.2	0.19
87	105	2.4	0.44	70	85	1.1	0.17
88	109	2.6	0.20	100	120	2.3	0.20
76	90	1.9	0.17	75	89	1.4	0.29
90	112	3.4	0.41	57	68	0.7	0.22
103	121	4.7	0.61	75	88	1.5	0.32
80	95	1.4	0.13	82	100	1.7	0.06
103	121	3.6	1.40	89	109	2.6	0.36
57	71	0.6	0.09	80	97	2.1	0.36
110	131	5.3	0.73	85	97	2.5	0.39
95	115	2.8	0.45	90	-88	2.6	0.07
92	111	2.0	0.27	83	98	1.9	0.42
90	103	2.5	0.15	76	93	1.5	0.32
100	118	2.8	0.44	86	103	1.8	0.17
88	104	2.6	0.27	90	110	2.2	0.11
93	112	2.7	0.11	83	95	1.6	0.31
85	100	2.7	0.49	78	90	1.4	0.29
80	92	1.7	0.17	72	87	1.2	0.27
78	94	2.1	0.12	80	93	1.5	0.28
74	90	1.2	0.15	72	90	1.2	0.09
				85	100	1.7	0.36
				70	83	1.2	0.27
				70	85	1.4	0.17
				76	85	1.4	0.31
				86	105	2.1	0.40
				75	88	1.1	0.27
				70	85	1.0	0.16
				70	83	0.9	0.18
				69	83	1.2	0.21
				70	81	1.1	0.14
				74	86	0.8	0.12
				72	88	1.5	0.19
				71	81	1.0	0.20
				67	80	1.0	0.14
				79	92	1.2	0.20
				78	93	1.4	0.24
				74	91	1.3	0.17
				75	85	1.2	0.18
				72	88	1.0	0.14
				70	83	1.0	0.15
				70	82	1.1	0.20
				68	81	0.9	0.18
				58	68	0.7	0.09
				70	80	1.0	0.18
				69	81	0.8	0.14
				61	74	0.7	0.16
Count =	Count =	Count =	Count =	Count =	Count =	Count =	Count =
82	82	82	82	108	108	108	108
Total =	Total =	Total =	Total =	Total =	Total =	Total =	Total =
7222.0	8659.0	203.1	23.6	8013.0	9341.0	142.5	24.6
Average =	Average =	Average =	Average =	Average =	Average =	Average =	Average =
88.07	105.60	2.48	0.29	74.19	86.49	1.32	0.23



10-14. Whole weight, tissue weight and total weight of caged bivalves.

Control, Rep 1			Control, Rep 2			Control, Rep 3		
Total Length (mm)	Total Weight (g)	Tissue Weight (g)	Total Length (mm)	Total Weight (g)	Tissue Weight (g)	Total Length (mm)	Total Weight (g)	Tissue Weight (g)
72	33.5	10.3	68	27.6	9.6	61	23.8	8.0
65	28.3	8.5	73	35.5	13.9	73	27.6	8.1
81	40.6	14.8	80	52.1	19.5	62	25.8	6.5
60	20.3	7.8	71	28.0	10.0	73	33.5	11.8
74	38.2	9.1	69	29.4	8.1	60	24.2	6.2
78	31.4	8.4	73	45.2	14.8	73	32.8	11.1
65	26.5	8.5	73	36.3	13.7	59	29.8	8.7
71	27.0	9.5	68	24.7	10.8	75	37.2	11.0
62	21.3	7.1	65	21.4	7.5	74	37.8	11.0
65	24.3	7.0	60	24.5	8.3	59	17.3	5.6
66	22.7	6.7	76	48.2	13.3	70	29.4	11.2
65	20.9	6.1	90	58.0	25.0	70	34.3	13.8
58	19.1	7.0	62	26.4	7.7	62	23.4	8.1
89	38.0	9.0	66	24.3	7.1	63	22.3	6.4
70	33.3	6.7	75	37.1	12.5	68	36.0	12.7
Count = 15	Count = 15	Count = 15	Count = 15	Count = 15	Count = 15	Count = 15	Count = 15	Count = 15
Total = 1041.0	Total = 425.3	Total = 126.5	Total = 81.0	Total = 518.7	Total = 181.8	Total = 1002.0	Total = 435.2	Total = 140.3
Average = 69.4	Average = 28.4	Average = 8.4	Average = 71.3	Average = 34.6	Average = 12.1	Average = 66.8	Average = 29.0	Average = 9.4

B3, Rep 1			B3, Rep 2			B3, Rep 3		
Total Length (mm)	Total Weight (g)	Tissue Weight (g)	Total Length (mm)	Total Weight (g)	Tissue Weight (g)	Total Length (mm)	Total Weight (g)	Tissue Weight (g)
75	43.3	16.3	84	49.1	16.3	71	44.6	15.7
67	32.0	9.2	83	52.7	12.9	69	43.2	11.2
76	41.2	13.3	67	36.4	7.9	79	47.1	14.5
65	25.2	11.0	74	35.1	13.0	58	26.9	8.4
78	51.4	16.2	73	38.7	13.0	64	35.2	7.8
84	58.0	18.2	75	43.2	12.3	86	62.4	20.6
65	33.7	18.6	77	47.6	17.2	63	32.0	10.5
75	39.4	12.2	78	44.1	13.3	75	37.2	11.3
74	40.1	13.9	64	35.7	10.5	75	38.9	13.4
70	43.5	10.7	88	49.6	17.5	80	50.7	19.3
76	48.0	14.8	75	29.7	9.9	73	42.6	14.8
68	38.1	10.9	76	54.4	15.8	73	36.1	12.6
75	37.1	14.4	86	54.6	20.6	79	43.4	14.4
69	34.8	10.0	70	33.3	9.0	64	29.0	8.9
83	49.5	17.8	75	40.2	13.9	57	27.6	7.0
Count = 15	Count = 15	Count = 15	Count = 15	Count = 15	Count = 15	Count = 15	Count = 15	Count = 15
Total = 1100.0	Total = 615.5	Total = 207.5	Total = 81.0	Total = 644.3	Total = 203.0	Total = 1066.0	Total = 596.8	Total = 190.4
Average = 73.3	Average = 41.0	Average = 13.8	Average = 76.3	Average = 43.0	Average = 13.5	Average = 71.1	Average = 39.8	Average = 12.7

B4, Rep 1			B4, Rep 2			B4, Rep 3		
Total Length (mm)	Total Weight (g)	Tissue Weight (g)	Total Length (mm)	Total Weight (g)	Tissue Weight (g)	Total Length (mm)	Total Weight (g)	Tissue Weight (g)
78	41.3	12.3	77	48.0	16.3	70	43.9	13.4
71	28.5	9.9	75	35.8	10.7	84	34.3	12.2
91	61.2	21.7	66	37.6	11.0	77	38.0	12.5
82	48.1	14.9	87	48.4	15.6	71	43.1	15.2
66	33.6	10.2	72	47.2	15.1	79	46.8	17.7
70	36.6	12.1	65	30.2	10.4	79	43.6	13.8
82	48.6	16.7	64	24.9	8.8	70	38.2	9.8
77	37.2	10.3	85	40.4	9.5	61	42.8	14.8
80	56.0	12.5	81	50.2	19.8	72	29.5	11.7
74	40.7	14.3	73	38.0	11.3	68	31.9	10.1
75	43.4	13.8	79	38.7	13.4	66	30.7	11.5
75	35.8	12.1	78	40.2	13.0	90	32.6	16.4
63	30.9	10.1	70	36.0	9.7	77	36.9	14.7
78	36.1	11.3	85	62.3	21.6	75	31.7	13.3
77	44.8	11.3	86	47.4	13.6	65	30.0	9.9
Count = 15	Count = 15	Count = 15	Count = 15	Count = 15	Count = 15	Count = 15	Count = 15	Count = 15
Total = 1139.0	Total = 622.7	Total = 193.3	Total = 1143.0	Total = 625.2	Total = 199.7	Total = 1124.0	Total = 553.9	Total = 197.0
Average = 75.9	Average = 41.5	Average = 12.9	Average = 76.2	Average = 41.7	Average = 13.3	Average = 74.9	Average = 36.9	Average = 13.1

B6, Rep 1			B6, Rep 2			B6, Rep 3		
Total Length (mm)	Total Weight (g)	Tissue Weight (g)	Total Length (mm)	Total Weight (g)	Tissue Weight (g)	Total Length (mm)	Total Weight (g)	Tissue Weight (g)
78	45.0	15.0	66	29.8	12.3	67	44.4	13.0
82	51.2	12.2	68	36.8	9.8	74	44.3	14.4
85	65.3	21.3	73	46.7	18.7	80	40.1	11.1
73	42.5	14.0	74	32.2	10.8	68	37.5	10.4
65	30.5	10.1	77	30.6	10.6	66	27.8	10.0
87	39.8	13.3	70	43.2	12.2	77	37.5	11.8
78	49.0	15.9	76	34.9	12.2	76	39.5	12.2
75	38.1	12.4	88	35.2	11.3	86	45.1	14.4
69	34.6	12.5	87	53.4	17.7	74	38.8	11.4
76	41.8	14.5	72	35.9	8.5	67	25.4	7.8
71	39.0	14.1	74	41.9	15.0	70	30.3	7.2
76	43.4	17.6	75	52.0	11.0	104	66.7	27.8
81	46.8	13.6	86	39.7	15.2	70	35.0	11.1
75	45.4	12.6	96	63.5	23.1	81	49.7	13.9
96	61.1	19.6	73	35.8	12.9	78	44.2	12.5
Count = 15	Count = 15	Count = 15	Count = 15	Count = 15	Count = 15	Count = 15	Count = 15	Count = 15
Total = 1167.0	Total = 673.4	Total = 218.5	Total = 1155.0	Total = 611.5	Total = 201.1	Total = 1138.0	Total = 606.1	Total = 189.1
Average = 77.8	Average = 44.9	Average = 14.6	Average = 77.0	Average = 40.8	Average = 13.4	Average = 75.9	Average = 40.4	Average = 12.6



Appendix

10-15. Fish and bivalve tissue concentrations by replicate for all constituents measured.

Constituent	Replicate	Fish Muscle		Fish Liver		Bivalve			
		TB3	TB6	TB3	TB6	Control	B3	B4	B6
General Chemistry (µg/dry g)									
% Lipids (detection limit = 0.01)	#1	1.33	1.53	38.78	28.65	9.25	5.96	7.38	6.45
	#2	1.47	2.29	38.57	27.72	7.55	5.83	6.99	7.48
	#3	<u>1.11</u>	<u>2.10</u>	<u>37.56</u>	<u>27.12</u>	<u>7.64</u>	<u>6.05</u>	<u>6.51</u>	<u>8.15</u>
	Mean =	1.30	1.97	38.30	27.83	8.15	5.95	6.96	7.36
	S.D. =	0.18	0.40	0.65	0.77	0.96	0.11	0.44	0.86
Mean for each Tissue =	1.638		33.067		7.103				
% Moisture (detection limit = 0.1)	#1	80.9	80.0	68.5	71.9	82.6	79.3	81.4	81.0
	#2	80.8	80.1	NS	NS	82.2	80.0	80.7	80.1
	#3	<u>80.4</u>	<u>80.0</u>	NS	NS	<u>83.6</u>	<u>80.4</u>	<u>81.2</u>	<u>80.0</u>
	Mean =	80.7	80.0	68.5	71.9	82.8	79.9	81.1	80.4
	S.D. =	0.3	0.1	NA	NA	0.7	0.6	0.4	0.6
Mean for each Tissue =	80.4		70.2		81.0				
Metals (µg/dry g)									
(detection limit = 0.025 µg/dry g)									
Arsenic	#1	7.131	3.884	8.455	6.064	7.862	9.487	9.285	9.706
	#2	7.282	4.249	8.041	6.229	7.123	9.548	9.730	9.102
	#3	<u>7.557</u>	<u>4.265</u>	<u>8.807</u>	<u>6.047</u>	<u>7.468</u>	<u>8.821</u>	<u>9.263</u>	<u>9.537</u>
	Mean =	7.323	4.133	8.434	6.113	7.484	9.285	9.426	9.448
	S.D. =	0.216	0.216	0.383	0.101	0.370	0.403	0.264	0.312
Mean for each Tissue =	5.728		7.274		8.911				
Cadmium	#1	0.041	0.036	8.121	9.869	4.340	4.852	5.750	5.515
	#2	0.042	0.033	8.104	9.612	4.179	5.839	6.099	5.231
	#3	<u>0.045</u>	<u>0.031</u>	<u>7.942</u>	<u>10.076</u>	<u>4.624</u>	<u>5.088</u>	<u>5.744</u>	<u>5.021</u>
	Mean =	0.043	0.033	8.056	9.852	4.381	5.260	5.864	5.256
	S.D. =	0.002	0.003	0.099	0.232	0.225	0.515	0.203	0.248
Mean for each Tissue =	0.038		8.954		5.190				
Chromium	#1	0.053	0.053	0.048	0.122	0.592	0.832	0.940	0.738
	#2	0.094	0.062	0.069	0.104	0.542	0.888	0.897	0.783
	#3	<u>0.043</u>	<u>0.059</u>	<u>0.167</u>	<u>0.112</u>	<u>0.662</u>	<u>0.953</u>	<u>0.956</u>	<u>0.726</u>
	Mean =	0.063	0.058	0.095	0.113	0.599	0.891	0.931	0.749
	S.D. =	0.027	0.005	0.064	0.009	0.060	0.061	0.031	0.030
Mean for each Tissue =	0.061		0.104		0.792				
Copper	#1	1.605	1.151	11.013	10.151	6.065	6.857	6.954	7.242
	#2	1.499	1.265	10.766	10.069	6.073	6.518	7.004	6.935
	#3	<u>2.970</u>	<u>1.161</u>	<u>10.757</u>	<u>10.478</u>	<u>5.701</u>	<u>7.080</u>	<u>7.077</u>	<u>7.086</u>
	Mean =	2.025	1.192	10.845	10.233	5.946	6.818	7.012	7.088
	S.D. =	0.820	0.063	0.145	0.216	0.213	0.283	0.062	0.154
Mean for each Tissue =	1.609		10.539		6.716				
Lead	#1	0.027	0.046	0.439	0.690	1.329	1.298	1.332	1.245
	#2	0.025	0.035	0.484	0.667	1.095	1.403	1.327	1.275
	#3	<u>0.031</u>	<u>0.025</u>	<u>0.402</u>	<u>0.721</u>	<u>1.249</u>	<u>1.557</u>	<u>1.196</u>	<u>1.156</u>
	Mean =	0.028	0.035	0.442	0.693	1.224	1.419	1.285	1.225
	S.D. =	0.003	0.011	0.041	0.027	0.119	0.130	0.077	0.062
Mean for each Tissue =	0.032		0.567		1.289				
Mercury (det. Limit = 0.00001 µg/dry g)	#1	0.1596	0.1902	0.1013	0.1259	0.0679	0.0415	0.0432	0.0418
	#2	0.1583	0.1842	0.1015	0.1218	0.0695	0.0394	0.0429	0.0375
	#3	<u>0.1679</u>	<u>0.1856</u>	<u>0.0961</u>	<u>0.1201</u>	<u>0.0693</u>	<u>0.0426</u>	<u>0.0420</u>	<u>0.0422</u>
	Mean =	0.1619	0.1867	0.0996	0.1226	0.0689	0.0411	0.0427	0.0405
	S.D. =	0.0052	0.0031	0.0031	0.0030	0.0009	0.0016	0.0006	0.0026
Mean for each Tissue =	0.174		0.111		0.048				
Nickel	#1	0.025	0.025	0.084	0.085	0.484	0.794	0.931	0.775
	#2	0.025	0.025	0.084	0.092	0.491	0.837	0.841	0.719
	#3	<u>0.025</u>	<u>0.075</u>	<u>0.099</u>	<u>0.118</u>	<u>0.556</u>	<u>0.796</u>	<u>0.912</u>	<u>0.709</u>
	Mean =	0.025	0.042	0.089	0.098	0.510	0.809	0.895	0.734
	S.D. =	0.000	0.029	0.009	0.017	0.040	0.024	0.047	0.036
Mean for each Tissue =	0.033		0.094		0.737				

NS=not enough tissue for replicate analysis.



Appendix

10-15. continued.

Constituent	Replicate	Fish Muscle		Fish Liver		Control	Bivalve		
		TB3	TB6	TB3	TB6		B3	B4	B6
Metals (µg/dry g)									
Selenium	#1	1.277	1.000	5.186	7.315	2.220	2.952	3.110	3.171
	#2	1.374	1.058	4.980	7.645	2.198	2.632	3.287	2.905
	#3	<u>1.251</u>	<u>0.917</u>	<u>5.808</u>	<u>7.602</u>	<u>2.314</u>	<u>2.942</u>	<u>3.102</u>	<u>3.209</u>
	Mean =	1.301	0.992	5.325	7.521	2.244	2.842	3.166	3.095
	S.D. =	0.065	0.071	0.431	0.179	0.062	0.182	0.105	0.166
	Mean for each Tissue =	1.15		6.42		2.837			
Silver	#1	0.025	0.025	0.095	0.285	0.168	0.184	0.210	0.292
	#2	0.025	0.025	0.102	0.287	0.232	0.267	0.267	0.197
	#3	<u>0.025</u>	<u>0.025</u>	<u>0.098</u>	<u>0.300</u>	<u>0.219</u>	<u>0.261</u>	<u>0.295</u>	<u>0.235</u>
	Mean =	0.025	0.025	0.098	0.291	0.206	0.237	0.257	0.241
	S.D. =	0.000	0.000	0.004	0.008	0.034	0.046	0.043	0.048
	Mean for each Tissue =	0.03		0.19		0.236			
Zinc	#1	18.53	17.22	67.80	72.81	92.71	126.91	128.00	128.44
	#2	19.21	15.28	66.75	71.51	83.72	120.15	122.08	135.08
	#3	<u>19.57</u>	<u>16.60</u>	<u>64.51</u>	<u>73.54</u>	<u>89.22</u>	<u>122.97</u>	<u>124.20</u>	<u>128.39</u>
	Mean =	19.106	16.365	66.349	72.619	88.547	123.339	124.759	130.636
	S.D. =	0.527	0.993	1.681	1.032	4.532	3.395	3.000	3.851
	Mean for each Tissue =	17.74		69.48		116.820			
Complex Organics (ng/dry Kg)									
Total DDT ¹	#1	10.6	20.3	956.9	435.2	52.6	19.6	23.3	15.9
	#2	10.8	20.4	854.8	429.5	47.7	18.3	16.7	18.7
	#3	<u>11.3</u>	<u>24.6</u>	<u>926.9</u>	<u>406.9</u>	<u>49.2</u>	<u>18.2</u>	<u>23.9</u>	<u>23.2</u>
	Mean =	10.9	21.8	912.9	423.9	49.8	18.7	21.3	19.3
	S.D. =	0.4	2.5	52.5	15.0	2.5	0.8	4.0	3.7
	Mean for each Tissue =	16.33		668.37		27.3			
Total Chlordane ¹	#1	0.0	0.0	10.6	0.0	0.0	0.0	0.0	0.0
	#2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	#3	<u>0.0</u>	<u>0.0</u>	<u>9.3</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>
	Mean =	0.0	0.0	6.6	0.0	0.0	0.0	0.0	0.0
	S.D. =	0.0	0.0	5.8	0.0	0.0	0.0	0.0	0.0
	Mean for each Tissue =	0.00		3.32		0.0			
Total HCHs ¹	#1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	#2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	#3	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>
	Mean =	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	S.D. =	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	Mean for each Tissue =	0.0		0.0		0.0			
Aldrin (detection limit = 1.0 ng/g)	#1	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0
	#2	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0
	#3	<u>1.0</u>	<u>1.0</u>	<u>1.0</u>	<u>1.0</u>	<u>1.0</u>	<u>1.0</u>	<u>1.0</u>	<u>1.0</u>
	Mean =	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0
	S.D. =	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	Mean for each Tissue =	1.0		1.0		1.0			
Dieldrin (detection limit = 1.0 ng/g)	#1	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0
	#2	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0
	#3	<u>1.0</u>	<u>1.0</u>	<u>1.0</u>	<u>1.0</u>	<u>1.0</u>	<u>1.0</u>	<u>1.0</u>	<u>1.0</u>
	Mean =	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0
	S.D. =	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	Mean for each Tissue =	1.0		1.0		1.0			
Heptachlor (detection limit = 1.0 ng/g)	#1	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0
	#2	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0
	#3	<u>1.0</u>	<u>1.0</u>	<u>1.0</u>	<u>1.0</u>	<u>1.0</u>	<u>1.0</u>	<u>1.0</u>	<u>1.0</u>
	Mean =	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0
	S.D. =	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	Mean for each Tissue =	1.0		1.0		1.0			

1. Complex organic derivatives are listed in Table 10-16.



Appendix

10-15. continued.

Constituent	Replicate	Fish Muscle		Fish Liver		Control	Bivalve		
		TB3	TB6	TB3	TB6		B3	B4	B6
Complex Organics (ng/dry Kg)									
Hexachlorobenzene (detection limit = 1.0 ng/g)	#1	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0
	#2	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0
	#3	<u>1.0</u>	<u>1.0</u>	<u>1.0</u>	<u>1.0</u>	<u>1.0</u>	<u>1.0</u>	<u>1.0</u>	<u>1.0</u>
	Mean =	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0
	S.D. =	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	Mean for each Tissue =	1.0		1.0		1.0			
Mirex (detection limit = 1.0 ng/g)	#1	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0
	#2	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0
	#3	<u>1.0</u>	<u>1.0</u>	<u>1.0</u>	<u>1.0</u>	<u>1.0</u>	<u>1.0</u>	<u>1.0</u>	<u>1.0</u>
	Mean =	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0
	S.D. =	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	Mean for each Tissue =	1.0		1.0		1.0			
Polychlorinated Biphenyls (PCBs) ¹	#1	0.0	0.0	0.0	29.8	0.0	0.0	0.0	0.0
	#2	0.0	0.0	0.0	36.5	0.0	0.0	0.0	0.0
	#3	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>37.6</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>
	Mean =	0.0	0.0	0.0	34.6	0.0	0.0	0.0	0.0
	S.D. =	0.0	0.0	0.0	4.2	0.0	0.0	0.0	0.0
	Mean for each Tissue =	0.00		17.32		0.0			
Arochlors ¹	#1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	#2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	#3	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>
	Mean =	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	S.D. =	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	Mean for each Tissue =	0.00		0.00		0.0			
Polynuclear Aromatic Hydrocarbons (PAHs) ¹	#1	34.3	27.8	224.5	100.8	77.5	64.7	67.2	80.2
	#2	29.6	31.5	221.1	141.8	65.1	58.9	71.8	68.5
	#3	<u>28.3</u>	<u>29.4</u>	<u>211.9</u>	<u>139.2</u>	<u>71.4</u>	<u>59.6</u>	<u>76.9</u>	<u>76.1</u>
	Mean =	30.7	29.6	219.2	127.3	71.3	61.1	72.0	74.9
	S.D. =	3.2	1.9	6.5	23.0	6.2	3.2	4.9	5.9
	Mean for each Tissue =	30.15		173.22		69.83			
1-Methylnaphthalene (det. Limit = 1 ng/dryg)	#1	2.5	1.3	7.2	5.7	4.0	2.6	3.8	3.1
	#2	2.2	1.8	4.2	9.1	4.1	2.7	2.7	4.0
	#3	<u>2.5</u>	<u>1.3</u>	<u>7.1</u>	<u>10.9</u>	<u>4.0</u>	<u>2.0</u>	<u>3.2</u>	<u>3.5</u>
	Mean =	2.4	1.5	6.2	8.6	4.0	2.4	3.2	3.5
	S.D. =	0.2	0.3	1.7	2.6	0.1	0.4	0.6	0.5
	Mean for each Tissue =	1.93		7.37		3.31			
1-Methylphenanthrene (det. Limit = 1 ng/dryg)	#1	2.3	1.9	10.7	5.9	7.3	6.0	6.2	5.9
	#2	1.8	1.9	14.2	13.2	4.5	6.5	6.5	5.8
	#3	<u>1.8</u>	<u>1.9</u>	<u>14.5</u>	<u>10.7</u>	<u>5.3</u>	<u>6.7</u>	<u>7.5</u>	<u>5.8</u>
	Mean =	2.0	1.9	13.1	9.9	5.7	6.4	6.7	5.8
	S.D. =	0.3	0.0	2.1	3.7	1.4	0.4	0.7	0.1
	Mean for each Tissue =	1.93		11.53		6.17			
2-Methylnaphthalene (det. Limit = 1 ng/dryg)	#1	3.9	2.1	18.8	13.9	9.4	4.6	7.7	6.5
	#2	4.8	3.7	1.0	18.3	8.4	8.2	7.8	10.2
	#3	<u>4.5</u>	<u>1.4</u>	<u>2.7</u>	<u>18.6</u>	<u>9.1</u>	<u>4.5</u>	<u>8.3</u>	<u>6.2</u>
	Mean =	4.4	2.4	7.5	16.9	9.0	5.8	7.9	7.6
	S.D. =	0.5	1.2	9.8	2.6	0.5	2.1	0.3	2.2
	Mean for each Tissue =	3.40		12.22		7.58			
2,3,5-Trimethylnaphthalene (det. Limit = 1 ng/dryg)	#1	1.0	1.0	1.0	1.0	1.6	1.7	1.4	1.6
	#2	1.0	1.0	1.0	13.7	1.5	1.2	1.3	1.0
	#3	<u>1.0</u>	<u>1.0</u>	<u>1.0</u>	<u>1.0</u>	<u>1.9</u>	<u>1.0</u>	<u>1.0</u>	<u>1.0</u>
	Mean =	1.0	1.0	1.0	5.2	1.7	1.3	1.2	1.2
	S.D. =	0.0	0.0	0.0	7.3	0.2	0.4	0.2	0.3
	Mean for each Tissue =	1.00		3.12		1.35			

1. Complex organic derivatives are listed in Table 10-16.



Appendix

10-15. continued.

Constituent	Replicate	Fish Muscle		Fish Liver		Control	Bivalve		
		TB3	TB6	TB3	TB6		B3	B4	B6
2,6-Dimethylnaphthalene (det. Limit = 1 ng/dry g)	#1	3.8	1.3	9.0	7.0	4.1	2.0	3.8	2.9
	#2	4.8	2.1	14.3	9.5	3.2	3.1	3.0	3.2
	#3	<u>3.0</u>	<u>1.7</u>	<u>10.7</u>	<u>9.2</u>	<u>3.9</u>	<u>2.1</u>	<u>3.5</u>	<u>2.8</u>
	Mean =	3.9	1.7	11.3	8.6	3.7	2.4	3.4	3.0
	S.D. =	0.9	0.4	2.7	1.4	0.5	0.6	0.4	0.2
Mean for each Tissue =	2.78		9.95		3.13				
Acenaphthene (det. Limit = 1 ng/dry g)	#1	1.0	1.5	1.0	1.0	3.1	1.0	1.0	1.0
	#2	1.0	2.3	1.0	1.0	2.2	1.0	1.0	1.0
	#3	<u>1.0</u>	<u>3.0</u>	<u>1.0</u>	<u>1.0</u>	<u>3.0</u>	<u>1.0</u>	<u>1.0</u>	<u>1.0</u>
	Mean =	1.0	2.3	1.0	1.0	2.8	1.0	1.0	1.0
	S.D. =	0.0	0.8	0.0	0.0	0.5	0.0	0.0	0.0
Mean for each Tissue =	1.63		1.00		1.44				
Biphenyl (det. Limit = 1 ng/dry g)	#1	2.0	1.7	7.1	8.3	5.2	2.4	3.3	2.9
	#2	2.5	2.2	6.8	12.3	4.1	2.2	2.7	3.5
	#3	<u>2.3</u>	<u>1.2</u>	<u>5.8</u>	<u>11.8</u>	<u>4.1</u>	<u>2.3</u>	<u>3.5</u>	<u>3.1</u>
	Mean =	2.3	1.7	6.6	10.8	4.5	2.3	3.2	3.2
	S.D. =	0.3	0.5	0.7	2.2	0.6	0.1	0.4	0.3
Mean for each Tissue =	1.98		8.68		3.28				
Benzo[a]anthracene (det. Limit = 1 ng/dry g)	#1	1.0	1.0	69.9	1.0	1.0	1.0	1.0	1.0
	#2	1.0	1.0	76.0	1.0	1.0	1.0	1.0	1.0
	#3	<u>1.0</u>	<u>1.0</u>	<u>73.6</u>	<u>1.0</u>	<u>1.0</u>	<u>1.0</u>	<u>1.0</u>	<u>1.0</u>
	Mean =	1.0	1.0	73.2	1.0	1.0	1.0	1.0	1.0
	S.D. =	0.0	0.0	3.1	0.0	0.0	0.0	0.0	0.0
Mean for each Tissue =	1.00		37.08		1.00				
Benzo[b]fluoranthene (det. Limit = 1 ng/dry g)	#1	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0
	#2	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0
	#3	<u>1.0</u>	<u>1.0</u>	<u>1.0</u>	<u>1.0</u>	<u>1.0</u>	<u>1.0</u>	<u>1.0</u>	<u>1.0</u>
	Mean =	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0
	S.D. =	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Mean for each Tissue =	1.00		1.00		1.00				
Benzo[e]pyrene (det. Limit = 1 ng/dry g)	#1	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0
	#2	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0
	#3	<u>1.0</u>	<u>1.0</u>	<u>1.0</u>	<u>1.0</u>	<u>1.0</u>	<u>1.0</u>	<u>1.0</u>	<u>1.0</u>
	Mean =	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0
	S.D. =	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Mean for each Tissue =	1.00		1.00		1.00				
Benzo[g,h,i]perylene (det. Limit = 1 ng/dry g)	#1	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0
	#2	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0
	#3	<u>1.0</u>	<u>1.0</u>	<u>1.0</u>	<u>1.0</u>	<u>1.0</u>	<u>1.0</u>	<u>1.0</u>	<u>1.0</u>
	Mean =	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0
	S.D. =	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Mean for each Tissue =	1.00		1.00		1.00				
Fluoranthene (det. Limit = 1 ng/dry g)	#1	8.3	5.3	12.0	20.8	13.9	17.6	6.7	14.0
	#2	3.0	4.4	31.2	21.2	7.6	13.1	18.8	15.1
	#3	<u>4.6</u>	<u>4.1</u>	<u>29.0</u>	<u>21.2</u>	<u>8.0</u>	<u>13.1</u>	<u>16.2</u>	<u>16.2</u>
	Mean =	5.3	4.6	24.1	21.1	9.8	14.6	13.9	15.1
	S.D. =	2.7	0.6	10.5	0.2	3.5	2.6	6.4	1.1
Mean for each Tissue =	4.95		22.57		13.36				
Naphthalene (det. Limit = 1 ng/dry g)	#1	12.5	5.6	40.5	27.7	38.9	10.8	15.9	11.5
	#2	15.3	7.6	38.0	39.8	31.2	12.3	11.2	15.0
	#3	<u>13.0</u>	<u>5.8</u>	<u>35.1</u>	<u>47.1</u>	<u>38.2</u>	<u>9.1</u>	<u>13.3</u>	<u>14.0</u>
	Mean =	13.6	6.3	37.9	38.2	36.1	10.7	13.5	13.5
	S.D. =	1.5	1.1	2.7	9.8	4.3	1.6	2.4	1.8
Mean for each Tissue =	9.97		38.03		18.45				
Perylene (det. Limit = 1 ng/dry g)	#1	1.0	1.0	27.9	1.0	1.0	1.0	1.0	1.0
	#2	1.0	1.0	54.0	1.0	1.0	1.0	1.0	1.0
	#3	<u>1.0</u>	<u>1.0</u>	<u>24.6</u>	<u>1.0</u>	<u>1.0</u>	<u>1.0</u>	<u>1.0</u>	<u>1.0</u>
	Mean =	1.0	1.0	35.5	1.0	1.0	1.0	1.0	1.0
	S.D. =	0.0	0.0	16.1	0.0	0.0	0.0	0.0	0.0
Mean for each Tissue =	1.00		18.25		1.00				

1. Complex organic derivatives are listed in Table 10-16.



Appendix

10-16. Complex organics (ng/dry g) in fish muscle and liver tissues.

Tissue ¹ Station Replicate	Fish Muscle						Fish Liver					
	Trawl Station TB3			Trawl Station TB6			Trawl Station TB3			Trawl Station TB6		
	1	2	3	1	2	3	1	2	3	1	2	3
DDT												
2,4'-DDD	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
2,4'-DDE	0.0	0.0	0.0	2.4	1.9	2.3	29.0	18.5	22.1	38.2	25.3	29.4
2,4'-DDT	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
4,4'-DDD	0.0	0.0	0.0	0.0	0.0	0.0	111.5	150.7	123.6	0.0	0.0	0.0
4,4'-DDE	10.6	10.8	11.3	17.9	18.5	22.3	772.9	640.3	732.6	397.0	404.2	377.5
<u>4,4'-DDT</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>43.5</u>	<u>45.3</u>	<u>48.6</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>
Sum =	10.6	10.8	11.3	20.3	20.4	24.6	956.9	854.8	926.9	435.2	429.5	406.9
Chlordane												
Chlordane-alpha	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Chlordane-gamma	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
cis-Nonachlor	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Oxychlordane	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
<u>trans-Nonachlor</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>10.6</u>	<u>0.0</u>	<u>9.3</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>
Sum =	0.0	0.0	0.0	0.0	0.0	0.0	10.6	0.0	9.3	0.0	0.0	0.0
Hexachlorocyclohexane (HCH)												
BHC-alpha	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
BHC-beta	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
BHC-delta	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
<u>BHC-gamma</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>
Sum =	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Polynuclear Aromatic Hydrocarbons (PAH's)												
PCB003	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB008	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB018	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB028	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB031	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB033	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB037	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB044	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB049	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB052	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB056(060)	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB066	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB070	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB074	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB077	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB081	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB087	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	6.2
PCB095	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB097	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB099	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	3.5	4.7	3.0
PCB101	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	5.5	5.1	6.1
PCB105	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB110	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB114	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB118	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB119	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB123	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0

1. Minimum detection limits, reporting limits and methods are listed in table 10-18.



Appendix

10-16. continued.

Tissue ¹ Station Replicate	Fish Muscle						Fish Liver					
	Trawl Station TB3			Trawl Station TB6			Trawl Station TB3			Trawl Station TB6		
	1	2	3	1	2	3	1	2	3	1	2	3
PCB126	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB128	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB138	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	8.0	10.9	9.6
PCB141	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB149	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2.0	0.0
PCB151	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB153	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	12.8	13.8	12.7
PCB156	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB157	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB158	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB167	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB168/132	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB169	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB170	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB174	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB177	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB180	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB183	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB187	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB189	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB194	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB195	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB199(200)	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB201	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB206	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
<u>PCB209</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>
Sum =	1.0	2.0	3.0	1.0	2.0	3.0	1.0	2.0	3.0	30.8	38.5	40.6
Aroclors												
Aroclor 1016	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Aroclor 1221	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Aroclor 1232	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Aroclor 1242	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Aroclor 1248	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Aroclor 1254	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Aroclor 1260	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>
sum=	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Polychlorinated Biphenyls (PCB's)												
Acenaphthylene	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Anthracene	2.0	2.9	2.2	2.4	2.4	2.4	18.2	20.3	19.4	9.0	0.0	9.4
Benz[a]anthracene	0.0	0.0	0.0	0.0	0.0	0.0	69.9	76.0	73.6	0.0	0.0	0.0
Benzo[a]pyrene	0.0	0.0	0.0	0.0	0.0	0.0	12.4	11.8	10.9	0.0	0.0	0.0
Benzo[b]fluoranthene	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Benzo[g,h,i]perylene	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Benzo[k]fluoranthene	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Chrysene	0.0	0.0	0.0	0.0	1.3	0.0	12.3	9.6	12.0	0.0	0.0	0.0
Dibenz[a,h]anthracene	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Fluorene	4.4	3.6	3.4	4.3	4.3	4.9	18.5	10.3	8.1	13.7	19.2	22.3
Indeno[1,2,3-c,d]pyrene	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Phenanthrene	24.3	20.7	19.4	18.3	19.9	19.5	80.7	80.7	75.6	62.2	97.0	94.5
<u>Pyrene</u>	<u>3.6</u>	<u>2.4</u>	<u>3.3</u>	<u>2.8</u>	<u>3.6</u>	<u>2.6</u>	<u>12.5</u>	<u>12.4</u>	<u>12.3</u>	<u>15.9</u>	<u>25.6</u>	<u>13.0</u>
Sum =	34.3	29.6	28.3	27.8	31.5	29.4	224.5	221.1	211.9	100.8	141.8	139.2

1. Minimum detection limits, reporting limits and methods are listed in table 10-18.



Appendix

10-17. Complex organics (ng/dry g) in caged bivalve tissues.

Tissue ¹ Station Replicate	Mussel Tissue											
	Control			Station B3			Station B4			Station B6		
	1	2	3	1	2	3	1	2	3	1	2	3
DDT & Derivatives												
2,4'-DDD	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
2,4'-DDE	7.9	3.6	5.5	2.2	0.0	0.0	0.0	0.0	3.6	0.0	0.0	3.9
2,4'-DDT	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
4,4'-DDD	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
4,4'-DDE	44.7	44.1	43.7	17.4	18.3	18.2	23.3	16.7	20.3	15.9	18.7	19.3
<u>4,4'-DDT</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>
Sum =	52.6	47.7	49.2	19.6	18.3	18.2	23.3	16.7	23.9	15.9	18.7	23.2
Chlordane												
Chlordane-alpha	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Chlordane-gamma	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
cis-Nonachlor	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Oxychlordane	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
<u>trans-Nonachlor</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>
Sum =	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Hexachlorocyclohexane (HCH)												
BHC-alpha	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
BHC-beta	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
BHC-delta	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
<u>BHC-gamma</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>
Sum =	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Biphenyls (PCB's)												
PCB003	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB008	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB018	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB028	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB031	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB033	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB037	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB044	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB049	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB052	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB056(060)	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB066	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB070	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB074	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB077	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB081	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB087	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB095	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB097	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB099	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB101	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB105	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB110	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB114	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB118	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB119	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB123	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB126	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0

1. Minimum detection limits, reporting limits and methods are listed in table 10-18.



Appendix

10-17. continued.

Tissue ¹ Station Replicate	Mussel Tissue											
	Control			Station B3			Station B4			Station B6		
	1	2	3	1	2	3	1	2	3	1	2	3
PCB128	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB138	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB141	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB149	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB151	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB153	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB156	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB157	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB158	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB167	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB168/132	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB169	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB170	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB174	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB177	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB180	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB183	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB187	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB189	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB194	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB195	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB199(200)	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB201	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PCB206	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
<u>PCB209</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>
Sum =	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Aroclors												
Aroclor 1016	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Aroclor 1221	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Aroclor 1232	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Aroclor 1242	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Aroclor 1248	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Aroclor 1254	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
<u>Aroclor 1260</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>
Sum =	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Polynuclear Aromatic Hydrocarbons (PAH's)												
Acenaphthylene	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Anthracene	6.0	4.9	4.9	4.3	4.0	3.5	5.3	6.1	5.6	6.3	6.2	4.9
Benz[a]anthracene	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Benzo[a]pyrene	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Benzo[b]fluoranthene	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Benzo[g,h,i]perylene	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Benzo[k]fluoranthene	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Chrysene	0.0	0.0	0.0	2.7	0.0	2.2	2.4	0.0	0.0	0.0	0.0	0.0
Dibenz[a,h]anthracene	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Fluorene	12.8	10.4	12.2	7.4	6.6	7.6	11.0	9.6	10.4	8.8	9.4	10.1
Indeno[1,2,3-c,d]pyrene	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Phenanthrene	47.0	42.0	47.5	40.7	40.3	38.2	42.7	44.9	49.3	41.0	43.8	45.4
<u>Pyrene</u>	<u>11.7</u>	<u>7.8</u>	<u>6.8</u>	<u>9.6</u>	<u>8.0</u>	<u>8.1</u>	<u>5.8</u>	<u>11.2</u>	<u>11.6</u>	<u>24.1</u>	<u>9.1</u>	<u>15.7</u>
Sum =	77.5	65.1	71.4	64.7	58.9	59.6	67.2	71.8	76.9	80.2	68.5	76.1



10-18 Tissue chemistry detection limits and methods

Parameter	MDL	RL	Units (dry wt.)	Method
General Chemistry				
Percent Lipids	0.01	0.05	%	Gravimetric
Percent Solids	0.1	0.1	%	SM 2540B
Trace Metals				
Arsenic	0.025	0.05	µg/g	EPA 6020
Cadmium	0.025	0.05	µg/g	EPA 6020
Chromium	0.025	0.05	µg/g	EPA 6020
Copper	0.025	0.05	µg/g	EPA 6020
Lead	0.025	0.05	µg/g	EPA 6020
Mercury	0.00001	0.00002	µg/g	EPA 245.7
Nickel	0.025	0.05	µg/g	EPA 6020
Selenium	0.025	0.05	µg/g	EPA 6020
Silver	0.025	0.05	µg/g	EPA 6020
Zinc	0.025	0.05	µg/g	EPA 6020
Chlorinated Pesticides				
2,4'-DDD	1	5	ng/g	EPA 8270C
2,4'-DDE	1	5	ng/g	EPA 8270C
2,4'-DDT	1	5	ng/g	EPA 8270C
4,4'-DDD	1	5	ng/g	EPA 8270C
4,4'-DDE	1	5	ng/g	EPA 8270C
4,4'-DDT	1	5	ng/g	EPA 8270C
Aldrin	1	5	ng/g	EPA 8270C
BHC-alpha	1	5	ng/g	EPA 8270C
BHC-beta	1	5	ng/g	EPA 8270C
BHC-delta	1	5	ng/g	EPA 8270C
BHC-gamma	1	5	ng/g	EPA 8270C
Chlordane-alpha	1	5	ng/g	EPA 8270C
Chlordane-gamma	1	5	ng/g	EPA 8270C
cis-Nonachlor	1	5	ng/g	EPA 8270C
Dieldrin	1	5	ng/g	EPA 8270C
Endosulfan sulfate	1	5	ng/g	EPA 8270C
Endosulfan-I	1	5	ng/g	EPA 8270C
Endosulfan-II	1	5	ng/g	EPA 8270C
Endrin	1	5	ng/g	EPA 8270C
Endrin aldehyde	1	5	ng/g	EPA 8270C
Endrin ketone	1	5	ng/g	EPA 8270C
Heptachlor	1	5	ng/g	EPA 8270C
Heptachlor epoxide	1	5	ng/g	EPA 8270C
Hexachlorobenzene	1	5	ng/g	EPA 8270C
Methoxychlor	1	5	ng/g	EPA 8270C
Mirex	1	5	ng/g	EPA 8270C
Oxychlordane	1	5	ng/g	EPA 8270C
Perthane	5	10	ng/g	EPA 8270C
trans-Nonachlor	1	5	ng/g	EPA 8270C
Polynuclear Aromatic Hydrocarbons (PAHs)				
1-Methylnaphthalene	1	5	ng/g	EPA 8270C
1-Methylphenanthrene	1	5	ng/g	EPA 8270C
2,3,5-Trimethylnaphthalene	1	5	ng/g	EPA 8270C
2,6-Dimethylnaphthalene	1	5	ng/g	EPA 8270C
2-Methylnaphthalene	1	5	ng/g	EPA 8270C
Acenaphthene	1	5	ng/g	EPA 8270C
Acenaphthylene	1	5	ng/g	EPA 8270C
Anthracene	1	5	ng/g	EPA 8270C
Benz[a]anthracene	1	5	ng/g	EPA 8270C
Benzo[a]pyrene	1	5	ng/g	EPA 8270C
Benzo[b]fluoranthene	1	5	ng/g	EPA 8270C
Benzo[e]pyrene	1	5	ng/g	EPA 8270C
Benzo[g,h,i]perylene	1	5	ng/g	EPA 8270C
Benzo[k]fluoranthene	1	5	ng/g	EPA 8270C
Biphenyl	1	5	ng/g	EPA 8270C
Chrysene	1	5	ng/g	EPA 8270C
Dibenz[a,h]anthracene	1	5	ng/g	EPA 8270C
Dibenzothiophene	1	5	ng/g	EPA 8270C
Fluoranthene	1	5	ng/g	EPA 8270C
Fluorene	1	5	ng/g	EPA 8270C
Indeno[1,2,3-c,d]pyrene	1	5	ng/g	EPA 8270C
Naphthalene	1	5	ng/g	EPA 8270C
Polynuclear Aromatic Hydrocarbons (Continued)				
Perylene	1	5	ng/g	EPA 8270C
Phenanthrene	1	5	ng/g	EPA 8270C
Pyrene	1	5	ng/g	EPA 8270C
Aroclors				
Aroclor 1016	10	20	ng/g	EPA 8270C
Aroclor 1221	10	20	ng/g	EPA 8270C
Aroclor 1232	10	20	ng/g	EPA 8270C
Aroclor 1242	10	20	ng/g	EPA 8270C
Aroclor 1248	10	20	ng/g	EPA 8270C
Aroclor 1254	10	20	ng/g	EPA 8270C
Aroclor 1260	10	20	ng/g	EPA 8270C
Polychlorinated Biphenyls (PCB's)				
PCB003	1	5	ng/g	EPA 8270C
PCB008	1	5	ng/g	EPA 8270C
PCB018	1	5	ng/g	EPA 8270C
PCB028	1	5	ng/g	EPA 8270C
PCB031	1	5	ng/g	EPA 8270C
PCB033	1	5	ng/g	EPA 8270C
PCB037	1	5	ng/g	EPA 8270C
PCB044	1	5	ng/g	EPA 8270C
PCB049	1	5	ng/g	EPA 8270C
PCB052	1	5	ng/g	EPA 8270C
PCB056(060)	1	5	ng/g	EPA 8270C
PCB066	1	5	ng/g	EPA 8270C
PCB070	1	5	ng/g	EPA 8270C
PCB074	1	5	ng/g	EPA 8270C
PCB077	1	5	ng/g	EPA 8270C
PCB081	1	5	ng/g	EPA 8270C
PCB087	1	5	ng/g	EPA 8270C
PCB095	1	5	ng/g	EPA 8270C
PCB097	1	5	ng/g	EPA 8270C
PCB099	1	5	ng/g	EPA 8270C
PCB101	1	5	ng/g	EPA 8270C
PCB105	1	5	ng/g	EPA 8270C
PCB110	1	5	ng/g	EPA 8270C
PCB114	1	5	ng/g	EPA 8270C
PCB118	1	5	ng/g	EPA 8270C
PCB119	1	5	ng/g	EPA 8270C
PCB123	1	5	ng/g	EPA 8270C
PCB126	1	5	ng/g	EPA 8270C
PCB128	1	5	ng/g	EPA 8270C
PCB138	1	5	ng/g	EPA 8270C
PCB141	1	5	ng/g	EPA 8270C
PCB149	1	5	ng/g	EPA 8270C
PCB151	1	5	ng/g	EPA 8270C
PCB153	1	5	ng/g	EPA 8270C
PCB156	1	5	ng/g	EPA 8270C
PCB157	1	5	ng/g	EPA 8270C
PCB158	1	5	ng/g	EPA 8270C
PCB167	1	5	ng/g	EPA 8270C
PCB168/132	1	5	ng/g	EPA 8270C
PCB169	1	5	ng/g	EPA 8270C
PCB170	1	5	ng/g	EPA 8270C
PCB174	1	5	ng/g	EPA 8270C
PCB177	1	5	ng/g	EPA 8270C
PCB180	1	5	ng/g	EPA 8270C
PCB183	1	5	ng/g	EPA 8270C
PCB187	1	5	ng/g	EPA 8270C
PCB189	1	5	ng/g	EPA 8270C
PCB194	1	5	ng/g	EPA 8270C
PCB195	1	5	ng/g	EPA 8270C
PCB199(200)	1	5	ng/g	EPA 8270C
PCB201	1	5	ng/g	EPA 8270C
PCB206	1	5	ng/g	EPA 8270C
PCB209	1	5	ng/g	EPA 8270C

